

Pairs of fields

A. Fornasiero

Talk on 9 June 2008, Freiburg, v. 1.2

1 Introduction

Definition. Given a language L , let L^2 be the expansion of L with a new unary predicate U . Given an L -theory T , let T^2 be the L^2 -expansion of T , whose models are given by pairs (B, A) , where $A \prec B \models T$, $A \neq B$, and U is interpreted by A .

Let B be a field, and A be an elementary substructure of B (in the language of rings $L_{Ring} := (0, 1, +, \cdot)$). What can we say about the theory of the pair (B, A) (i.e., what can we say about T_{Field}^2)?

We will focus on 2 cases: $T = T_{ACF}$, and $T = T_{RCF}$.

Let T_{ACF_p} be the theory of algebraically closed fields of characteristic p .

1.1 Theorem (Robinson '59). $T_{ACF_p}^2$ is complete (and decidable), for every $p \geq 0$.

T_{RCF}^2 is not complete: for instance, A could be dense in B , or not.

Definition. Given a theory T expanding T_{RCF} , let T^{dense} be the extension of T^2 by adding the condition that A is dense in B .

1.2 Theorem (Robinson '59, van den Dries '98). *If T is o -minimal and complete, then T^{dense} is complete. In particular, T_{RCF}^{dense} is complete.*

1.3 Theorem (Macintyre '69). *Given a prime p , let T_p be the theory of p -adically closed fields. Then, T_p^{dense} is complete.*

1.4 Theorem (Macintyre '69). *If (B_i, A_i) are pairs of Henselian valued fields of residue characteristic 0, such that $A_1 \equiv A_2 \equiv B_1 \equiv B_2$, and A_i are dense proper subfields of B_i , then $(B_1, A_1) \equiv (B_2, A_2)$.*

2 Proof of Thm. 1.1

(Modified from Keisler '64).

Given a pair of fields (B, A) , let $\dim(B/A)$ be the transcendence degree of B over A .

2.1 Lemma. *Let κ be an uncountable cardinal, and (B, A) be a κ -saturated model of T_{ACF}^2 . Then, $\dim B/A \geq \kappa$.*

Proof. Let E be a basis of B/A . If, for contradiction, $|E| < \kappa$, let $q(x)$ be the partial unary L^2 -type over E , saying for every $P \in \mathbb{Z}[x, \bar{y}, \bar{z}]$ polynomial and every \bar{e} tuple of E ,

$$\forall \bar{y} (U(\bar{y}) \wedge P(\cdot, \bar{y}, \bar{e}) \neq 0 \rightarrow P(x, \bar{y}, \bar{e}) \neq 0).$$

The type q is consistent; therefore, it has a realization $b \in B$. However, $b \notin \text{acl}(A \cup E)$, absurd. \square

2.2 Lemma. *Let κ be an uncountable cardinal, $p \geq 0$, and (B, A) and (B', A') be saturated models of $T_{ACF_p}^2$ of cardinality κ . Then, (B, A) and (B', A') are isomorphic (as L_{Ring}^2 -structures).*

Proof. It is clear that A and A' are isomorphic (as fields), because T_{ACF_p} is a complete theory; let $\phi : A \cong A'$ be such an isomorphism.

By the previous Lemma, $\dim(B/A) = \dim(B'/A') = \kappa$. Thus, there exists an isomorphism of fields $\psi : B \cong B'$ extending ϕ (by well known facts about algebraically closed fields). Therefore, (B, A) and (B', A') are isomorphic. \square

Proof of Thm. 1.1. Assume that GCH , the Generalized Continuum Hypothesis, is true. Then, for every 2 models (B, A) and (D, C) of $T_{ACF_p}^2$, there exists saturated models (B^*, A^*) and (D^*, C^*) of the same cardinality, such that $(B, A) \preceq (B^*, A^*)$, and $(D, C) \preceq (D^*, C^*)$. By the previous Lemma, $(B^*, A^*) \cong (D^*, C^*)$, and therefore $(B, A) \equiv (D, C)$, proving completeness.

If GCH is not true, notice that the statement $\Omega := "T_{ACF_p}^2 \text{ is complete}"$ is an arithmetical one (i.e., it is an assertion about natural numbers, involving only quantifiers over \mathbb{N}). Since the constructible universe L satisfies GCH , Ω is true in L . By absoluteness, Ω is true also in the "true" universe V (that is, if ψ is a sentence in L_{Ring} , then, by Gödel completeness theorem, in L we can prove either ψ or $\neg\psi$ from $T_{ACF_p}^2$; such a proof is also a proof in V). \square

3 Proof of Thm. 1.2

(From van den Dries '98)

Let T be a complete o-minimal theory, expanding a field, and having universal axioms. Given (B, A) a pair of models of T , let $\dim(B/A)$ be the o-minimal dimension of B over A .¹

3.1 Proposition. *Let $\kappa > |T|$ be an infinite cardinal, and (B, A) be a κ -saturated model of T^{dense} . Then, $\dim(B/A) \geq \kappa$.*

Proof. If $T = T_{RCF}$, the proposition follows from Lemma 2.1. The general case is postponed. \square

Definition. Let $(B, A) \subseteq (D, C) \models T^2$. We say that B and C are **free** over A if every set $Y \subseteq B$ that is independent over C remains independent over A .²

3.2 Remark. If B and C are free over A , then

1. $C \cap B = A$;
2. C and B are free over A (symmetry);
3. if $Z \subseteq C$, then $(B, A) \subseteq (B\langle Z \rangle, A\langle Z \rangle) \subseteq (D, C)$, and $B\langle Z \rangle$ and C are free over $A\langle Z \rangle$.

3.3 Remark. If $(B, A) \preceq (B^*, A^*) \models T^2$, then B and A^* are free over A .

3.4 Remark. Let $p(x)$ be a 1-type over $A \models T$. If p is not realized in A , then p is “open”: that is, for every $L(A)$ -formula contained in p and for every $b \models p$ ($b \in B \succ A$), there exists an open interval I with end-points in A , such that $b \in I \subset \phi(B)$.

Let $\kappa > |T|$ be a cardinal number, and (B^*, A^*) and (D^*, C^*) be κ -saturated models of T^{dense} . Define Γ be the set of all isomorphisms $i : (B, A) \cong (D, C)$ between substructures of (B^*, A^*) and (D^*, C^*) respectively, such that $|B| = |D| < \kappa$, B and A^* are free over A , and D and C^* are free over C .

3.5 Lemma. Γ has the back-and-forth property.

Theorem 1.2 follows easily from the above Lemma.

¹Remember that T has definable Skolem functions, and that the definable closure dcl has the Exchange Property. Thus, dcl is a pre-geometry, and \dim is the dimension induced by that pre-geometry.

²Free models are also called linearly disjoint in the case $T = T_{RCF}$.

Proof. Let $i : (B, A) \cong (D, C)$ be in Γ . Given $b^* \in B^* \setminus B$, we want to extend i to $j \in \Gamma$, such that b^* is in the domain of j .

CASE 1: $b^* := a^* \in A^* \setminus B$. Define $B' := B\langle a^* \rangle$, $A' := A\langle a^* \rangle$. Notice that B' and A^* are free over A' (and, in particular, $B' \cap A^* = A'$). Let $p(x)$ be the L -type of a^* over B , and $q(x)$ be the image of p via i . Notice that $q(x)$ is consistent, and therefore, by saturation, it has a realization $d^* \in D^* \setminus D$. Since A^* is dense in D^* , $q(x)$ has also a realization $c^* \in C^* \setminus D$. Define $D' := D\langle c^* \rangle$, $C' := C\langle c^* \rangle$. Notice that D' and C^* are free over C' . Hence, there exists a unique isomorphisms $j : (B', A') \cong (D', C')$ in Γ extending i and mapping a^* to c^* .

Let $K := A^*\langle B \rangle = B\langle A^* \rangle$, and $K' := C^*\langle D \rangle = D\langle C^* \rangle$.

CASE 2: $b^* \in K$. Then, $b \in B\langle a_1^*, \dots, a_n^* \rangle$, for some $n \in \mathbb{N}$ and $a_1^*, \dots, a_n^* \in A^*$. Apply the Case 1 construction n times to extend i .

CASE 3: $b^* \in B^* \setminus K$. Define $B' := B\langle b^* \rangle$, $A' := A$. Notice that B' and A^* are free over A' . Let $p(x)$ be the L -type of b^* over B , and $q(x)$ be the image of p via i . By Proposition 3.1, $K' \neq D^*$. Hence, by saturation, q has a realization $d^* \in D^* \setminus K'$. Define $D' := D\langle d^* \rangle$, $C' := C$. Notice that D' and C^* are free over C' . Hence, there exists a unique isomorphisms $j : (B', A') \cong (D', C')$ in Γ extending i and mapping b^* to d^* . \square

3.6 Corollary (of the proof). *Let $i \in \Gamma$, and $\bar{b} := (b_1, \dots, b_n)$ be in the domain of i . Then, \bar{b} realizes the same L^2 -type over \emptyset as $i(\bar{b})$.*

Call a L^2 -formula **special** if it is of the form $\exists \bar{y}(\psi(\bar{x}, \bar{y}) \wedge U(\bar{y}))$, where ψ is an L -formula.

3.7 Corollary. *Every L^2 -formula $\phi(\bar{x})$ is equivalent, modulo T^{dense} , to a Boolean combination of special formulae. I.e., for every L^2 -formula $\phi(\bar{x})$ there exists a Boolean combination of special formulae $\theta(\bar{x})$, such that $T^{\text{dense}} \vdash \forall \bar{x}(\phi(\bar{x}) \leftrightarrow \theta(\bar{x}))$.*

Proof. By a compactness argument, it suffices to show that, if $\bar{b} \in (B^*)^n$ and $\bar{d} \in (D^*)^n$ satisfy the same special formulae, then \bar{b} and \bar{d} realize the same L^2 -types over \emptyset .

W.l.o.g., we can assume that b_1, \dots, b_r are independent over A^* , where that $r := \dim(\bar{b}/A^*) = r$.

1 *Claim.* $\dim(\bar{d}/C^*) = r$, and d_1, \dots, d_r are independent over C^* .

In fact, if for instance, $d_r \in A^*\langle d_1, \dots, d_{r-1} \rangle$, then $d_r = f(d_1, \dots, d_{r-1}, \bar{c})$ for some \emptyset -definable function f and some $\bar{c} \in (C^*)^l$. By hypothesis, there exists $\bar{a} \in (A^*)^l$ such that $b_r = f(b_1, \dots, b_{r-1}, \bar{a})$, absurd.

Let $\bar{a} \in A^m$ be such that $\dim(\bar{b}/A) = r$, where A is the substructure of A^* generated by \bar{a} . Notice that $A\langle\bar{b}\rangle$ and A^* are free over A . Let $p(\bar{x}, \bar{y})$ be the L -type of (\bar{a}, \bar{b}) over \emptyset , $p'(\bar{x}, \bar{y})$ be the partial L^2 -type extending p with “ $U(\bar{x})$ ”, and $q(\bar{x}) := p'(\bar{x}, \bar{d})$. By hypothesis on \bar{d} , q is consistent; let $\bar{c} \in (C^*)^m$ be a realization of q , and C be the definable closure of \bar{c} .

Notice that $\dim(\bar{d}/C) = r$, and $C\langle\bar{d}\rangle$ and C^* are free over C . Thus, using the Case 3 construction r times, we obtain an isomorphism $j : (A\langle\bar{b}\rangle, A) \cong (C\langle\bar{d}\rangle, C)$ in Γ . Thus, (\bar{b}, \bar{a}) and (\bar{d}, \bar{c}) realize the same L^2 -type over \emptyset . \square

3.1 Sketch of proof of Prop. 3.1

Call a subset $X \subseteq B$ **small** if it is of the form $f(A^n)$, for some $T(B)$ -definable function f .

3.8 Lemma. *Let $(B, A) \models T^2$, $f : B^{n+1} \rightarrow B$ be A -definable in B , and $b \in B \setminus A$. Then, there exist $a_0, \dots, a_n \in A$ such that*

$$a_0 + a_1 b + \dots + a_n b^n \notin f(A^n \times \{b\}).$$

In particular, $f(A^n \times \{b\}) \neq B$.

Proof. See [1, Lemma 1.1]. \square

3.9 Lemma. *Let $(B, A) \models T^{\text{dense}}$. Then, each small subset of B is a proper subset of B .*

Proof. Let $g : B^n \rightarrow B$ be T -definable in B . We have to prove that $g(A^n) \neq B$. After increasing A , we can assume that $B = A\langle b \rangle$, for some $b \in B \setminus A$. Hence, $g(xv)$ is of the form $f(\bar{x}, b)$, for some A -definable function f . The conclusion follows from the previous Lemma. \square

We can now prove Prop. 3.1 in the same way as Lemma 2.1. In fact, let E be a basis of B/A , and assume, for contradiction, that $|E| < \kappa$. Let $p(x)$ be the partial unary L^2 -type over E , saying, for every function $f(\bar{y}, \bar{z})$ T -definable over \emptyset , and every \bar{e} tuple of E ,

$$\forall \bar{y} (U(\bar{y}) \rightarrow x \neq f(\bar{y}, \bar{e})).$$

By the previous Lemma, p is consistent (because $f(A^n, \bar{e})$ is small, and a finite union of small sets is small); therefore, by saturation, it has a realization $b \in B$. However, $b \notin A\langle E \rangle$, absurd.

References

- [1] van den Dries, Lou. *Dense pairs of o-minimal structures*. Fund. Math. 157 (1998), no. 1, 61–78.
- [2] Keisler, H. Jerome. *Complete theories of algebraically closed fields with distinguished subfields*. Michigan Math. J. 11 (1964), 71–81.
- [3] Macintyre, Angus. *Dense embeddings I: A theorem of Robinson in a general setting*. Model theory and algebra (A memorial tribute to Abraham Robinson), pp. 200–219. Lecture Notes in Math., Vol. 498, Springer, Berlin, 1975.
- [4] Robinson, A. *Solution of a problem of Tarski*. Fund. Math. 47 (1959), 179–204.