

A general correlation inequality and the Almost Sure Local Limit Theorem for random sequences in the domain of attraction of a stable law ^{*}

Rita Giuliano[†]

Zbigniew S. Szewczak[‡]

Abstract

In the present paper we obtain a new correlation inequality and use it for the purpose of extending the theory of the Almost Sure Local Limit Theorem to the case of lattice random sequences in the domain of attraction of a stable law. In particular, we prove ASLLT in the case of the normal domain of attraction of α -stable law, $\alpha \in (1, 2)$.

Keywords: Almost Sure Local Limit Theorem, domain of attraction, stable law, characteristic function, correlation inequality.

2010 Mathematical Subject Classification: Primary 60F15, Secondaries 60E07, 62H20.

1 Introduction

In the recent paper [10], the author proves a correlation inequality and an Almost Sure Local Limit Theorem (ASLLT) for i.i.d. square integrable random variables taking values in a lattice. The sequence of partial sums of such variables are of course in the domain of attraction of the normal law, which is stable of order $\alpha = 2$.

The aim of the present paper is to give an analogous correlation inequality (Theorem 3.1) for the more general case of random sequences in the domain of attraction of a stable law of order $\alpha \leq 2$ and to apply it for the purpose of extending the theory of ASLLT. Notice that in our situation the summands need not be square integrable. Our correlation inequality turns out to be of the typical form needed in the theory of Almost Sure (Central and Local) Limit Theorems (see Corollary 3.3 and Remark 3.4). Our work is based on a careful use of the form of the characteristic function, and is completely different from the one used in [10] (Mc Donald's method of extraction of the Bernoulli part of a random variable).

Acknowledgement. We are grateful to an anonymous referee whose suggestions have led to a substantial improvement of the presentation.

2 The assumptions and some preliminaries

In this paper we shall be concerned with a sequence of i.i.d. random variables $(X_n)_{n \geq 1}$ such that their common distribution F is in the domain of attraction of G , where G is a stable distribution

^{*}The financial supports of the Research Grant PRIN 2008 *Probability and Finance*, of the INDAM-GNAMPA and of the Polish National Science Centre, Grant N N201 608740 are gratefully acknowledged.

[†]Address: Dipartimento di Matematica "L. Tonelli", Università di Pisa, Largo Bruno Pontecorvo 5, I-56127 Pisa, Italy. e-mail: giuliano@dm.unipi.it

[‡]Address: Nicolaus Copernicus University, Faculty of Mathematics and Computer Science, ul. Chopina 12/18 87-100 Toruń, Poland. e-mail: zssz@mat.uni.torun.pl

with exponent α ($0 < \alpha \leq 2$, $\alpha \neq 1$). This means that, for a suitable choice of constants a_n and b_n , the distribution of

$$T_n := \frac{X_1 + \cdots + X_n - a_n}{b_n}$$

converges weakly to G . It is well known (see [6], p. 46) that in such a case we have $b_n = L(n)n^{1/\alpha}$, where L is slowly varying in Karamata's sense. For $\alpha > 1$ we shall assume that X_1 is centered; by Remark 2 p. 402 of [1], this implies that $a_n = 0$, for every α .

We shall suppose that X_1 takes values in the lattice $\mathcal{L}(a, d) = \{a + kd, k \in \mathbb{Z}\}$ where d is the maximal span of the distribution; hence $S_n := X_1 + \cdots + X_n$ takes values in the lattice $\mathcal{L}(na, d) = \{na + kd, k \in \mathbb{Z}\}$.

For every n , let κ_n be a number of the form $na + kd$ and let

$$\lim_{n \rightarrow \infty} \frac{\kappa_n}{b_n} = \kappa.$$

Observe that Theorem 4.2.1 p. 121 in [6] implies that

$$\sup_n \left\{ \sup_k b_n P(S_n = k) \right\} = C < \infty. \quad (1)$$

Throughout this paper we assume that

$$x^\alpha P(X > x) = (c_1 + o(1))l(x); \quad x^\alpha P(X \leq -x) = (c_2 + o(1))l(x), \quad \alpha \in (0, 2], \quad (2)$$

where l is slowly varying as $x \rightarrow \infty$ and c_1 and c_2 are two suitable non-negative constants, $c_1 + c_2 > 0$, related to the stable distribution G .

Let ϕ be the characteristic function of F . By [1], Theorem 1, for $\alpha \neq 1$ it has the form

$$\phi(t) = \exp \left\{ -c|t|^\alpha h\left(\frac{1}{|t|}\right) \left(1 - i\beta \text{sign}(t) \tan \frac{\pi\alpha}{2}\right) + o\left(|t|^\alpha h\left(\frac{1}{|t|}\right)\right) \right\}, \quad (3)$$

where $c = \Gamma(1 - \alpha)(c_1 + c_2) \cos \frac{\pi\alpha}{2} > 0$ and $\beta = \frac{c_1 - c_2}{c_1 + c_2} \in [-1, 1]$ are two constants and $h(x) = l(x)$ if $\alpha \in (0, 2)$ and $c = \frac{1}{2}$, $\beta = 0$, $h(x) = E[X^2 1_{\{|X| \leq x\}}]$ if $\alpha = 2$. This formula implies that

$$\begin{aligned} \log |\phi(t)| &= \Re(\log \phi(t)) = -c|t|^\alpha h\left(\frac{1}{|t|}\right) (1 + o(1)), \\ \arg(\phi(t)) &= \Im(\log \phi(t)) = -c|t|^\alpha h\left(\frac{1}{|t|}\right) \left(-\beta \text{sign}(t) \tan \frac{\pi\alpha}{2} + o(1)\right) \end{aligned} \quad (4)$$

hence

$$\lim_{t \rightarrow 0} \left| \frac{\arg(\phi(t))}{\log |\phi(t)|} \right| = \left| \beta \tan \frac{\pi\alpha}{2} \right|. \quad (5)$$

We notice that $L(n) = h^{\frac{1}{\alpha}}(b_n)$ for $\alpha \in (0, 2)$ (by Remark 2 p. 402 in [1]), while $L(n) = \sqrt{E[X^2 1_{\{|X| \leq b_n\}}]}$ for $\alpha = 2$.

Remark 2.1 For the case $\alpha = 2$ we need that $x \mapsto x^2 P(|X| > x)$ is a slowly varying function, a stronger assumption than the slow variation of $x \mapsto E[X^2 1_{\{|X| \leq x\}}]$ (which in turn is equivalent to the CLT, see Corollary 1 p. 578 in [3]). To see this, consider the following distribution:

$$P(X = n) = \frac{C}{n^2 2^n}, \quad n \geq 1, \quad C = \sum_{k \geq 1} \frac{1}{k^2 2^k}.$$

It is easy to check that in this case $x \mapsto x^2 P(|X| > x)$ is not slowly varying.

Remark 2.2 Let $\tilde{h} \sim h$ as $x \rightarrow +\infty$. Then, by (4),

$$\log |\phi(t)| = -c|t|^\alpha h\left(\frac{1}{|t|}\right)(1 + o(1)) = -c|t|^\alpha \tilde{h}\left(\frac{1}{|t|}\right) \cdot \frac{h\left(\frac{1}{|t|}\right)}{\tilde{h}\left(\frac{1}{|t|}\right)}(1 + o(1)) = -c|t|^\alpha \tilde{h}\left(\frac{1}{|t|}\right)(1 + o(1)).$$

This means that h is unique up to equivalence; thus, by Theorem 1.3.3. p. 14 of [2] we can assume that h is continuous (even C^∞) on $[a, \infty)$ for some $a > 0$.

An analogous observation is in force for $\arg(\phi(t))$.

Remark 2.3 Thus we deal with a subclass of strictly stable distributions. Denoting by ψ the characteristic function of G , we know from [11], Theorem C.4 on p.17 that $\log \psi$ (for strictly stable distributions) has the form

$$\log \psi(t) = -c|t|^\alpha \exp\left\{-i \left(\frac{\pi}{2}\right) \theta \alpha \operatorname{sign}(t)\right\},$$

where $|\theta| \leq \min\{1, \frac{2}{\alpha} - 1\}$ and $c > 0$. For $\alpha = 1$ and $|\theta| = 1$ we get degenerate distribution and in this case we say that X_n is relatively stable (see e.g. [9]). Almost sure variant of relative stability for dependent strictly stationary sequences will be discussed elsewhere.

Let $\delta > -1$ and $p > 0$ two given numbers; we shall use the equality

$$\int_0^{+\infty} t^\delta e^{-pt^\alpha} dt = \frac{\Gamma(\frac{\delta+1}{\alpha})}{\alpha} \cdot \frac{1}{p^{\frac{\delta+1}{\alpha}}} = C \cdot \frac{1}{p^{\frac{\delta+1}{\alpha}}}. \quad (6)$$

In what follows, with the symbols C , c and so on we shall mean positive constants the value of which may change from case to case.

3 The correlation inequality

We assume that $(X_n)_{n \geq 1}$ is a sequence of i.i.d. random variables verifying the following conditions: (2), $\alpha \neq 1$ and $\mu = E[X_1] = 0$ when $\alpha > 1$. Recall that the norming constants are $a_n = 0$ and $b_n = L(n)n^{1/\alpha}$ with L slowly varying. With no loss of generality, we shall assume throughout that $d = 1$.

Theorem 3.1 (i) *In the above setting we have*

$$\begin{aligned} & b_m b_n \left| P(S_m = \kappa_m, S_n = \kappa_n) - P(S_m = \kappa_m)P(S_n = \kappa_n) \right| \\ & \leq C \left\{ \left(\frac{n}{n-m} \right)^{1/\alpha} \frac{L(n)}{L(n-m)} + 1 \right\}. \end{aligned}$$

(ii) *In addition to the previous hypotheses assume that the function h appearing in (2) and (3) verifies*

$$\liminf_{x \rightarrow \infty} h(x) =: \ell > 0.$$

Then there exists $\epsilon > 0$ such that, putting

$$M(x) = \sup_{\frac{1}{\epsilon} \leq y \leq x} h(y), \quad x \geq \frac{1}{\epsilon},$$

we have $M(x) < \infty$ for every x and

$$\begin{aligned} & b_m b_n \left| P(S_m = \kappa_m, S_n = \kappa_n) - P(S_m = \kappa_m)P(S_n = \kappa_n) \right| \\ & \leq CL(n) \left\{ n^{1/\alpha} \left(\frac{1}{e^{(n-m)c}} + \frac{1}{e^{nc}} \right) + \frac{\frac{m}{n}}{\left(1 - \frac{m}{n}\right)^{1+\frac{1}{\alpha}}} \left(1 + M(n^{1+\frac{1}{\alpha}})\right) + \frac{\left(\frac{m}{n}\right)^{\frac{\eta}{\alpha}} L^\eta(m)}{\left(1 - \frac{m}{n}\right)^{\frac{\eta+1}{\alpha}}} \right\} \quad (7) \end{aligned}$$

for every pair (m, n) of integers, with $m \geq 1$, $n > m + \epsilon^{-\frac{\alpha}{\alpha+1}}$, and for every $\eta \in (0, 1]$.

Remark 3.2 If h is ultimately increasing, then condition (ii) of Theorem 3.1 is automatically satisfied. A quick look at the proof (see below) shows that if h is increasing and continuous, the inequality (7) holds for $1 \leq m < n$.

Proof of Theorem 3.1.

(i) We write

$$\begin{aligned} & b_m b_n \left| P(S_m = \kappa_m, S_n = \kappa_n) - P(S_m = \kappa_m)P(S_n = \kappa_n) \right| \\ & = \left\{ b_m P(S_m = \kappa_m) \right\} \cdot \left\{ b_n \left| P(S_{n-m} = \kappa_n - \kappa_m) - P(S_n = \kappa_n) \right| \right\} \\ & \leq C \cdot b_n \left| P(S_{n-m} = \kappa_n - \kappa_m) - P(S_n = \kappa_n) \right|, \end{aligned}$$

by (1). Inequality (i) follows since

$$\begin{aligned} & b_n \left| P(S_{n-m} = \kappa_n - \kappa_m) - P(S_n = \kappa_n) \right| \leq b_n \left(P(S_{n-m} = \kappa_n - \kappa_m) + P(S_n = \kappa_n) \right) \\ & = \left(\frac{b_n}{b_{n-m}} \cdot b_{n-m} P(S_{n-m} = \kappa_n - \kappa_m) + b_n P(S_n = \kappa_n) \right) \leq C \left(\frac{b_n}{b_{n-m}} + 1 \right), \end{aligned}$$

by (1) again.

(ii) Let ϕ be the characteristic function of F . By the inversion formula (see Theorem 4, p. 511 of [3]) we can write

$$\begin{aligned} & b_n \left| P(S_{n-m} = \kappa_n - \kappa_m) - P(S_n = \kappa_n) \right| = \frac{b_n}{2\pi} \left| \int_{-\pi}^{\pi} \left\{ e^{-it(\kappa_n - \kappa_m)} \phi^{n-m}(t) - e^{-it\kappa_n} \phi^n(t) \right\} dt \right| \\ & \leq C b_n \int_{-\pi}^{\pi} \left| e^{it\kappa_m} \phi^{n-m}(t) - \phi^n(t) \right| dt. \end{aligned}$$

Recall the expression (3) of ϕ where, by Remark 2.2, one can choose h continuous on $[a, \infty)$ for some $a > 0$. The additional assumption $\ell > 0$ allows to take $\epsilon \in (0, \frac{1}{a}]$ such that, for $|t| < \epsilon$ we have

$$h\left(\frac{1}{|t|}\right) > \frac{A}{2} > 0. \quad (8)$$

We write

$$\int_{-\pi}^{\pi} \left| e^{it\kappa_m} \phi^{n-m}(t) - \phi^n(t) \right| dt = \int_{|t| < \epsilon} + \int_{\epsilon < |t| < \pi} = I_1 + I_2.$$

Now

$$I_2 \leq \int_{\epsilon < |t| < \pi} |\phi(t)|^{n-m} dt + \int_{\epsilon < |t| < \pi} |\phi(t)|^n dt.$$

Since $d = 1$, by Theorem 1.4.2 p. 27 of [6] we have $|\phi(t)| < 1$ for $0 < |t| < 2\pi$. Hence a constant $c > 0$ exists such that, for $\epsilon < |t| < \pi$ we have $|\phi(t)| < e^{-c}$, which gives

$$\int_{\epsilon < |t| < \pi} |\phi(t)|^n dt < 2\pi e^{-nc}; \quad \int_{\epsilon < |t| < \pi} |\phi(t)|^{n-m} dt < 2\pi e^{-(n-m)c},$$

so that

$$I_2 \leq C \left(e^{-(n-m)c} + e^{-nc} \right). \quad (9)$$

Now we evaluate I_1 ,

$$\begin{aligned} I_1 &= \int_{-\epsilon}^{\epsilon} \left| e^{it\kappa_m} \phi^{n-m}(t) - \phi^n(t) \right| dt \\ &\leq \int_{-\epsilon}^{\epsilon} \left| e^{it\kappa_m} \phi^{n-m}(t) - e^{it\kappa_m} \phi^n(t) \right| dt + \int_{-\epsilon}^{\epsilon} \left| e^{it\kappa_m} \phi^n(t) - \phi^n(t) \right| dt \\ &= \int_{-\epsilon}^{\epsilon} \left| \phi^{n-m}(t) - \phi^n(t) \right| dt + \int_{-\epsilon}^{\epsilon} \left| e^{it\kappa_m} - 1 \right| \cdot |\phi(t)|^n dt. \end{aligned} \quad (10)$$

For a complex number $A = \rho e^{i\theta}$ we have

$$\begin{aligned} |A^x - A^y| &= \left\{ (\rho^x - \rho^y)^2 + 2\rho^{x+y}(1 - \cos \theta(x-y)) \right\}^{1/2} \leq |\rho^x - \rho^y| + \left\{ 2\rho^{x+y}(1 - \cos \theta(x-y)) \right\}^{1/2} \\ &\leq |\rho^x - \rho^y| + \rho^{\frac{x+y}{2}} |\theta| |x-y|. \end{aligned}$$

Applying with $A = \phi(t)$, $x = n - m$ and $y = n$ we get

$$\int_{|t| < \epsilon} \left| \phi^{n-m}(t) - \phi^n(t) \right| dt \leq C \left(\int_{|t| < \epsilon} \left| |\phi(t)|^{n-m} - |\phi(t)|^n \right| dt + m \int_{|t| < \epsilon} \left| \arg \phi(t) \right| \cdot |\phi(t)|^{n-(m/2)} dt \right). \quad (11)$$

Applying Lagrange Theorem to the first summand we find that for a suitable $\xi \in (n - m, n)$ we have, for every $\delta < \epsilon$

$$\begin{aligned} \int_{|t| < \epsilon} \left| |\phi(t)|^{n-m} - |\phi(t)|^n \right| dt &\leq \int_{|t| < \delta} 2 dt + m \int_{\delta < |t| < \epsilon} \left| \frac{d}{dx} \{ |\phi(t)|^x \} \right|_{x=\xi} dt \\ &= 4\delta + m \int_{\delta < |t| < \epsilon} |\log |\phi(t)|| \cdot |\phi(t)|^\xi dt \leq 4\delta + m \int_{\delta < |t| < \epsilon} |\log |\phi(t)|| \cdot |\phi(t)|^{(n-m)} dt \\ &\leq 4\delta + C_1 m \int_{\delta < |t| < \epsilon} |t|^{\alpha} h\left(\frac{1}{|t|}\right) \cdot e^{-C_2(n-m)|t|^{\alpha} h\left(\frac{1}{|t|}\right)} dt, \end{aligned} \quad (12)$$

using the relation (4). By reporting the inequality (8) into (12), and recalling that h is continuous, hence bounded on $[\delta, \epsilon]$, we obtain

$$\begin{aligned} \int_{|t|<\epsilon} \left| |\phi(t)|^{n-m} - |\phi(t)|^n \right| dt &\leq 4\delta + C_1 m M\left(\frac{1}{\delta}\right) \int_{\delta < |t| < \epsilon} |t|^\alpha \cdot e^{-C_2(n-m)|t|^\alpha} dt \\ &\leq C \left\{ \delta + \frac{m}{(n-m)^{1+\frac{1}{\alpha}}} \cdot M\left(\frac{1}{\delta}\right) \right\}, \end{aligned}$$

by (6), for any $\delta < \epsilon$. Taking $\delta = \frac{1}{(n-m)^{1+\frac{1}{\alpha}}}$, we get

$$\begin{aligned} \int_{|t|<\epsilon} \left| |\phi(t)|^{n-m} - |\phi(t)|^n \right| dt &\leq C \left\{ \frac{1}{(n-m)^{1+\frac{1}{\alpha}}} + \frac{m}{(n-m)^{1+\frac{1}{\alpha}}} \cdot M\left((n-m)^{1+\frac{1}{\alpha}}\right) \right\} \\ &\leq C \frac{m}{(n-m)^{1+\frac{1}{\alpha}}} \left(1 + M\left(n^{1+\frac{1}{\alpha}}\right) \right), \end{aligned} \quad (13)$$

M being non-decreasing.

For the second summand in (11) we can proceed as follows: by (5),

$$|\arg \phi(t)| \leq C |\log |\phi(t)||, \quad \forall t.$$

Hence, arguing as before

$$\begin{aligned} m \int_{|t|<\epsilon} |\arg \phi(t)| \cdot |\phi(t)|^{n-(m/2)} dt &\leq C m \int_{|t|<\epsilon} |\log \phi(t)| \cdot |\phi(t)|^{n-(m/2)} dt \\ &\leq C \frac{m}{(n-\frac{m}{2})^{1+\frac{1}{\alpha}}} \left\{ 1 + M\left((n-\frac{m}{2})^{1+\frac{1}{\alpha}}\right) \right\} \leq C \frac{m}{(n-m)^{1+\frac{1}{\alpha}}} \left(1 + M\left(n^{1+\frac{1}{\alpha}}\right) \right), \end{aligned} \quad (14)$$

as before. Thus, by (11) (13), (14) we obtain

$$\int_{-\epsilon}^{\epsilon} \left| \phi^{n-m}(t) - \phi^n(t) \right| dt \leq C \frac{m}{(n-m)^{1+\frac{1}{\alpha}}} \left(1 + M\left(n^{1+\frac{1}{\alpha}}\right) \right). \quad (15)$$

Let's turn to the second summand in (10). By the well known inequality

$$|e^{it} - 1| \leq 2^{1-\eta} |t|^\eta, \quad \forall \eta \in (0, 1]$$

(see [7], p. 200), we have

$$\int_{-\epsilon}^{\epsilon} \left| e^{it\kappa_m} - 1 \right| \cdot |\phi(t)|^n dt \leq |\kappa_m|^\eta 2^{1-\eta} \int_{-\epsilon}^{\epsilon} |t|^\eta \cdot |\phi(t)|^{n-m} dt \leq C \frac{|\kappa_m|^\eta}{(n-m)^{\frac{\eta+1}{\alpha}}} \leq C \frac{m^{\frac{\eta}{\alpha}} L^\eta(m)}{(n-m)^{\frac{\eta+1}{\alpha}}}, \quad (16)$$

again by (6) and the fact that

$$\kappa_m \sim \kappa b_m = \kappa L(m) m^{1/\alpha}, \quad m \rightarrow \infty.$$

Summing the estimates (9), (15) and (16) we get, for every $\eta \in (0, 1]$

$$\begin{aligned} \int_{-\pi}^{-\pi} \left| e^{it\kappa_m} \phi^{n-m}(t) - \phi^n(t) \right| dt \\ \leq C \left\{ \left(e^{-(n-m)c} + e^{-nc} \right) + \frac{m}{(n-m)^{1+\frac{1}{\alpha}}} \left(1 + M\left(n^{1+\frac{1}{\alpha}}\right) \right) + \frac{m^{\frac{\eta}{\alpha}} L^\eta(m)}{(n-m)^{\frac{\eta+1}{\alpha}}} \right\}. \end{aligned}$$

Multiplying by $b_n = L(n)n^{1/\alpha}$ gives the conclusion. \square

Corollary 3.3 For large m and $n \geq 2m$, for every $\delta < \frac{1}{\alpha}$ and for every $\eta \in (0, 1]$ we have

$$b_m b_n \left| P(S_m = \kappa_m, S_n = \kappa_n) - P(S_m = \kappa_m)P(S_n = \kappa_n) \right| \leq C \tilde{L}(n) \cdot \left(\frac{m}{n}\right)^\rho,$$

with $\tilde{L}(n) = L(n)(1 + M(n^{1+\frac{1}{\alpha}}) + L^\eta(n))$ and $\rho := \min\{\eta(\frac{1}{\alpha} - \delta), 1\}$.

Proof of Corollary 3.3. Let ϵ be the number identified in (ii) of Theorem 3.1 and c the constant appearing in the second member of (7); let $x_0 > \epsilon^{-\frac{\alpha}{\alpha+1}}$ be such that $e^{cx} \geq x^{2/\alpha}$ for $x \geq x_0$. For $m \geq x_0$ we have also $n - m \geq x_0 \geq \epsilon^{-\frac{\alpha}{\alpha+1}}$ (since $n - m \geq m$). Then (ii) of Theorem 3.1 holds and

$$\frac{n^{1/\alpha}}{e^{nc}} \leq \frac{n^{1/\alpha}}{e^{(n-m)c}} \leq \frac{n^{1/\alpha} m^{1/\alpha}}{(n-m)^{2/\alpha}} = \frac{\left(\frac{m}{n}\right)^{1/\alpha}}{\left(1 - \frac{m}{n}\right)^{2/\alpha}} \leq 2^{2/\alpha} \cdot \left(\frac{m}{n}\right)^{1/\alpha}. \quad (17)$$

Moreover

$$\frac{\frac{m}{n}}{\left(1 - \frac{m}{n}\right)^{1+1/\alpha}} \leq \frac{\frac{m}{n}}{\left(\frac{1}{2}\right)^{1+1/\alpha}} = (2^{1+\frac{1}{\alpha}}) \left(\frac{m}{n}\right); \quad (18)$$

similarly

$$\frac{\left(\frac{m}{n}\right)^{\frac{\eta}{\alpha}} L^\eta(m)}{\left(1 - \frac{m}{n}\right)^{\frac{\eta+1}{\alpha}}} \leq 2^{\frac{\eta+1}{\alpha}} L^\eta(m) \left(\frac{m}{n}\right)^{\frac{\eta}{\alpha}}. \quad (19)$$

Recall the well known representation of slowly varying functions (see for instance [2], p.12):

$$L(x) = \gamma(x) \exp \left\{ \int_1^x \frac{\varepsilon(t)}{t} dt \right\},$$

where $\gamma(x) \rightarrow \gamma$ (a finite constant) and $\varepsilon(x) \rightarrow 0$ as $x \rightarrow \infty$.

We deduce from it that, for every $\delta > 0$, $n \geq 2m$ and large m we have

$$\begin{aligned} \frac{m^\delta L(m)}{n^\delta L(n)} &\leq C \exp \left\{ \delta \log m + \int_1^m \frac{\varepsilon(t)}{t} dt - \delta \log n - \int_1^n \frac{\varepsilon(t)}{t} dt \right\} = C \exp \left\{ \delta \log \frac{m}{n} + \int_n^m \frac{\varepsilon(t)}{t} dt \right\} \\ &= C \exp \left\{ \delta \log \frac{m}{n} - \int_m^n \frac{\varepsilon(t)}{t} dt \right\} \leq C \exp \left\{ \delta \log \frac{m}{n} + \int_m^n \frac{\delta}{2t} dt \right\} = C \exp \left\{ \delta \log \frac{m}{n} - \frac{\delta}{2} \log \frac{m}{n} \right\} \\ &= C \left(\frac{m}{n}\right)^{\frac{\delta}{2}} \leq C \left(\frac{1}{2}\right)^{\frac{\delta}{2}}. \end{aligned}$$

It follows that

$$L^\eta(m) \left(\frac{m}{n}\right)^{\frac{\eta}{\alpha}} \leq C L^\eta(n) \left(\frac{m}{n}\right)^{\eta(\frac{1}{\alpha} - \delta)}. \quad (20)$$

From (19) and (20) we obtain

$$\frac{\left(\frac{m}{n}\right)^{\frac{\eta}{\alpha}} L^\eta(m)}{\left(1 - \frac{m}{n}\right)^{\frac{\eta+1}{\alpha}}} \leq C L^\eta(n) \left(\frac{m}{n}\right)^{\eta(\frac{1}{\alpha} - \delta)}. \quad (21)$$

Now the desired conclusion follows from (17), (18) and (21) and the inequality in (ii) of Proposition 3.1. □

Remark 3.4 Let $(Y_n)_{n \geq 1}$ be i.i.d. centered random variables with second moments. It is well known that correlation inequalities of the form

$$\left| \text{Cov}(Y_m, Y_n) \right| \leq C \left(\frac{m}{n} \right)^\rho \quad (22)$$

for some positive constant ρ are useful tools in order to prove Almost Sure Theorems with logarithmic weights, i.e. statements of the form

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{Y_n}{n} = 0. \quad (23)$$

See for instance [4], Theorem (2.9) as a reference.

The correlation inequality of Corollary 3.3 is similar to (22), but notice that the coefficient \tilde{L} need not be bounded. See also Remark 4.3.

Remark 3.5 In the case $\alpha = 2$ the correlation inequality of Corollary 3.3 furnishes $\rho = \eta(\frac{1}{2} - \epsilon) < \frac{1}{2}$, while in [10] the better exponent $\rho = \frac{1}{2}$ is found. Nevertheless, we are able to prove an Almost Sure Local Theorem even with the weaker exponent, as we shall see in the next section.

4 Application to the Almost Sure Local Limit Theorem

In this section we apply our main result to prove a suitable form of the Almost Sure Local Limit Theorem. Denote by g the α -stable density function related to the distribution function G . We point out that here we consider only the case $\alpha > 1$. Precisely

Theorem 4.1 *Let $(X_n)_{n \geq 1}$ be a centered, independent and identically lattice distributed (i.i.l.d.) random sequence with span $d = 1$; assume moreover that (2) holds with $\alpha \in (1, 2]$ and that there exists $\gamma \in (0, 2)$ such that*

$$\sum_{k=a}^b \frac{L(k) \{1 + M(k^{1+\frac{1}{\alpha}}) + L^\eta(k)\}}{k} \leq C(\log^\gamma b - \log^\gamma a),$$

for some $\eta \in (0, 1]$. If the condition (ii) of Theorem 3.1 is satisfied, then

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{b_n}{n} 1_{\{S_n = \kappa_n\}} = g(\kappa).$$

Example 4.2 Let $h(x) = \log^\sigma x$, with $0 < \sigma < \frac{\alpha}{1+\alpha}$. Notice that

$$\frac{\sigma}{\alpha} < \sigma \wedge (1 - \sigma) \quad (24)$$

Remark 2 p. 402 in [1] assures that

$$b_n^\alpha = n \log^\sigma b_n.$$

Putting $f(x) = \frac{x^\alpha}{\log^\sigma x}$ and observing that f is strictly increasing for $x > e^{\frac{\sigma}{\alpha}}$, this means that

$$L(n) = \frac{b_n}{n^{\frac{1}{\alpha}}} = \frac{f^{-1}(n)}{n^{\frac{1}{\alpha}}}, \quad (25)$$

for sufficiently large n . It is not difficult to check that for sufficiently large n

$$L(n) \leq \log^\delta n, \quad \forall \delta > \frac{\sigma}{\alpha}.$$

In fact by (25) this is equivalent to

$$n \leq f(n^{\frac{1}{\alpha}} \cdot \log^\delta n) = \frac{n \log^{\alpha\delta} n}{\left(\frac{1}{\alpha} \log n + \delta \log \log n\right)^\sigma},$$

which clearly holds for $\alpha\delta > \sigma$. Thus

$$L(n)\{1 + M(n^{1+\frac{1}{\alpha}}) + L^\eta(n)\} \leq \log^\delta n \left\{1 + \left(1 + \frac{1}{\alpha}\right)^\sigma \log^\sigma n + \log^{\delta\eta} n\right\} \leq C(\log n)^{2\delta \vee (\delta + \sigma)}.$$

If $\delta < \sigma$ we have $2\delta \vee (\delta + \sigma) = \delta + \sigma$, hence

$$\sum_{k=a}^b \frac{L(k)\{1 + M(k^{1+\frac{1}{\alpha}}) + L^\eta(k)\}}{k} \leq C \sum_{k=a}^b \frac{(\log k)^{\delta + \sigma}}{k} \leq C(\log^\gamma b - \log^\gamma a)$$

with $\gamma = \delta + \sigma + 1 < 2$ for any $\delta \in \left(\frac{\sigma}{\alpha}, \sigma \wedge (1 - \sigma)\right)$ (see (24)).

Remark 4.3 Let $Y_n = b_n \left(1_{\{S_n = \kappa_n\}} - P(S_n = \kappa_n)\right)$. Theorem 4.1 states exactly relation (23): just observe that

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{b_n P(S_n = \kappa_n)}{n} = g(\kappa),$$

by Theorem 4.2.1 p. 121 of [6]. Of course, our statement requires an auxiliary hypothesis, due to the fact that in the second member of our correlation inequality we have a supplementary factor, $\tilde{L}(n)$, which need not be bounded, as observed before (Remark 3.4).

Remark 4.4 If $L \equiv$ a constant (i.e. F belongs to the domain of *normal* attraction of G , according to the definition on p. 92 of [6]), then the assumption in Theorem 4.1 and condition (ii) of Theorem 3.1 are automatically satisfied; thus we get the following nice result

Corollary 4.5 *If $(X_n)_{n \geq 1}$ is a centered i.i.l.d. random sequence with span $d = 1$, (2) holds with $\alpha \in (1, 2)$ and $L \equiv c$, then*

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{c}{n^{1-\frac{1}{\alpha}}} 1_{\{S_n = \kappa_n\}} = g(\kappa).$$

Proof of Theorem 4.1. We shall denote $\tilde{L}(n) := L(n)\{1 + M(n^{1+\frac{1}{\alpha}}) + L^\eta(n)\}$ as in Corollary 3.3 of the previous section. By Ex. 1.11.4 p. 58 of [2], M is slowly varying, hence the same happens for $n \mapsto M(n^{1+\frac{1}{\alpha}})$ and for \tilde{L} . Put

$$Z_n := \sum_{k=2^{n-1}}^{2^n-1} \frac{Y_k}{k},$$

where Y_n is defined in Remark (4.3). Following the same argument as in [5], we must prove that

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n Z_i}{n} = 0.$$

We shall use the Gaal–Koksma Strong Law of Large Numbers, i.e. (see [8], p. 134); here is the precise statement:

Theorem 4.6 Let $(Z_n)_{n \geq 1}$ be a sequence of centered random variables with finite variance. Suppose that there exists a constant $\beta > 0$ such that, for all integers $m \geq 0, n > 0$,

$$\mathbf{E} \left[\left(\sum_{i=m+1}^{m+n} Z_i \right)^2 \right] \leq C((m+n)^\beta - m^\beta), \quad (26)$$

for a suitable constant C independent of m and n . Then, for each $\delta > 0$,

$$\sum_{i=1}^n Z_i = O(n^{\beta/2}(\log n)^{2+\delta}), \quad P - a.s.$$

Remark 4.7 It is easy to see that Theorem 4.6 is in force even if the bound (26) holds only for all integers $m \geq h_0, n > 0$, where h_0 is an integer strictly greater than 0: just take $Z_i = 0$ for $i = 1, 2, \dots, h_0$ and use Theorem 4.6.

We go back to the proof of Theorem 4.1, where we shall repeatedly use Remark 4.7 without mentioning it. Since

$$\mathbf{E} \left[\left(\sum_{i=m+1}^{m+n} Z_i \right)^2 \right] = \sum_{i=m+1}^{m+n} \mathbf{E}[Z_i^2] + 2 \sum_{m+1 \leq i < j \leq m+n} \mathbf{E}[Z_i Z_j], \quad (27)$$

we bound separately these two summands. We have first

$$\mathbf{E}[Z_i^2] = \sum_{h,k=2^{i-1}}^{2^i-1} \frac{1}{hk} \mathbf{E}[Y_h Y_k] = \sum_{h=2^{i-1}}^{2^i-1} \frac{1}{h^2} \mathbf{E}[Y_h^2] + 2 \sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk} \mathbf{E}[Y_h Y_k]. \quad (28)$$

Now, by (1)

$$\mathbf{E}[Y_h^2] = b_h^2 \left\{ P(S_h = \kappa_h) - P^2(S_h = \kappa_h) \right\} \leq b_h^2 P(S_h = \kappa_h) \leq C b_h = C \cdot L(h) h^{1/\alpha}.$$

Fix any $\epsilon \in (0, 1 - \frac{1}{\alpha})$ and let h_0 be such that $L(t) < t^\epsilon$ for $t \geq h_0$. Let m be such that $2^m \geq h_0$; for $h \geq 2^{i-1} \geq 2^m$ we have from the above that $\mathbf{E}[Y_h^2] \leq C h^{\epsilon+(1/\alpha)}$, which gives

$$\sum_{h=2^{i-1}}^{2^i-1} \frac{1}{h^2} \mathbf{E}[Y_h^2] \leq C \sum_{h=2^{i-1}}^{2^i-1} \frac{1}{h^{2-\epsilon-1/\alpha}} \leq C \cdot \frac{2^i - 2^{i-1}}{(2^{i-1})^{2-\epsilon-1/\alpha}} = \frac{C}{(2^{i-1})^{1-\epsilon-1/\alpha}} \leq C. \quad (29)$$

Moreover, by (i) of Theorem 3.1, we have

$$\sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk} \mathbf{E}[Y_h Y_k] \leq C \sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk} \left(\frac{k}{k-h} \right)^{1/\alpha} \frac{L(k)}{L(k-h)} + C \sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk}. \quad (30)$$

Now

$$\sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk} = \sum_{k=2^{i-1}}^{2^i-1} \frac{1}{k} \sum_{h=2^{i-1}}^{k-1} \frac{1}{h} \leq \left(\sum_{k=2^{i-1}}^{2^i-1} \frac{1}{k} \right)^2 \leq C; \quad (31)$$

and

$$\sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk} \left(\frac{k}{k-h} \right)^{1/\alpha} \frac{L(k)}{L(k-h)} \leq 2 \sum_{k=2^{i-1}}^{2^i-1} \frac{L(k)}{k^{2-1/\alpha}} \sum_{j=1}^{k-2^{i-1}} \frac{1}{j^{1/\alpha}} \frac{1}{L(j)} \quad (32)$$

(we have used the fact that $\frac{k}{2} < 2^{i-1} \leq h$).

Now we are concerned with the inner sum in the last member of (32). The function

$$t \mapsto U(t) = \frac{1}{[t]^{1/\alpha}} \frac{1}{L([t])}, \quad t \geq 1$$

is regularly varying with exponent $-\frac{1}{\alpha}$; hence, from Theorem 1 p. 281 part (b) of [3] we deduce that, for every $p \geq \frac{1}{\alpha} - 1$

$$\frac{k^{p+1}U(k)}{\int_1^k x^p U(x) dx} \rightarrow p - \frac{1}{\alpha} + 1, \quad k \rightarrow \infty.$$

Since

$$\int_1^k x^p U(x) dx = \sum_{j=2}^k \int_{j-1}^j x^p U(x) dx \geq \sum_{j=2}^k (j-1)^p \int_{j-1}^j U(x) dx = \sum_{j=1}^{k-1} j^p U(j),$$

we get

$$\liminf_{k \rightarrow \infty} \frac{k^{p+1}U(k)}{\sum_{j=1}^{k-1} j^p U(j)} \geq \lim_{k \rightarrow \infty} \frac{k^{p+1}U(k)}{\int_1^k x^p U(x) dx} = \left(p - \frac{1}{\alpha} + 1\right), \quad k \rightarrow \infty.$$

In particular, for $p = 0$ we obtain (remember that $\frac{1}{\alpha} < 1$)

$$\frac{k}{k^{1/\alpha} L(k)} = kU(k) \geq C \sum_{j=1}^{k-1} U(j) = C \sum_{j=1}^{k-1} \frac{1}{j^{1/\alpha} L(j)},$$

whence

$$\sum_{j=1}^{k-2^{i-1}} \frac{1}{j^{1/\alpha} L(j)} \leq \sum_{j=1}^{k-1} \frac{1}{j^{1/\alpha} L(j)} \leq C \frac{k}{k^{1/\alpha} L(k)},$$

and continuing (32) we obtain

$$\sum_{k=2^{i-1}}^{2^i-1} \frac{L(k)}{k^{2-1/\alpha}} \sum_{j=1}^{k-2^{i-1}} \frac{1}{j^{1/\alpha} L(j)} \leq C \sum_{k=2^{i-1}}^{2^i-1} \frac{L(k)}{k^{2-1/\alpha}} \frac{k}{k^{1/\alpha} L(k)} = C. \quad (33)$$

Summarizing, from (30), (31), (32) and (33) we have found

$$\sum_{2^{i-1} \leq h < k \leq 2^i-1} \frac{1}{hk} \mathbf{E}[Y_h Y_k] \leq C, \quad (34)$$

so that by (28), (29) and (34) we get

$$\mathbf{E}[Z_i^2] \leq C; \quad (35)$$

this implies

$$\sum_{i=m+1}^{m+n} \mathbf{E}[Z_i^2] \leq Cn \quad (36)$$

which bounds the first sum in (27). Now we consider the second one, i.e.

$$\sum_{m+1 \leq i < j \leq m+n} \mathbf{E}[Z_i Z_j].$$

We start with a bound for the summand $\mathbf{E}[Z_i Z_j]$ when $j \geq i + 2$. In this case we have

$$h \leq 2^i \leq 2^{j-2} \leq \frac{k}{2}.$$

Let m be such that $2^m > x_0$, where x_0 is as in Corollary 3.3. For $i \geq m + 1$, the same Corollary assures that

$$\mathbf{E}[Z_i Z_j] = \sum_{h=2^{i-1}}^{2^i-1} \sum_{k=2^{j-1}}^{2^j-1} \frac{1}{hk} \mathbf{E}[Y_h Y_k] \leq C \sum_{h=2^{i-1}}^{2^i-1} \frac{1}{h^{1-\rho}} \sum_{k=2^{j-1}}^{2^j-1} \frac{\tilde{L}(k)}{k^{1+\rho}}. \quad (37)$$

The function

$$V(t) = \frac{\tilde{L}([t])}{[t]^{1+\rho}}, \quad t \geq 1$$

is regularly varying with exponent $-(1 + \rho)$. Hence, by Theorem 1 p. 281 part (a) of [3], we have

$$\frac{k^{p+1}V(k)}{\int_k^\infty x^p V(x) dx} \rightarrow -p + \rho$$

if $-p + \rho \geq 0$ and $\int_k^\infty x^p V(x) dx$ is finite. In particular we can take $p = 0$, since

$$\int_1^\infty V(x) dx = \sum_{j=1}^\infty \frac{\tilde{L}(j)}{j^{1+\rho}} < +\infty,$$

and we obtain

$$\frac{\frac{\tilde{L}(k)}{k^\rho}}{\int_k^\infty V(x) dx} = \frac{kV(k)}{\int_k^\infty V(x) dx} \rightarrow \rho. \quad (38)$$

Now

$$\int_k^\infty V(x) dx = \int_k^\infty \frac{\tilde{L}([x])}{[x]^{1+\rho}} dx = \sum_{j=k}^\infty \int_j^{j+1} \frac{\tilde{L}([x])}{[x]^{1+\rho}} dx \geq \sum_{j=k}^\infty \frac{\tilde{L}(j)}{(j+1)^{1+\rho}} \geq \left(\frac{1}{2}\right)^{1+\rho} \sum_{j=k}^\infty \frac{\tilde{L}(j)}{j^{1+\rho}} \quad (39)$$

and similarly

$$\int_k^\infty V(x) dx \leq \sum_{j=k}^\infty \frac{\tilde{L}(j)}{j^{1+\rho}}. \quad (40)$$

From (38), (39) and (40) we deduce that there exist two constants $0 < C_1 < C_2$ such that, for every sufficiently large k ,

$$C_1 \frac{\tilde{L}(k)}{k^\rho} < \sum_{j=k}^\infty \frac{\tilde{L}(k)}{k^{1+\rho}} < C_2 \frac{\tilde{L}(k)}{k^\rho}.$$

Going back to (37), we find for sufficiently large i

$$\mathbf{E}[Z_i Z_j] \leq 2^{i\rho} \left(C_2 \frac{\tilde{L}(2^{j-1})}{2^{(j-1)\rho}} - C_1 \frac{\tilde{L}(2^j)}{2^{j\rho}} \right),$$

and now, by (35)

$$\begin{aligned} \sum_{m+1 \leq i < j \leq m+n} \mathbf{E}[Z_i Z_j] &= \sum_{m+3 \leq i+2 \leq j \leq m+n} \mathbf{E}[Z_i Z_j] + \sum_{i=m+1}^{m+n-1} \mathbf{E}[Z_i Z_{i+1}] = \\ &\leq \sum_{j=m+2}^{m+n} \left(C_2 \frac{\tilde{L}(2^{j-1})}{2^{(j-1)\rho}} - C_1 \frac{\tilde{L}(2^j)}{2^{j\rho}} \right) \sum_{i=m+1}^{j-1} 2^{i\rho} + \sum_{i=m+1}^{m+n-1} \mathbf{E}[Z_i^2]^{1/2} \mathbf{E}[Z_{i+1}^2]^{1/2} \\ &\leq C \left(\sum_{j=m+1}^{m+n-1} \tilde{L}(2^j) + n \right) \leq C \{ (m+n)^\gamma - m^\gamma + n \}. \end{aligned} \quad (41)$$

Now we insert (36) and (41) into (27) and obtain

$$\mathbf{E}\left[\left(\sum_{i=m+1}^{m+n} Z_i\right)^2\right] \leq C\{(m+n)^\gamma - m^\gamma + n\} \leq C\{(m+n)^{\gamma \vee 1} - m^{\gamma \vee 1}\},$$

and we conclude by Theorem 4.6. □

Remark 4.8 As clearly stated at the beginning of this section, Theorem 4.1 holds in the case $\alpha > 1$. We believe that this is due to the particular arguments used for the proof, and that it is possible to extend the ASLLT also to the case $\alpha < 1$. The critical case $\alpha = 1$ remains unexplored till now. Another not yet investigated situation is for $\alpha = 2$ with $x \mapsto E[X^2 1_{\{|X| \leq x\}}]$ slowly varying and $E[X^2] = \infty$ with $x \mapsto x^2 P(|X| > x)$ not slowly varying. Hopefully, we shall treat these cases in another paper.

References

- [1] **Aaronson, J., Denker, M.** (1998) *Characteristic functions of random variables attracted to 1-stable laws*, Ann. Probab. 26, 1, 399–415.
- [2] **Bingham, N. H., Goldie, C. M., Teugels, J. L.** (1987) *Regular variation* Encyclopedia of Mathematics and Its Applications, vol. 27, Cambridge University Press, Cambridge.
- [3] **Feller, W.** (1971) *An Introduction to Probability Theory and its Applications*, Vol. 2., second ed., John Wiley and Sons, Inc., New York.
- [4] **Giuliano Antonini, R., Weber, M.** (2008) *The theta-dependence coefficient and an Almost Sure Limit Theorem for Random Iterative Models*, Stat. Prob. Lett. 78, 564–575.
- [5] **Giuliano Antonini, R., Weber, M.** (2011) *Almost Sure Local Limit Theorems with rate*, Stoch. Anal. Appl. 29 no. 5, 779–798.
- [6] **Ibragimov, I. A., Linnik, Y. V.** (1971) *Independent and Stationary Sequences of Random Variables*, J. F. C. Kingman, ed. Wolters-Noordhoff, Gröningen.
- [7] **Loève, M.** (1960) *Probability Theory*, second ed., D. Van Nostrand Co., Inc., Princeton.
- [8] **Philipp, W., Stout, W.** (1975) *Almost sure invariance principles for partial sums of weakly dependent random variables*, Memoirs Amer. Math. Soc. 161.
- [9] **Szewczak, Z. S.** (2012) *Relative stability in strictly stationary random sequences*, Stochastic Process. Appl. 122 8, 2811–2829.
- [10] **Weber, M.** (2011) *A sharp correlation inequality with applications to almost sure local limit theorem*, Probab. Math. Statist. 31, 79–98.
- [11] **Zolotarev, V.M.** (1986) *One-dimensional stable distributions*, Translations of Mathematical Monographs, vol. 65, AMS, Providence.