

Plane-like minimizers and differentiability of the stable norm

A. Chambolle ^{*} M. Goldman [†] M. Novaga [‡]

Abstract

In this paper we investigate the strict convexity and the differentiability properties of the stable norm, which corresponds to the homogenized surface tension for a periodic perimeter homogenization problem (in a regular and uniformly elliptic case). We prove that it is always differentiable in totally irrational directions, while in other directions, it is differentiable if and only if the corresponding plane-like minimizers satisfying a strong Birkhoff property foliate the torus. We also discuss the issue of the uniqueness of the correctors for the corresponding homogenization problem.

1 Introduction

We wish to investigate here a conjecture raised by Caffarelli and De La Llave in [20] concerning the differentiability of the so-called stable norm (or minimal action functional) in geometric Weak KAM theory. When considering the area functional in the Euclidean space, it is a classical result that hyperplanes are minimizers under compact perturbations. Caffarelli and De La Llave [20] proved the existence of plane-like minimizers for more general integrands of the form

$$\int_{\partial^* E} F(x, \nu^E) d\mathcal{H}^{d-1}(x) + \int_E g(x) dx$$

where $F(x, \nu)$ is periodic in x , convex and one-homogeneous in ν and where g is a periodic function. It is then possible to define the homogenized energy $\varphi(p)$ which represents the average energy of a plane-like minimizer in the direction p . Our aim is to study the regularity properties of this function φ . Our main results are the following:

- If p is “totally irrational” then $\nabla\varphi(p)$ exists.

^{*}CMAP, Ecole Polytechnique, CNRS, Palaiseau, France, email: antonin.chambolle@cmap.polytechnique.fr

[†]CMAP, Ecole Polytechnique, CNRS, Palaiseau, France, email: michael.goldman@cmap.polytechnique.fr

[‡]Dipartimento di Matematica, Università di Padova, via Trieste 63, 35121 Padova, Italy, email: novaga@math.unipd.it

- The same occurs for any p such that the plane-like minimizers satisfying the strong Birkhoff property (see Section 4.1) give rise to a foliation of the space.
- If there is a gap in this lamination and if $(q_1, \dots, q_k) \in \mathbb{Z}^d$ is a maximal family of independent vectors such that $q_i \cdot p = 0$, then $\partial\varphi(p)$ is a convex set of dimension k , and φ is differentiable in the directions which are orthogonal to $\{q_1, \dots, q_k\}$. In particular if p is not totally irrational then φ is not differentiable at p .
- φ^2 is strictly convex.

We also discuss the uniqueness of the minimizers (see Theorem 4.21 and Appendix B).

Our approach is based on the cell formula for φ , introduced by the first author and Thouroude in [21] (see (6)). This formula provides a characterization of φ as the support function of some convex set C which gives the expression of its subgradient. Since φ is a convex function, it is differentiable at a point p if and only if $\partial\varphi(p)$ is reduced to a point. The set C being made of the integral over the unit cell of the calibrations in the direction p , the problem of the differentiability of φ reduces to the investigation of whether two different calibrations in a given direction p can have or not a different mean. The idea is to prove that every calibration calibrates every plane-like minimizer satisfying the strong Birkhoff property (see Section 4.1) and is thus prescribed on the union of these sets which form a lamination of the space. If there is no gap in this lamination (i.e. if it is a foliation) then the value of any calibration is fixed everywhere and thus its mean is also prescribed and φ is differentiable at p . If p is totally irrational, every gap of the lamination must have finite volume. This can be used to show that the integral over a gap of two different calibrations must coincide, implying again the differentiability of φ . For non totally irrational vectors, the gap will be of infinite volume and using different heteroclinic minimizers inside the gap, one can show that there exists two different calibrations with different means, proving that φ is non-differentiable.

One of the interesting features of our work is the connections it makes between the plane-like minimizers constructed in [21] (as level sets of appropriate correctors in a periodic homogenization problem), and the class of recurrent plane-like minimizers satisfying the strong Birkhoff property (see Section 4.1). We also give a clear presentation of the notion of calibration, and clarify the structure of the subgradient of functionals with linear growth (see also [5]), which could be of independent interest. Among the by-products of our analysis we obtain a proof of Mañé's conjecture in this setting (see Theorem 4.21 and Appendix B).

In [9, 25] the authors study the differentiability properties of the stable norm of the homology class $H_{d-1}(M, \mathbb{R})$ for compact orientable Riemannian manifolds M . If M is the d -dimensional torus with a Riemannian metric, the stable norm is exactly our function φ . In this case, our results generalize [9, 25] in the sense that we consider interfacial energies which do not necessarily come from a Riemannian metric, and we also allow the presence of a volume term. In some sense, a contribution of our work is also a simplification of some of the arguments

of [9, 25], which comes from the use of the cell formula (6).

The stable norm is a geometric analog of the minimal action functional of KAM theory whose differentiability has first been studied by Aubry and Mather [8, 29] for geodesics on the two dimensional torus. The results of Aubry and Mather have then been extended by Moser [32], in the framework of non-parametric integrands, and more recently by Senn and Bessi [35, 15]. In this context, the study of the set of non-selfintersecting minimizers, which correspond to our plane-like minimizers satisfying the Birkhoff property has been performed by Moser and Bangert [31, 13], whereas the proof of the strict convexity of the minimal action has been recently shown by Senn [36]. Another related problem is the homogenization of periodic Riemannian metrics (geodesics are objects of dimension one whereas in our problem the hypersurfaces are of codimension one). We refer to [19, 28, 18] for more information on this problem.

The plan of the paper is the following. In Section 2 we recall some well known facts about functions of bounded variation, sets of finite perimeter, convex anisotropies, pairings between BV functions and bounded vector fields, and introduce the notion of Class A minimizers, plane-like minimizers. We define the stable norm, which is, in fact, the homogenized surface tension of a periodic homogenization problem and can be recovered with a cell formula. In Section 3 we give some general properties of Class A Minimizers. In Section 4 we define the calibrations and prove an important regularity result (Theorem 4.5). In Section 4.1 we introduce the Birkhoff property and prove that every calibration calibrates every plane-like minimizer which satisfies this strong Birkhoff property. In Section 4.2 we construct heteroclinic surfaces in every gap and use them in Section 5 to study the strict convexity and differentiability properties of φ . In Section 6 we give some simple examples of energies for which the existence or absence of gaps can be proven. Finally in Section 7 we show that the set of stable norms that can be attained starting from isotropic energies is dense in the set of all symmetric anisotropies. Appendix A gives an elementary proof of a separation result in \mathbb{Z}^d , while in Appendix B it is shown a generic uniqueness result for the correctors minimizing the cell formula.

Acknowledgments.

The second author acknowledges very interesting discussions with G. Thouroude and B. Merlet. The third author acknowledges partial support by the Fondazione CaRiPaRo Project “Nonlinear Partial Differential Equations: models, analysis, and control-theoretic problems”. Some of the main ideas of this work have been clarified during a fruitful stay of the first two authors at the University of Padova.

2 Notation and main assumptions

2.1 Basic notation

We let $Q' = (0, 1)^d$, $Q = [0, 1]^d$ and \mathbb{T} be the torus $(\mathbb{R}^d/\mathbb{Z}^d)$. For $m \in \mathbb{N}$, we let \mathcal{H}^m be the Hausdorff measure of dimension m . Given a vector $p \in \mathbb{R}^d$ we say that

- p is totally irrational if there is no $q \in \mathbb{Z}^d \setminus \{0\}$ such that $p \cdot q = 0$,
- p is not totally irrational if there exists such a $q \in \mathbb{Z}^d \setminus \{0\}$,
- p is rational if $p \in \mathbb{R} \cdot \mathbb{Z}^d$,
- p is irrational if it is not rational.

For a given $p \in \mathbb{R}^d$ which is not totally irrational we let

$$\Gamma(p) := \{q \in \mathbb{Z}^d : q \cdot p = 0\}.$$

Then, there exists $(q_1, \dots, q_k) \in \Gamma(p)$ such that $\Gamma(p) = \text{Span}_{\mathbb{Z}}(q_1, \dots, q_k)$. By a Gram-Schmidt procedure we can find $(\bar{q}_1, \dots, \bar{q}_k) \in \Gamma(p)$ such that $(\bar{q}_1, \dots, \bar{q}_k)$ is an orthogonal basis of $V^r(p) := \text{Span}_{\mathbb{R}}(q_1, \dots, q_k)$. Notice that in general $\text{Span}_{\mathbb{Z}}(\bar{q}_1, \dots, \bar{q}_k) \neq \Gamma(p)$. We let $V^i(p) := (\mathbb{R}p \oplus V^r(p))^\perp$ be the set of irrational directions, that is, $V^i(p) \cap \mathbb{Z}^d = \{0\}$.

For $R > 0$ we let B_R be the open ball of radius R . Finally let \mathbb{S}^{d-1} be the unit sphere of \mathbb{R}^d . In this paper we will take as a convention that all the sets are oriented by their inward normal.

2.2 BV functions

We briefly recall the definition of function of bounded variation and set of finite perimeter. For a complete presentation we refer to [4, 24].

Definition 2.1. *Let Ω be an open set of \mathbb{R}^d , we will say that a function $u \in L^1(\Omega)$ is a function of bounded variation if*

$$\int_{\Omega} |Du| := \sup_{\substack{z \in \mathcal{C}_c^1(\Omega) \\ |z|_\infty \leq 1}} \int_{\Omega} u \operatorname{div} z \, dx < +\infty.$$

We denote by $BV(\Omega)$ the set of functions of bounded variation in Ω (when $\Omega = \mathbb{R}^d$ we simply write BV instead of $BV(\mathbb{R}^d)$). We define similarly the set $BV(\mathbb{T})$ of periodic functions of bounded variation by replacing the set $\mathcal{C}_c^1(\Omega)$ by $\mathcal{C}^1(\mathbb{T})$ in the definition of the total variation. We say that a set $E \subseteq \mathbb{R}^d$ is of finite perimeter if its characteristic function χ_E is of bounded variation and will denote its perimeter in an open set Ω by $P(E, \Omega) := \int_{\Omega} |D\chi_E|$ and will write simply $P(E)$ when $\Omega = \mathbb{R}^d$.

A function $v \in BV(\mathbb{T})$ can equivalently be seen as a Q -periodic function, with $\int_{\mathbb{T}} |Dv| = \int_Q |Dv|$ (which is also equal to $\int_Q |Dv|$ iff $|Dv|(\partial Q) = 0$, and $\int_{Q+x} |Dv|$ for a.e. x). We also let BV_{loc} be the set of functions of locally bounded variation. Similarly we will say that a set is of locally finite perimeter if its characteristic function is in BV_{loc} .

Definition 2.2. *Let E be a set of finite perimeter and let $t \in [0; 1]$. We then define*

$$E^{(t)} := \left\{ x \in \mathbb{R}^d : \lim_{r \downarrow 0} \frac{|E \cap B_r(x)|}{|B_r(x)|} = t \right\}.$$

We denote by $\partial E := (E^{(0)} \cup E^{(1)})^c$ the measure theoretical boundary of E . We define the reduced boundary of E by:

$$\partial^* E := \left\{ x \in \text{Spt}(|D\chi_E|) : \nu^E(x) := \lim_{r \downarrow 0} \frac{D\chi_E(B_r(x))}{|D\chi_E|(B_r(x))} \text{ exists and } |\nu^E(x)| = 1 \right\} \subseteq E^{(\frac{1}{2})}.$$

The vector $\nu^E(x)$ is the measure theoretical inward normal to the set E . When no confusion can be made, we simply denote ν^E by ν .

Proposition 2.3. *If E is a set of finite perimeter then $D\chi_E = \nu \mathcal{H}^{d-1} \llcorner \partial^* E$, $P(E) = \mathcal{H}^{d-1}(\partial^* E)$ and $\mathcal{H}^{d-1}(\partial E \setminus \partial^* E) = 0$.*

An important link between BV functions and sets of finite perimeter is given by the Coarea Formula:

Proposition 2.4. *Let $u \in BV(\Omega)$ then*

$$|Du|(\Omega) = \int_{\mathbb{R}} \mathcal{H}^{d-1}(\partial^* \{u > t\} \cap \Omega) dt.$$

2.3 Anisotropies

Let $F(x, p) : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ be a convex one-homogeneous function in the second variable such that there exists c_0 with

$$c_0 |p| \leq F(x, p) \leq \frac{1}{c_0} |p| \quad \forall (x, p) \in \mathbb{R}^d \times \mathbb{R}^d.$$

We say that F is elliptic if for some $\delta > 0$, the function $F - \delta|p|$ is still a convex function. We denote by $W(x) := \{p : F(x, p) \leq 1\}$ the unit ball of $F(x, \cdot)$ (when F does not depend on the space variable x we will just denote it by W). We define the polar function of F by

$$F^\circ(x, z) := \sup_{\{F(x, p) \leq 1\}} z \cdot p$$

so that $(F^\circ)^\circ = F$. If we denote by $F^*(x, z)$ the convex conjugate of F with respect to the second variable then $\{F^*(x, z) = 0\} = \{F^\circ \leq 1\}$. If $F(x, \cdot)$ is differentiable then, for every $p \in \mathbb{R}^d$,

$$F(x, p) = p \cdot \nabla_p F(x, p)$$

and

$$z \in \{F^\circ(x, \cdot) \leq 1\} \text{ with } p \cdot z = F(x, p) \iff z = \nabla_p F(x, p).$$

If F is elliptic and $\mathcal{C}^2(\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\})$ then F° is also elliptic and $\mathcal{C}^2(\mathbb{R}^d \times \mathbb{R}^d \setminus \{0\})$. Moreover for every $x, y, z \in \mathbb{R}^d$, there holds

$$F^2(x, y) - F^2(x, z) \geq 2(F(x, z)\nabla_p F(x, z)) \cdot (y - z) + C|y - z|^2 \quad (1)$$

for some constant C not depending on x (and the same holds for F°). Inequality (1) just state that F^2 is uniformly convex. We refer to [34] for a proof of these results and much more about convex bodies.

2.4 Pairings between measures and bounded functions

We fix in the following an elliptic anisotropy F . Following Anzellotti [7] we define a generalized trace $[z, Du]$ for functions u with bounded variation and bounded vector fields z with divergence in L^d .

Definition 2.5. *let Ω an open set with Lipschitz boundary, let $u \in BV(\Omega)$ and let $z \in L^\infty(\Omega, \mathbb{R}^d)$ with $\operatorname{div} z \in L^d(\Omega)$. We define the distribution $[z, Du]$ by*

$$\langle [z, Du], \psi \rangle = - \int_{\Omega} u \psi \operatorname{div} z - \int_{\Omega} u z \cdot \nabla \psi \quad \forall \psi \in \mathcal{C}_c^\infty(\Omega).$$

If $u \in BV(\mathbb{T})$ and $z \in L^\infty(\mathbb{T}, \mathbb{R}^d)$, with $\operatorname{div} z \in L^d(\mathbb{T})$, we can easily define the distribution $[z, Du]$ in a similar way.

Theorem 2.6. *The distribution $[z, Du]$ is a bounded Radon measure on Ω and if ν is the inward unit normal to Ω , there exists a function $[z, \nu] \in L^\infty(\partial\Omega)$ such that the generalized Green's formula holds,*

$$\int_{\Omega} [z, Du] = - \int_{\Omega} u \operatorname{div} z - \int_{\partial\Omega} [z, \nu] u d\mathcal{H}^{d-1}.$$

The function $[z, \nu]$ is the generalized (inward) normal trace of z on $\partial\Omega$. If $u \in BV(\mathbb{T})$ and $z \in L^\infty(\mathbb{T}, \mathbb{R}^d)$, with $\operatorname{div} z \in L^d(\mathbb{T})$, there holds

$$\int_{\mathbb{T}} [z, Du] = - \int_{\mathbb{T}} u \operatorname{div} z.$$

Given $z \in L^\infty(\mathbb{T}, \mathbb{R}^d)$, with $\operatorname{div} z \in L^d(\mathbb{T})$, we can also define the generalized trace of z on ∂E , where E is a set of locally finite perimeter. Indeed, for every bounded open set Ω with smooth boundary, we can define as above the measure $[z, D\chi_E]$ on Ω . Since this measure is absolutely continuous with respect to $|D\chi_E| = \mathcal{H}^{d-1} \llcorner \partial^* E$ we have

$$[z, D\chi_E] = \psi_z(x) \mathcal{H}^{d-1} \llcorner \partial^* E$$

with $\psi_z \in L^\infty(\partial^* E)$ independent of Ω . We denote by $[z, \nu] := \psi_z$ the generalized (inward) normal trace of z on ∂E . If E is a bounded set of finite perimeter, by taking Ω strictly containing E , we have the generalized Gauss-Green Formula

$$\int_E \operatorname{div} z = - \int_{\partial^* E} [z, \nu] d\mathcal{H}^{d-1}.$$

We notice that there has been a lot of interest in defining the trace of bounded vector fields with divergence a bounded measure, on boundaries of sets of finite perimeters [3, 22], since this is related to the study of conservations laws.

2.5 Plane-like minimizers and the stable norm

Given a set E of locally finite perimeter, we consider the energy

$$\mathcal{E}(E, A) := \int_{\partial^* E \cap A} F(x, \nu^E) d\mathcal{H}^{d-1} + \int_{E \cap A} g(x) dx \quad (2)$$

with F, g Q -periodic continuous functions, and $\int_Q g = 0$. Here ν^E is the inner normal to $\partial^* E$, so that $\int_{\partial^* E \cap A} F(x, \nu^E) d\mathcal{H}^{d-1} = \int_A F(x, D\chi_E)$. We also assume that F is convex, one-homogeneous with respect to its second variable, and satisfies for some $c_0 > 0$

$$c_0 |p| \leq F(x, p) \leq c_0^{-1} |p| \quad (3)$$

for any $(x, p) \in \mathbb{R}^d \times \mathbb{R}^d$. A fundamental assumption throughout the paper is that the energy is coercive, in the sense that

$$\mathcal{E}(E, Q) \geq \delta P(E, Q) \quad (4)$$

for some $\delta > 0$ independent of E . This will be ensured if (3) holds and g is small enough in some appropriate norm. When $F(x, p) = |p|$ and $g = 0$, the energy \mathcal{E} is just the perimeter. In that case, it is well known that planes are minimizers under compact perturbations. Indeed, the Bernstein Theorem states that, if $d \leq 7$, the only minimizers of the perimeter under compact perturbations are the hyperplanes (see [24]). In [20], Caffarelli and De la Llave proved that, for general energies \mathcal{E} , even if hyperplanes are not minimizers anymore there still exist plane-like minimizers.

Definition 2.7. *We say that a set E of locally finite perimeter is a Class A Minimizer of \mathcal{E} if, for any $R > 0$, the set E minimizes $\mathcal{E}(E, B_R)$ under compact perturbation in B_R .*

Theorem 2.8 ([20]). *There exists $M > 0$ depending only on c_0 and δ such that for every $p \in \mathbb{R}^d \setminus \{0\}$ and $a \in \mathbb{R}$, there exists a Class A Minimizer E of \mathcal{E} such that*

$$\left\{ x \cdot \frac{p}{|p|} > a + M \right\} \subseteq E \subseteq \left\{ x \cdot \frac{p}{|p|} > a - M \right\}. \quad (5)$$

Moreover ∂E is connected.

Definition 2.9. *If E satisfies (5) for some $M > 0$, we say that E is a plane-like set. If E is a Class A Minimizer of \mathcal{E} satisfying (5) we say that E is a plane-like minimizer.*

The existence of Class A Minimizers is closely related to the existence of minimizers of functionals of the form

$$\int_{\mathbb{R}^d} G(x, u, \nabla u)$$

satisfying $\sup |u(x) - p \cdot x| < +\infty$ for some p (the vector p is often called the rotation vector) in Weak KAM theory (see [31, 35]). The analogous of the minimal action functional of Weak KAM theory in our setting is the so-called stable norm introduced by Federer.

Definition 2.10. *Let $p \in \mathbb{R}^d \setminus \{0\}$ and let E be a plane-like minimizer of \mathcal{E} in the direction p . We set*

$$\varphi(p) := |p| \lim_{R \rightarrow \infty} \frac{1}{\mathcal{H}^{d-1}(\partial B_R)} \mathcal{E}(E, B_R).$$

Caffarelli and De La Llave proved that this limit exists and does not depend on E . In [21], the first author and Thouroude proved the cell formula:

$$\varphi(p) = \min \left\{ \int_{\mathbb{T}} F(x, p + Dv(x)) + \int_{\mathbb{T}} g(x)(v(x) + p \cdot x) dx : v \in BV(\mathbb{T}) \right\}. \quad (6)$$

It is obvious from (6) that φ is a convex, one-homogeneous function. It is also shown in [21] that the minimizers of (6) give an easy way to construct plane-like minimizers:

Proposition 2.11. *Let v_p be a minimizer of (6) then for every $s \in \mathbb{R}$, the set $\{v_p(x) + p \cdot x > s\}$ is a plane-like minimizer of \mathcal{E} in the direction p .*

We will make the following additional hypotheses on F, g :

- F is $\mathcal{C}^{2,\alpha}(\mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}))$, and $g \in \mathcal{C}^{1,\alpha}(\mathbb{R}^d)$,
- F is elliptic (that is $F(x, p) - C|p|$ is a convex function of p)

Under these assumptions, one can show that there exists a periodic vector field σ of class $\mathcal{C}^{2,\alpha}$, with $\operatorname{div} \sigma = g$ and such that $F'(x, p) = F(x, p) - \sigma(x) \cdot p \geq c'_0 |p|$, for some $c'_0 > 0$. The proof follows the same idea as in [21] (which is based on [17]). From (4) and (3), it follows that $\int_{\mathbb{T}} F(x, Dv) + \int_{\mathbb{T}} (1 + \varepsilon)gv dx \geq \delta/2 \int_{\mathbb{T}} |Dv|$ if $\varepsilon > 0$ is small enough. Hence, $(1 + \varepsilon)g \in \partial H(0)$, the subgradient at zero of $H(v) := \int_{\mathbb{T}} F(x, Dv)$. This is a 1-homogeneous, convex, l.s.c. functional defined on $L^p(\mathbb{T})$ for any $p \in [1, \infty]$ (letting $H(v) = +\infty$ for $v \notin BV(\mathbb{T})$), which is the support function of

$$K_H = \{\operatorname{div} \sigma : \sigma \in \mathcal{C}^\infty(\mathbb{T}; \mathbb{R}^d), F^\circ(x, \sigma(x)) \leq 1 \forall x \in \mathbb{T}\}.$$

From Hahn-Banach's theorem, one deduces that $(1 + \varepsilon)g$ is in the closure of K_H for the topology $(L^{p'}, L^p)$. For $p < \infty$, large, it coincides with the strong closure, hence $(1 + \varepsilon)g$

is the L^p -limit of a sequence $\operatorname{div} \sigma_n$, with σ_n smooth and $F^\circ(x, \sigma_n(x)) \leq 1$. We solve then $\Delta u_n = g - \operatorname{div}(\sigma_n/(1+\varepsilon))$ in the torus. On one hand, $u_n \in \mathcal{C}^{3,\alpha}(\mathbb{T})$ by elliptic regularity. On the other hand, $\|u_n\|_{2,p} \leq C\|g - \operatorname{div}(\sigma_n/(1+\varepsilon))\|_p$, which goes to zero with n . In particular, $\|\nabla u_n\|_\infty$ is arbitrarily small. We choose n so large that this quantity is less than $c_0\varepsilon/(2+2\varepsilon)$, and let $\sigma = \sigma_n/(1+\varepsilon) + \nabla u_n \in \mathcal{C}^{2,\alpha}(\mathbb{T})$. Then for any p , $\sigma(x) \cdot p \leq F(x, p)/(1+\varepsilon) + \nabla u_n \cdot p \leq F(x, p) - c_0\varepsilon/(1+\varepsilon)|p| + c_0\varepsilon|p|/(2+2\varepsilon)$, so that the claim holds with $c'_0 = c_0\varepsilon/(2+2\varepsilon) > 0$. For this reason, we can replace without loss of generality F with F' and g with zero in (2) without changing anything to the problem. To simplify the notation we will therefore assume that $g = 0$ in the rest of the paper.¹

In the following we let

$$X := \{z \in L^\infty(\mathbb{T}, \mathbb{R}^d) : \operatorname{div} z = 0, F^\circ(x, z(x)) \leq 1 \text{ a.e.}\}.$$

The next result is the starting point of our analysis on the differentiability properties of φ .

Proposition 2.12. *There holds*

$$\varphi(p) = \sup_{z \in X} \left(\int_{\mathbb{T}} z \right) \cdot p.$$

Proof. For every $v \in BV(\mathbb{T})$ and every $z \in X$,

$$\begin{aligned} \int_{\mathbb{T}} [Dv + p, z] &\leq \int_{\mathbb{T}} F(x, Dv + p) && \text{hence} \\ - \int_{\mathbb{T}} v \operatorname{div} z + \left(\int_{\mathbb{T}} z \right) \cdot p &\leq \int_{\mathbb{T}} F(x, Dv + p) && \text{and since } \operatorname{div} z = 0, \\ \left(\int_{\mathbb{T}} z \right) \cdot p &\leq \int_{\mathbb{T}} F(x, Dv + p) \\ \left(\int_{\mathbb{T}} z \right) \cdot p &\leq \varphi(p) && \text{taking the infimum on } v, \text{ thus} \\ \sup_{z \in X} \left(\int_{\mathbb{T}} z \right) \cdot p &\leq \varphi(p). \end{aligned}$$

To prove the opposite inequality let $v_p \in BV(\mathbb{T})$ be a minimizer in the definition of $\varphi(p)$ and let $z \in X$ be such that

$$\int_{\mathbb{T}} [Dv_p + p, z] = \int_{\mathbb{T}} F(x, Dv_p + p),$$

then

$$\varphi(p) = \int_{\mathbb{T}} F(x, Dv_p + p) \leq \sup_z \left(\int_{\mathbb{T}} z \right) \cdot p.$$

□

¹The hypothesis $g \in \mathcal{C}^{1,\alpha}$ could be relaxed to g Lipschitz. Indeed, the regularity hypothesis on F is mainly there to ensure that regularity theory and maximum principle hold for the plane-like minimizers (see Proposition 3.4).

Proposition 2.12 shows that φ is the support function of the convex set

$$C := \left\{ \int_Q z(x) dx : z \in X \right\}, \quad (7)$$

so that

$$\varphi(p) = \sup_{\xi \in C} \xi \cdot p.$$

Observe that C is trivially closed in \mathbb{R}^d , since the associated vector fields z are uniformly bounded in L^∞ , and the constraints pass to the limit.

The subgradient of φ at $p \in \mathbb{R}^d$ is given by

$$\partial\varphi(p) = \{\xi \in C : \xi \cdot p = \varphi(p)\}. \quad (8)$$

Any $\xi \in \partial\varphi(p)$ is associated to a field z as in (7). Our analysis relies on the following fact

$$\varphi \text{ is differentiable at } p \iff \partial\varphi(p) \text{ is a singleton.}$$

3 Properties of Class A Minimizers

We first start by recalling some well known facts about Class A and plane-like minimizers (see [2, 20, 21]).

Proposition 3.1. *Let E be a Class A Minimizer. Then the reduced boundary ∂^*E is of class $C^{2,\alpha}$ and $\mathcal{H}^{d-3}(\partial E \setminus \partial^*E) = 0$. Moreover, there exists $\gamma > 0$ and $\beta > 0$ such that*

- if $x \in \bar{E}$ then $|B_r(x) \cap E| \geq \gamma r^d$ for every $r > 0$,
- if $x \in E^c$ then $|B_r(x) \setminus E| \geq \gamma r^d$ for every $r > 0$,
- if $x \in \partial E$ then $\beta r^{d-1} \leq |D\chi_E|(B_r(x)) \leq \frac{1}{\beta} r^{d-1}$ for every $r > 0$.

As a consequence, we will assume in the following that our Class A Minimizers are all open sets (indeed, we can identify them with their points of density 1, which clearly are an open set because of the second density estimate). Moreover, the topological boundary ∂E agrees with the measure theoretical boundary of E .

The stability of plane-like minimizers under convergence is a crucial point in the theory.

Proposition 3.2. *Let E_n be a sequence of plane-like minimizers satisfying (5) with a uniform M and converging in the L^1_{loc} topology to a set E , then E is also a plane-like minimizer. Moreover, $E_n \rightarrow \bar{E}$, $E_n^c \rightarrow E^c$, and of $\partial E_n \rightarrow \partial E$ in the Kuratowski sense.*

Proof. The stability of the plane-like minimizers is a well known fact (see [20, Section 9]). The Kuratowski (or local Hausdorff) convergence easily follows from the uniform density

estimates for plane-like minimizers (Proposition 3.1). Indeed, let $\varepsilon > 0$ be fixed and let $x \in E \cap \{y : d(y, \partial E) > \varepsilon\}$. If x is not in E_n then by the density estimates we have

$$|E_n \Delta E| \geq |B_\varepsilon(x) \setminus E_n| \geq \gamma \varepsilon^d.$$

This is impossible if n is big enough because $|E_n \Delta E|$ tends to zero. Similarly, we can show that for n big enough, all the points of $E^c \cap \{y : d(y, \partial E) > \varepsilon\}$ are outside E_n . This shows that $\partial E_n \subseteq \{y : d(y, \partial E) \leq \varepsilon\}$. Inverting E_n and E , the same argument proves that $\partial E \subseteq \{y : d(y, \partial E_n) \leq \varepsilon\}$ giving the Kuratowski convergence of ∂E_n to ∂E . \square

Another simple (and classical) consequence of the density estimates is the following

Proposition 3.3. *Let $u \in BV_{\text{loc}}(\mathbb{R}^d)$ then for every $R > 0$,*

$$Spt(|Du|) \cap B_R = B_R \cap \overline{\bigcup_s \partial^* \{u > s\}}.$$

where in the union we consider only the levels for which $\{u > s\}$ has finite perimeter in B_R . If in addition v_p is a minimizer of (6) and $u(x) = v_p(x) + p \cdot x$, then

- $P(\{u > s\} \cap B_R) < +\infty$ for every $s \in \mathbb{R}$;
- $\overline{\partial^* \{u > s\}} = \partial \{u > s\}$;
- the function u^+ is u.s.c., and u^- is l.s.c.;
- $Spt(|Du|) \cap B_R = \left(\bigcup_{s \in \mathbb{R}} \partial \{u > s\} \cap B_R \right) \cup \left(\bigcup_{s \in \mathbb{R}} \partial \{u \geq s\} \cap B_R \right)$.

Here, $u^\pm(x)$ are classically defined as the approximate upper and lower limits of u at x , see [4].

Proof. Let us show the first assertion. For $u \in BV_{\text{loc}}(\mathbb{R}^d)$, if $x \notin Spt(|Du|)$, then there exists $\rho > 0$ with $B_\rho(x) \subseteq \{Du = 0\}$ and thus u is constant on $B_\rho(x)$, which implies that

$$x \notin \overline{\bigcup_s \partial^* \{u > s\} \cap B_R}.$$

On the contrary, if $x \in Spt(|Du|)$ then for every $\rho > 0$, by the Coarea Formula,

$$|Du|(B_\rho(x)) = \int_{\mathbb{R}} \mathcal{H}^{d-1}(\partial^* \{u > s\} \cap B_\rho(x)) ds > 0$$

thus for every $\rho > 0$ there exists $x_\rho \in B_\rho(x) \cap \partial^* \{u > s_\rho\}$ for some s_ρ since x_ρ tends to x when $\rho \rightarrow 0$, this proves the other inclusion.

Given a minimizer v_p , the other properties follow easily from the density estimates. \square

The following maximum principle for minimizers is a cornerstone of the theory.

Proposition 3.4. *Let $E_1 \subseteq E_2$ be two Class A Minimizers with connected boundary, then $\mathcal{H}^{d-3}(\partial E_1 \cap \partial E_2) = 0$.*

Proof. We shall now prove that $\partial^* E_1 \cap \partial^* E_2 = \emptyset$. Let us assume by contradiction that $\partial^* E_1 \cap \partial^* E_2 \neq \emptyset$ then we can find $\bar{x} \in \partial^* E_1 \cap \partial^* E_2$ such that $\partial^* E_1 \cap B_r(\bar{x}) \neq \partial^* E_2 \cap B_r(\bar{x})$ for every $r > 0$. Since $E_1 \subseteq E_2$, E_1 and E_2 have the same tangent space at \bar{x} and they can be seen as graphs over the same domain D of two functions $v_1, v_2 \in C^{2,\alpha}(D)$, with $v_2 \geq v_1$. For $(y, r, p) \in \mathbb{R}^{d-1} \times \mathbb{R} \times \mathbb{R}^{d-1}$, let

$$\tilde{F}(y, r, p) := F((y, r), (-p, 1)).$$

The functions $v_i \in C^{2,\alpha}(D)$, $i = 1, 2$, (locally) minimize the functional

$$\int_D \tilde{F}(y, u, \nabla u) dy$$

and thus solve the elliptic PDE with Hölder continuous coefficients

$$\frac{\partial \tilde{F}}{\partial r}(y, v_i, \nabla v_i) - \operatorname{div}[\nabla_p \tilde{F}(y, v_i, \nabla v_i)] = 0. \quad (9)$$

Consider the function $w = v_2 - v_1$. Up to reducing the domain, we can assume that $\bar{x} \in \partial D$, $w > 0$ in D and $w(\bar{x}) = 0$. We must then have $\nabla w(\bar{x}) = 0$. Let

$$\begin{aligned} A(x) &:= \int_0^1 \nabla_p^2 \tilde{F}(x, v_2(x), \nabla v_2(x) - \nabla v_1(x)) dt \\ B_1(x) &:= \int_0^1 \nabla_p \frac{\partial}{\partial r} \tilde{F}(x, tv_2(x) + (1-t)v_1(x), \nabla v_2(x)) dt \\ B_2(x) &:= \int_0^1 \nabla_p \frac{\partial}{\partial r} \tilde{F}(x, v_2(x), t\nabla v_2(x) + (1-t)\nabla v_1(x)) dt \\ c_1(x) &:= \int_0^1 \frac{\partial^2}{\partial r^2} \tilde{F}(x, tv_2 + (1-t)v_1, \nabla v_2) dt \end{aligned}$$

then w satisfies the linear non-degenerate elliptic PDE,

$$-\operatorname{div}(A(x)\nabla^2 w) - \operatorname{div}(B_1(x)w) + B_2(x) \cdot \nabla w + c_1(x)w = 0.$$

By Hopf's Lemma [23, Lemma 3.4], this implies that $\nabla w(\bar{x}) \cdot \nu_D < 0$, which gives a contradiction. Thus ∂E_1 and ∂E_2 can only intersect in singular points which are of $(d-3)$ -Hausdorff measure zero. \square

Remark 3.5. In the case of isotropic functionals i.e. $F(x, \nu) = a(x)|\nu|$, [37] shows that in fact two minimizers which are contained one in the other cannot touch at all.

Proposition 3.6. *Let $d = 2$ and let E be a Class A Minimizer, then E is a plane-like minimizer and ∂E is connected.*

Proof. By Proposition 3.4, ∂E is of class $C^{2,\alpha}$ and is composed of a locally finite union of curves. Moreover each of these curves has infinite length by the minimality of E . Let γ be such a curve and H be a plane-like minimizer with connected boundary. Due to the regularity and minimality of E and H , ∂H can intersect γ in at most one point. Since by Theorem 2.8 there exist such plane-like minimizers inside every strip of width M , it follows that γ is included in such a strip. Using again the minimality of E , we then get that ∂E is connected and E is plane-like. \square

Proposition 3.6 is reminiscent of Bernstein Theorem, and the same result also holds for $d = 3$ under the additional assumption that F does not depend on x [39]. However, in [30] it is shown that it is non longer true in four dimensions, even for a function F independent of x .

4 Calibrations

We now introduce the notion of calibration.

Definition 4.1. *We say that a vector field $z \in X$ is a periodic calibration of a set E of locally finite perimeter if, for every open set A , we have*

$$\int_A [z, D\chi_E] = \int_A F(x, D\chi_E).$$

When no confusion can be made, by calibration we mean a periodic calibration.

The constant interest towards calibrations in the study of minimal surfaces, comes from the following result:

Theorem 4.2. *If E is a set for which there exists a calibration (not necessarily periodic), then E is a Class A Minimizer.*

Proof. It follows by integration by parts, using $\operatorname{div} z = 0$. \square

Calibrations are very stable objects, as shown by the following Proposition:

Proposition 4.3. *Let E_n be Class A minimizers converging in the L^1_{loc} -topology to a set E . Assume that the sets E_n are calibrated by $z_n \in X$ and that z_n converges to a field z weakly-* in L^∞ . Then z calibrates E (which is thus also a minimizer).*

Proof. First, observe that $z \in X$. A way to check this is to show that for $p \in \mathbb{Q}^d$, the average of $z \cdot p$ in any ball $B_\rho(x)$ is less than the average of $F(\cdot, p)$ (since it is true for z_n). For a.e. x it follows that $z(x) \cdot p \leq F(x, p)$ for all $p \in \mathbb{Q}^d$ (hence \mathbb{R}^d), that is, $F^\circ(x, z(x)) \leq 1$.

Let now $\varphi \in C_c^\infty(\mathbb{R}^d)$, $\varphi \geq 0$. Then since $z \in X$,

$$\begin{aligned} \int_{\partial^* E_n} \varphi F(x, \nu^{E_n}) d\mathcal{H}^{d-1} &= \int_{\mathbb{R}^d} \varphi [z_n, D\chi_{E_n}] = - \int_{E_n} z_n \cdot \nabla \varphi \\ &\xrightarrow{n \rightarrow \infty} - \int_E z \cdot \nabla \varphi = \int_{\mathbb{R}^d} \varphi [z, D\chi_E] \leq \int_{\partial^* E} \varphi F(x, \nu^E) d\mathcal{H}^{d-1} \end{aligned}$$

and the reverse inequality follows by lower-semicontinuity of the total variation. Hence z is a calibration for E . □

A natural way of producing calibrations for a set is through the cell problem (6). By Proposition 2.12 there exists $z \in X$ such that, for any minimizer v_p of (6),

$$\int_{\mathbb{T}} [z, Dv_p + p] = \int_{\mathbb{T}} F(x, p + Dv_p) = \varphi(p). \quad (10)$$

We say that such a vector field z is a calibration in the direction p . Conversely, if $z \in X$ and $v \in BV(\mathbb{T})$ are such that

$$\int_{\mathbb{T}} [z, Dv + p] = \int_{\mathbb{T}} F(x, p + Dv), \quad (11)$$

then v is a minimizer of (6) and z is a calibration in the direction p .

Proposition 4.4. *Let $z \in X$ be a calibration in the direction p , then for every minimizer v_p of (6) and every $s \in \mathbb{R}$, z calibrates the set $E_s := \{v_p + p \cdot x > s\}$. Conversely for $v \in BV(\mathbb{T})$, if $z \in X$ calibrates all the sets E_s then v is a minimizer of (6).*

Proof. If z is a calibration in the direction p then as noticed above, z calibrates all the solution v_p of (6). Let v_p be one of these solutions then by the Coarea Formula and [7, Prop. 2.7], for every Borel set $A \subseteq \mathbb{R}^d$

$$\begin{aligned} \int_{\mathbb{R}} \int_A F(x, D\chi_{E_s}) \, ds &= \int_A F(x, Dv_p + p) \\ &= \int_A [z, Dv_p + p] \\ &= \int_{\mathbb{R}} \int_A [z, D\chi_{E_s}] \, ds \\ &\leq \int_{\mathbb{R}} \int_A F(x, D\chi_{E_s}) \, ds \end{aligned}$$

and thus for almost every $s \in \mathbb{R}$, z calibrates E_s . Since for every s , $E_s = \cup_{s' > s} E_{s'}$, by Proposition 4.3, z calibrates in fact every E_s . The converse implication follows by the same argument. □

Theorem 4.5. *Let E be a Class A Minimizer, let $z \in X$ and let \bar{x} be a Lebesgue point of z . Then, if z calibrates E and if $\bar{x} \in \partial E$, we have that $\bar{x} \in \partial^* E$ and*

$$z(\bar{x}) = \nabla_p F(\bar{x}, \nu^E(\bar{x})). \quad (12)$$

Proof. Letting $z_\rho(y) = z(\bar{x} + \rho y)$, the assumption that \bar{x} is a Lebesgue point of z yields that $z_\rho \rightarrow \bar{z}$ in $L^1(B_R)$, hence also weakly-* in $L^\infty(B_R)$, for any $R > 0$, where $\bar{z} \in \mathbb{R}^d$ is a constant vector.

As usual, we let $E_\rho := (E - \bar{x})/\rho$ and we observe that E_ρ minimizes

$$\int_{\partial E_\rho \cap B_R} F(\bar{x} + \rho y, \nu^{E_\rho}(y)) d\mathcal{H}^{d-1}(y)$$

with respect to compactly supported perturbations of the set (in the fixed ball B_R). In particular, the sets E_ρ (and the boundaries ∂E_ρ) satisfy uniform density bounds, and hence are compact with respect to both local L^1 and Hausdorff convergence.

Hence, up to extracting a subsequence, we can assume that $E_\rho \rightarrow \bar{E}$, with $0 \in \partial \bar{E}$. Proposition 4.3 shows that \bar{z} is a calibration for the energy $\int_{\partial \bar{E} \cap B_R} F(\bar{x}, \nu^{\bar{E}}(y)) d\mathcal{H}^{d-1}(y)$, and that \bar{E} is a plane-like minimizer calibrated by \bar{z} .

It follows that $[\bar{z}, \nu^{\bar{E}}] = F(\bar{x}, \nu^{\bar{E}}(y))$ for \mathcal{H}^{d-1} -a.e. y in $\partial \bar{E}$, but since \bar{z} is a constant, we deduce that $\bar{E} = \{y \cdot \bar{\nu} \geq 0\}$ with $\bar{\nu} = F(\bar{\nu}) \nabla_p F^\circ(\bar{x}, \bar{z})$. In particular the limit \bar{E} is unique, hence we obtain the global convergence of $E_\rho \rightarrow \bar{E}$, without passing to a subsequence.

We want to deduce that $\bar{x} \in \partial^* E$, with $\nu^E(\bar{x}) = F(\bar{x}, \nu^E(\bar{x})) \nabla_p F^\circ(\bar{x}, \bar{z})$, which is equivalent to (12). The last identity is obvious from the arguments above, so that we only need to show that

$$\lim_{\rho \rightarrow 0} \frac{D\chi_{E_\rho}(B_1)}{|D\chi_{E_\rho}|(B_1)} = \bar{\nu}. \quad (13)$$

Assume we can show that

$$\lim_{\rho \rightarrow 0} |D\chi_{E_\rho}|(B_R) = |D\chi_{\bar{E}}|(B_R) \quad (= \omega_{d-1} R^{d-1}) \quad (14)$$

for any $R > 0$, then for any $\psi \in C_c^\infty(B_R; \mathbb{R}^d)$ we would get

$$\begin{aligned} \frac{1}{|D\chi_{E_\rho}|(B_R)} \int_{B_R} \psi \cdot D\chi_{E_\rho} &= - \frac{1}{|D\chi_{E_\rho}|(B_R)} \int_{B_R \cap E_\rho} \operatorname{div} \psi(x) dx \\ &\rightarrow - \frac{1}{|D\chi_{\bar{E}}|(B_R)} \int_{B_R \cap \bar{E}} \operatorname{div} \psi(x) dx = \frac{1}{|D\chi_{\bar{E}}|(B_R)} \int_{B_R} \psi \cdot D\chi_{\bar{E}} \end{aligned}$$

and deduce that the measure $D\chi_{E_\rho}/(|D\chi_{E_\rho}|(B_R))$ weakly- $*$ converges to $D\chi_{\bar{E}}/(|D\chi_{\bar{E}}|(B_R))$.

Using again (14)), we then obtain that

$$\lim_{\rho \rightarrow 0} \frac{D\chi_{E_\rho}(B_R)}{|D\chi_{E_\rho}|(B_R)} = \bar{\nu} \quad (15)$$

for almost every $R > 0$. Since $D\chi_{E_\rho}(B_{\mu R})/(|D\chi_{E_\rho}|(B_{\mu R})) = D\chi_{E_\rho/\mu}(B_R)/(|D\chi_{E_\rho/\mu}|(B_R))$ for any $\mu > 0$, (15) holds in fact for any $R > 0$ and (13) follows, so that $\bar{x} \in \partial^* E$.

It remains to show (14). First, we observe that, by minimality of E_ρ and \bar{E} plus the Hausdorff convergence of ∂E_ρ in balls, we can easily show the convergence of the energies

$$\lim_{\rho \rightarrow 0} \int_{\partial E_\rho \cap B_R} F(\bar{x} + \rho y, \nu^{E_\rho}(y)) d\mathcal{H}^{d-1}(y) = \int_{\partial \bar{E} \cap B_R} F(\bar{x}, \nu^{\bar{E}}(y)) d\mathcal{H}^{d-1}(y) \quad (16)$$

and, by the continuity of F ,

$$\lim_{\rho \rightarrow 0} \int_{\partial E_\rho \cap B_R} F(\bar{x}, \nu^{E_\rho}(y)) d\mathcal{H}^{d-1}(y) = \int_{\partial \bar{E} \cap B_R} F(\bar{x}, \nu^{\bar{E}}(y)) d\mathcal{H}^{d-1}(y). \quad (17)$$

Then, (13) follows from a variant of Reshetnyak's continuity theorem where instead of using the Euclidean norm as reference norm, we use the uniformly convex norm $F(\bar{x}, \cdot)$. Notice that this variant is covered by the original version of Reshetnyak [33]. For the reader's convenience we sketch the proof here, following very closely the proof of [4, Thm. 2.39]. In what follows, the point \bar{x} is fixed and thus we will not specify the dependence of the functions on \bar{x} (for example $F(p)$ will stand for $F(\bar{x}, p)$).

Let now $\mu_\rho := \nu^{E_\rho} \mathcal{H}^{d-1} \llcorner \partial^* E_\rho$, $\mu := \nu^{\bar{E}} \mathcal{H}^{d-1} \llcorner \partial^* \bar{E}$, $\theta_\rho := \frac{\nu^{E_\rho}}{F(\nu^{E_\rho})}$, $\theta = \frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}$ and $W := \{F(p) \leq 1\}$. Then we define the measures η_ρ on $B_R \times \partial W$ by setting $\eta_\rho := F(\mu_\rho) \otimes \delta_{\theta_\rho(x)}$. The sequence η_ρ is bounded and thus there exists a weakly-* converging subsequence to a measure η . Let $\pi : B_R \times \partial W \rightarrow B_R$ be the projection, then $F(\mu_\rho) = \pi_{\#} \eta_\rho$ and thus by [4, Rk. 1.71], $F(\mu_\rho)$ weakly-* converges to $\pi_{\#} \eta$, therefore by (17) and [4, Prop. 1.80] we get $\pi_{\#} \eta = F(\mu)$.

By the Disintegration Theorem [4, Th. 2.28], there exists a $F(\mu)$ -measurable map $x \rightarrow \eta_x$ such that $\eta_x(\partial W) = 1$ and $\eta = F(\mu) \otimes \eta_x$. Arguing exactly as in [4, Th. 2.38], we have

$$\int_{\partial W} y d\eta_x = \theta(x) \quad \text{for } F(\mu) - a.e. \ x \in B_R. \quad (18)$$

The anisotropic ball W being strictly convex and $\theta(x)$ being on its boundary, this will imply that indeed, $\eta_x = \delta_{\theta(x)}$ which will conclude the proof. Since F is strictly convex, for every $y \in \partial W$, (1), yields

$$F(y)^2 - F^2\left(\frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right) \geq 2F\left(\frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right) \left(\nabla F\left(\frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right)\right) \cdot \left(y - \frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right) + C \left|y - \frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right|^2,$$

from which it follows

$$2 \left(1 - \left(\nabla F\left(\frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right)\right) \cdot y\right) \geq C \left|y - \frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right|^2.$$

Integrating this inequality on ∂W and using (18) we get

$$0 = 2 \left(1 - \left(\nabla F\left(\frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right)\right) \cdot \left(\int_{\partial W} y d\eta_x\right)\right) \geq C \int_{\partial W} \left|y - \frac{\nu^{\bar{E}}}{F(\nu^{\bar{E}})}\right|^2$$

hence $\eta_x = \delta_{\theta(x)}$. The proof of (14) now easily follows. Indeed, since $\eta_\rho(B_R \times \partial W)$ converges to $F(\mu)(B_R) = \eta(B_R \times \partial W)$, using [4, Prop 1.80] we find

$$\begin{aligned} \lim_{\rho \rightarrow 0} \int |\theta_\rho(x)| dF(\mu_\rho)(x) &= \int_{B_R \times \partial W} |\theta_\rho(x)| d\eta_\rho(x, y) \\ &= \int_{B_R \times \partial W} |y| d\eta(x, y) = \int_{B_R} |\theta(x)| dF(\mu)(x). \end{aligned}$$

Since $|\theta_\rho(x)| dF(\mu_\rho)(x) = d|D\chi_{E_\rho}|(x)$ and $|\theta(x)| dF(\mu)(x) = |D\chi_{\bar{E}}|(x)$, this gives (14). \square

Remark 4.6. In the isotropic case $a(x)|\nu|$, Auer and Bangert proved a similar result [9, Th. 4.2]. In that case, the monotonicity formula directly implies the convergence of the blow-up to a cone which is calibrated by the constant \bar{z} and is thus a plane. For minimal surfaces this classically implies that $\bar{x} \in \partial^* E$.

Thanks to Proposition 3.4, one can order the minimizers which are calibrated by a given vector field.

Proposition 4.7. *Let $z \in X$ calibrates two plane-like minimizers E_1 and E_2 with connected boundaries. Then, either $E_1 \subseteq E_2$, or $E_2 \subseteq E_1$. As a consequence $\mathcal{H}^{d-3}(\partial E_1 \cap \partial E_2) = 0$.*

Proof. If z calibrates E_1 and E_2 then it also calibrates $E_1 \cap E_2$ and $E_1 \cup E_2$ which are then also Class A minimizers by Theorem 4.2. Since $E_1 \cap E_2 \subseteq E_1$, by Proposition 3.4, either $E_1 \cap E_2 = E_1$ in which case $E_1 \subseteq E_2$, or $\mathcal{H}^{d-3}(\partial(E_1 \cap E_2) \cap \partial E_1) = 0$ which implies that $E_2 \subseteq E_1$. \square

4.1 Calibrations and the Birkhoff property

We now define the class of plane-like minimizers that we are going to consider in the analysis of the differentiability properties of φ . If $E = \{x : v_p(x) + p \cdot x > s\}$ for some v_p which minimizes (6), we have that $E + q = \{v_p(x) + p \cdot x > s + p \cdot q\}$ for all $q \in \mathbb{Z}^d$, therefore $E = E + q$ if $p \cdot q = 0$, $E \subseteq E + q$ if $p \cdot q < 0$, and $E + q \subseteq E$ if $p \cdot q > 0$. This is called the *Birkhoff property*. Notice that in this case we also have $E = \bigcup_{\{q \cdot p > 0, q \in \mathbb{Z}^d\}} (E + q)$.

Definition 4.8. *Following [25, 35, 12] we give the following definitions:*

- we say that $E \subseteq \mathbb{R}^d$ satisfies the Birkhoff property if, for any $q \in \mathbb{Z}^d$, either $E \subseteq E + q$ or $E + q \subseteq E$;
- we say that E satisfies the strong Birkhoff property in the direction $p \in \mathbb{Z}^d$ if $E \subseteq E + q$ when $p \cdot q \leq 0$ and $E + q \subseteq E$ when $p \cdot q \geq 0$;
- we say that a plane-like minimizer E in the direction p is recurrent if either p is rational and E has the strong Birkhoff property, or if

$$E = \bigcup_{q \cdot p > 0, q \in \mathbb{Z}^d} (E + q) \quad \text{or} \quad E = \bigcap_{q \cdot p < 0, q \in \mathbb{Z}^d} (E + q). \quad (19)$$

Remark 4.9. Observe that if E satisfies the Birkhoff property, there exists $p \in \mathbb{R}^d$ such that if $q \in \mathbb{Z}^d$, $q \cdot p > 0$, then $E + q \subseteq E$, while $E + q \supseteq E$ if $q \cdot p < 0$ (the difference with the strong Birkhoff property is in the fact that when $q \cdot p = 0$, then one might not have $E + q = E$). The vector p (up to multiplication with a positive scalar) is uniquely determined, unless $E + q = E$ for all $q \in \mathbb{Z}^d$. See [12], or Lemma A.1 in Appendix A for an elementary proof of this claim.

Remark 4.10. A recurrent plane-like minimizer always enjoys the strong Birkhoff property (since the set $\bigcup_{q \cdot p > 0}(E + q)$, for instance, obviously does).

We will let $\mathcal{CA}(p)$ be the set of all the plane-like minimizers in the direction p which satisfy the strong Birkhoff property.

The following result can be deduced from [20].

Lemma 4.11. *If E is a Class A minimizer which satisfies the Birkhoff property, and $E \neq \emptyset$, $E \neq \mathbb{R}^d$, then it is plane-like in the direction given by Remark 4.9. Moreover it satisfies (5) with a constant M depending only on the anisotropy F and the dimension d .*

Proof. All the arguments can be found in the proofs of Proposition 8.3, Proposition 8.4 and Lemma 8.5 in [20]. First, if for some $a \in \mathbb{R}^d$, $a + [0, 1)^d \subseteq E$, then E contains the half-space $\{x \cdot p > a \cdot p + \sum_i |p_i|\} \subseteq \bigcup_{q \cdot p > 0}(q + a) + [0, 1)^d$, and similarly, if $(a + [0, 1)^d) \cap E = \emptyset$, then E is contained in a half-space.

Assume for instance that E does not contain a half-space, hence that $a + [0, 1)^d \cap E^c \neq \emptyset$ for all $a \in \mathbb{R}^d$. Then, by the density estimate, $|E^c \cap (a + [-1/2, 3/2)^d)| > \delta > 0$ for any a and some constant δ which depends only on c_0 and the dimension d . Now, we also have that $b + [0, 1)^d \cap E \neq \emptyset$ for some $b \in \mathbb{R}^d$, otherwise E would be empty. Then, for any $q \in \mathbb{Z}^d$ with $q \cdot p \geq 0$, $(q + b + [0, 1)^d) \cap E \subseteq (q + b + [0, 1)^d) \cap (q + E) \neq \emptyset$. Again it follows that $|(q + b + [-1/2, 3/2)^d) \cap E| \geq \delta > 0$. We deduce that the energy $\int_{q+b+[-1/2, 3/2)^d} F(x, D\chi_E)$ is bounded from below, by some constant $\delta' > 0$. Hence, if $B_R(x_R)$ is a large ball contained in $\{x : (x - b) \cdot p \geq 0\}$, the energy in the ball is bounded below by $N\delta'$, where $N := \#\{q \in \mathbb{Z}^d : q + b + [-1/2, 3/2)^d \subseteq B_R(x_R)\} \cong R^d$. However, by Class A minimality it is also less than $c_0^{-1} d\omega_d R^{d-1}$, a contradiction. It follows that E satisfies (5), with a constant M independent on E . \square

Proposition 4.12. *Let E be a Class A Minimizer with the Birkhoff property, then E has a periodic calibration.*

Proof. Let $R > 0$ and $k \geq 1$, and let

$$v_k(x) := \sum_{q \in \mathbb{Z}^d, |q| \leq k} \chi_{E+q} \in BV(B_R)$$

where in the sum, we drop the terms which are 1 a.e. on B_R . Thanks to the Birkhoff property, the sets $E + q$, $|q| \leq k$ are exactly the level sets of v_k . Consider now $v \in BV(B_R)$ such that $v - v_k$ has support in B_R . For $s \in \mathbb{R}$, the level set $\{v > s\}$ is a compactly supported perturbation of the level set $\{v_k > s\}$. Since this latter set is a Class A Minimizer, one has

$$\int_{B_R} F(x, D\chi_{\{v_k > s\}}) \leq \int_{B_R} F(x, D\chi_{\{v > s\}}). \quad (20)$$

Hence,

$$\begin{aligned} \int_{B_R} F(x, Dv) &= \int_{-\infty}^{\infty} \int_{B_R} F(x, D\chi_{\{v>s\}}) ds \\ &\geq \int_{-\infty}^{\infty} \int_{B_R} F(x, D\chi_{\{v_k>s\}}) ds = \int_{B_R} F(x, Dv_k), \end{aligned} \quad (21)$$

so that v_k is minimizing in B_R , with its own boundary datum. This yields the existence of a calibration $z_k^R \in L^\infty(B_R; \mathbb{R}^d)$, such that $F^\circ(x, z_k^R(x)) \leq 1$ a.e., $\operatorname{div} z_k^R = 0$, and $[z_k^R, Dv_k] = F(x, Dv_k)$ (in the sense of measures). By construction, the latter property is equivalent to $[z_k^R, D\chi_{E+q}] = F(x, D\chi_{E+q})$ for any $q \in \mathbb{Z}^d$ with $|q| \leq k$, that is, z_k^R is also a calibration for each minimizing set $E + q$, inside B_R .

Now, we let $k \rightarrow \infty$: up to a subsequence, z_k^R will converges, weakly-* in $L^\infty(B_R; \mathbb{R}^d)$, to some z_R which will be a calibration for all the sets $E+q$, $q \in \mathbb{Z}^d$, in the ball B_R (cf Prop. 4.3). Then we can send $R \rightarrow \infty$, in that case z_R (extended by zero out of B_R) converges again, weakly-* in $L^\infty(\mathbb{R}^d; \mathbb{R}^d)$, to some z which is a calibration for all the sets $E + q$, $q \in \mathbb{Z}^d$, in any ball. Let us now show that z may be chosen to be periodic: indeed, clearly, $z(x - q)$ is also a calibration for E and all its translates, for any $q \in \mathbb{Z}^d$. One may consider for any k

$$z'_k(x) := \frac{1}{\#\{q \in \mathbb{Z}^d : |q| \leq k\}} \sum_{q \in \mathbb{Z}^d, |q| \leq k} z(x - q). \quad (22)$$

which again, will be a calibration for E and all its translates. Passing to the limit, it converges (up to subsequences) to a new calibration z' , which is now periodic. \square

Proposition 4.13. *Let E be a plane-like minimizer with a periodic calibration, then E has the Birkhoff property and ∂E is connected.*

Proof. Without loss of generality we can assume that E satisfies (5) with $a = 0$. Since E has a periodic calibration, by Theorem 4.2 it is a Class A Minimizer. Moreover, since every connected component E is also calibrated, any of them is a Class A Minimizer. Let E_0 be the connected component of E which contains the half space $\{x \cdot p > M\}$ and let E_1 be another connected component of E . By Proposition 3.1, for every $R > 0$ we have $\mathcal{E}(E_1, B_R) \geq c_0 \beta R^{d-1}$. However, since $E_1 \subseteq \{|x \cdot p| \leq M\}$, the minimality of E_1 also yields $\mathcal{E}(E_1, B_R) \leq CM R^{d-2}$. This is a contradiction if R is large enough, and it follows that E is connected.

An analogous argument gives that E^c is also connected, thus implying the thesis. The Birkhoff property is deduced from Proposition 4.7, applied to E and $q + E$, $q \in \mathbb{Z}^d$. \square

From Lemma 4.11 and Propositions 4.12 and 4.13 we obtain the following

Corollary 4.14. *Let E be a Class A Minimizer with the Birkhoff property, then ∂E is connected.*

Remark 4.15. An interesting question raised by Bangert in [13] for non-parametric integrands is whether every plane-like minimizer necessarily satisfies the Birkhoff property. In [13, Th. 8.4], Bangert proves that, in the non-parametric case, it is true for totally irrational vectors p . Propositions 4.12 and 4.13 show that, in the parametric case, this question is equivalent to understand if every plane-like minimizer has a periodic calibration. See also [26] where a nice relation is given between this question of Bangert and De Giorgi's conjecture.

We also show the following result:

Proposition 4.16. *E is a recurrent plane-like minimizer in the direction p if and only if there exists a minimizer v_p of (6) such that*

$$E = \{x : v_p(x) + p \cdot x > 0\} \quad \text{or} \quad E = \{x : v_p(x) + p \cdot x \geq 0\}.$$

Proof. The “if part” is straightforward, as already observed.

If E is a recurrent plane-like minimizer (hence with the strong Birkhoff property), by Proposition 4.12 E has a periodic calibration z . We build a function v_k , $k \geq 1$, as follows: since E has the strong Birkhoff property, we can define in $BV_{\text{loc}}(\mathbb{R}^d)$ a function v_k such that $\{v_k \geq p \cdot q\} = E + q$ for all q with $|q| \leq k$. Indeed, E being plane-like, if $p \cdot q > 0$ one cannot have $E + q = E$, otherwise repeating the translation one would reach a contradiction. Hence the function

$$v_k(x) := \sup\{p \cdot q : |q| \leq k, x \in E + q\} \tag{23}$$

has actually the right level sets. Now, since E is plane-like, its oscillation is also uniformly bounded, and in fact it is locally uniformly bounded in L^∞ . By construction, z is a calibration for v_k , which means in particular that for any $R > 0$,

$$\int_{B_R} F(x, Dv_k) = \int_{B_R} [z, Dv_k] = - \int_{\partial B_R} v_k [z, \nu^{B_R}] \leq 2^d C_R c_0^{-1}$$

where C_R is a uniform bound for $\|v_k\|_{L^\infty(B_R)}$. Hence the v_k are uniformly bounded in $BV(B_R)$: up to a subsequence, we may assume that $v_k \rightarrow v$ in $L^1_{\text{loc}}(\mathbb{R}^d)$ with $v \in BV_{\text{loc}}(\mathbb{R}^d)$. If $|q| \leq k$, then $E + q \subseteq \{v_k \geq p \cdot q\}$. Passing to the limit it follows that $E + q \subseteq \{v \geq p \cdot q\}$. Conversely, if $(E + q)^c \subseteq \{v_k < p \cdot q\}$. Hence $(E + q)^c \subseteq \{v_k \leq p \cdot q\}$. Then, for any $q \in \mathbb{Z}^d$,

$$\{v > p \cdot q\} \subseteq E + q \subseteq \{v \geq p \cdot q\}. \tag{24}$$

If p is rational, then it is obvious that equality holds in (24), since, in fact, one can check that v_k does not change when k is large enough. If p is not rational, since E is recurrent we can assume that $E = \bigcup_{p \cdot q > 0} (E + q)$, and we shall prove that $E \subseteq \{v > 0\}$. This will imply that $E = \{v > 0\}$, and $E + q = \{v > p \cdot q\}$ for every $q \in \mathbb{Z}^d$.

If $x \in E$ then, for some $q \in \mathbb{Z}^d$ with $p \cdot q > 0$, we have $x \in E + q$ and thus, for $k \geq |q|$, we also have $v_k(x) \geq p \cdot q > 0$. Since the sequence $v_k(x)$ is increasing, we get $v(x) > 0$ and thus $E \subseteq \{v > 0\}$. The case $E = \bigcap_{p \cdot q < 0} (E + q)$ is similar and gives $E = \{v \geq 0\}$.

Let us show that $v - p \cdot x$ is periodic. It is enough to show that for almost every t and for all $q \in \mathbb{Z}^d$, we have $\{v \geq t\} + q \subseteq \{x : v \geq t + p \cdot q\}$. Then, letting $v_p(x) = v(x) - p \cdot x$, we will deduce that

$$v_p(x) \geq t - p \cdot x \implies v_p(x + q) \geq t + p \cdot q - p \cdot (x + q) = t - p \cdot x$$

for almost every t, x and for all $q \in \mathbb{Z}^d$, yielding that v_p is periodic (indeed, being \mathbb{Z}^d countable, the converse also holds for a.e. t and x).

For a.e. t we have $E_t = \{v \geq t\} = \{v > t\}$. In that case, E_t is the Kuratowski limit of $E + q_n$ for some sequence (q_n) in \mathbb{Z}^d , with $p \cdot q_n \rightarrow t$ as $n \rightarrow +\infty$. In particular, E_t is calibrated by z . Now, for k large enough and fixed n , $E + q_n + q = \{v_k \geq p \cdot (q_n + q)\}$, hence in the limit $E + q_n + q \subseteq \{v \geq t + p \cdot q\}$. Passing then to the limit in n we find that $\{v \geq t\} + q \subseteq \{v \geq t + p \cdot q\}$, which is our claim.

As already observed, almost all level sets of v are calibrated by z . It follows easily that v_p is a minimizer of (6). \square

An example of a non recurrent plane-like minimizer satisfying the strong Birkhoff property would be a set E with gaps on both side of its boundary, however we do not know whether this situation can occur.

Thanks to Proposition 4.16 we can now prove that every calibration calibrates every plane-like minimizer with the strong Birkhoff property, which implies that $\mathcal{CA}(p)$ is stable under union or intersection.

Theorem 4.17. *Let z be a calibration in the direction p , then z calibrates every plane-like minimizer with the strong Birkhoff property.*

Proof. By Proposition 4.16 we know that every recurrent plane-like minimizer is of the form $\{v_p + p \cdot x > s\}$ or $\{v_p + p \cdot x \geq s\}$, for some periodic function v_p minimizing (6), which implies that it is calibrated by every calibration.

We can thus assume that p is irrational and that we are given a non recurrent plane-like minimizer E , but which satisfies the strong Birkhoff property. Then, the set $\tilde{E} := \bigcup_{q \cdot p > 0} (E + q)$ is a plane-like recurrent Class A Minimizer (Proposition 3.2), hence calibrated by z . Moreover, it satisfies

$$\tilde{E} \subseteq E \quad \text{and} \quad E \subseteq \tilde{E} + q \quad \forall q \in \mathbb{Z}^d, \quad q \cdot p < 0. \quad (25)$$

Using the notation of Section 2.1, we define the quotient torus with respect to the rational directions orthogonal to p , $\mathbb{T}_r := \mathbb{R}^d / \Gamma(p)$.

If we are given a calibration z in the direction p , since it is periodic and since E and \tilde{E} are invariant by translations in $\Gamma(p)$, we can identify them with their equivalence class in \mathbb{T}_r . Thanks to (25), the measure $|E \setminus \tilde{E}|$ is then finite. Reminding that $V^r(p) = \text{Span}_{\mathbb{R}} \Gamma(p)$, we let for every $x \in \mathbb{R}^d$, $f(x) := \min_{x^r \in V^r(p)} |x - x^r|$. Then, the projection of f on \mathbb{T}_r

(still denoted by f) is well defined, and satisfies $|\nabla f| = 1$ a.e. in \mathbb{T}^r . Given $s \in \mathbb{R}$ we let $C_s := \{x \in \mathbb{T}^r : f(x) \leq s\}$. Then, it follows (from the Coarea Formula) that

$$|E \setminus \tilde{E}| = \int_0^\infty \mathcal{H}^{d-1}((E \setminus \tilde{E}) \cap \partial C_s) ds.$$

Hence, there exists a sequence $s_i \rightarrow +\infty$ such that

$$\mathcal{H}^{d-1}((E \setminus \tilde{E}) \cap \partial C_{s_i}) \leq \frac{1}{s_i}. \quad (26)$$

Now, Proposition 4.12 shows that every plane-like minimizer which satisfies the Birkhoff property is calibrated by a periodic calibration, and we can easily deduce that its projection in \mathbb{T}^r is also minimizing (with respect to compact perturbation inside the cylinder). In particular, comparing the energy of E with the energy of $(E \setminus C_{s_i}) \cup (\tilde{E} \cap C_{s_i})$, it follows from (26)

$$\int_{\partial^* E \cap C_{s_i}} F(x, \nu^E) \leq \int_{\partial^* \tilde{E} \cap C_{s_i}} F(x, \nu^{\tilde{E}}) + \frac{c_0^{-1}}{s_i}. \quad (27)$$

Integrating by parts on $(E \setminus \tilde{E}) \cap C_{s_i}$ and recalling that z calibrates \tilde{E} , we get (using (26) once more)

$$\int_{\partial^* E \cap C_{s_i}} [z, \nu^E] = \int_{\partial^* \tilde{E} \cap C_{s_i}} [z, \nu^{\tilde{E}}] - \int_{\partial C_{s_i} \cap (E \setminus \tilde{E})} [z, \nu^{C_{s_i}}] \geq \int_{\partial^* \tilde{E} \cap C_{s_i}} F(x, \nu^{\tilde{E}}) - \frac{c_0^{-1}}{s_i}. \quad (28)$$

Estimates (27), (28) yield

$$\int_{\partial^* E \cap C_{s_i}} F(x, \nu^E) \leq \int_{\partial^* E \cap C_{s_i}} [z, \nu^E] + 2 \frac{c_0^{-1}}{s_i}$$

and since $[z, \nu^E] \leq F(x, \nu^E)$ a.e. on $\partial^* E$, we easily deduce that in any C_R , we must have $\int_{\partial^* E \cap C_R} [z, \nu^E] = \int_{\partial^* E \cap C_R} F(x, \nu^E)$ so that z calibrates E . \square

Corollary 4.18. *For every $p \in \mathbb{R}^d \setminus \{0\}$, given E, F two plane-like minimizers in the direction p , if E has the Birkhoff property, and F the strong Birkhoff property, then either $E \subseteq F$ or $F \subseteq E$.*

Proof. It follows from Proposition 4.12, Corollary 4.14, Proposition 4.7 and Theorem 4.17. \square

Corollary 4.19. *For every $p \in \mathbb{R}^d \setminus \{0\}$, the plane-like minimizers of $\mathcal{CA}(p)$ form a lamination of \mathbb{R}^d (possibly with gaps).*

The following Lemma is reminiscent of [12, Lemma 4.4] and shows that for irrational vectors p , every recurrent plane-like minimizer is in the orbit of every other minimizer with the Birkhoff property.

Lemma 4.20. *Let $p \in \mathbb{R}^d \setminus (\mathbb{R} \cdot \mathbb{Z}^d)$ be an irrational vector. Then for every recurrent plane-like minimizer E satisfying*

$$E = \bigcup_{q \in \mathbb{Z}^d, q \cdot p > 0} (E + q) \quad (29)$$

and for every plane-like minimizer F with the Birkhoff property, there holds

$$E = \bigcup_{q \in \mathbb{Z}^d, q \cdot p > \alpha} (F + q) \quad (30)$$

where

$$\alpha := \inf\{q \cdot p : q \in \mathbb{Z}^d \text{ and } F + q \subseteq E\}.$$

In particular, for every u_p and v_p minimizing (6) and every $s, t \in \mathbb{R}$, there exists $\alpha \in \mathbb{R}$ such that $\{u_p + p \cdot x > t\} = \{v_p + p \cdot x > s + \alpha\}$. A similar result holds for the recurrent sets satisfying the second equality in (19).

Proof. Let E, F as above. By Corollary 4.18, for all $q \in \mathbb{Z}^d$, either $E \subseteq F + q$ or $F + q \subseteq E$. Notice that α is well-defined since E and F are plane-like (in the direction p). Let

$$\tilde{F} := \bigcup_{q \in \mathbb{Z}^d, q \cdot p > \alpha} (F + q) \subseteq E \quad (31)$$

and assume that the inclusion in (31) is strict. Thanks to (29), there exists $\bar{q} \in \mathbb{Z}^d$ with $\bar{q} \cdot p > 0$ and such that for every $q \in \mathbb{Z}^d$ with $p \cdot q > \alpha$ there holds

$$F + q \subseteq \tilde{F} \subseteq E + \bar{q} \subseteq E$$

and thus

$$F + (q - \bar{q}) \subseteq E$$

which contradicts the definition of α .

Applying this to the recurrent sets $\{u_p + p \cdot x > t\}$ and $\{v_p + p \cdot x > t + \alpha\}$ which both satisfy (29) and recalling that $\{v_p + p \cdot x > s + \alpha\} = \bigcup_{q \in \mathbb{Z}^d, p \cdot q > \alpha} \{v_p + p \cdot x > s\} + q$, we conclude the proof of the lemma. \square

As a consequence we get the following uniqueness result.

Theorem 4.21. *Let $p \in \mathbb{R}^d \setminus (\mathbb{R} \cdot \mathbb{Z}^d)$ be an irrational vector, then the minimizer v_p of (6) is unique up to an additive constant.*

Proof. Let u_p and v_p be two minimizers of (6). By Lemma 4.20 there exists $\alpha \in \mathbb{R}$ such that $\{v_p + p \cdot x > 0\} = \{u_p + p \cdot x > \alpha\}$. Then, for any $t \in \mathbb{R}$ there holds

$$\begin{aligned} \{v_p + p \cdot x > t\} &= \bigcup_{q \in \mathbb{Z}^d, q \cdot p > t} \{v_p + p \cdot x > 0\} + q = \bigcup_{q \in \mathbb{Z}^d, q \cdot p > t} \{u_p + p \cdot x > \alpha\} + q \\ &= \bigcup_{q \in \mathbb{Z}^d, q \cdot p > t + \alpha} \{u_p + p \cdot x > 0\} + q = \{u_p + p \cdot x > t + \alpha\}. \end{aligned}$$

It follows that $u_p = v_p + \alpha$. \square

Remark 4.22. In our context, the measure $Dv_p + p$ plays the role of Mather’s measures in Weak KAM Theory, and of the minimizing currents in the non-parametric setting. In that context, Bessi and Massart [16] proved that for irrational directions every non self-intersecting minimizer gives rise to the same minimizing current. In some sense their result is stronger than ours since the measure $Dv_p + p$ only accounts for the recurrent minimizers. In the same paper they also prove Mañé’s conjecture [27], namely that the uniqueness result generically holds also in the rational directions. See Appendix B for a similar result in our context.

From Theorem 4.17, Theorem 4.5 and Proposition 3.3 we have

Theorem 4.23. *Let $p \in \mathbb{R}^d \setminus \{0\}$ and $\Lambda := \bigcup \{\partial^* E : E \in \mathcal{CA}(p)\}$ then for every calibration $z \in X$ in the direction p we have*

$$z = \nabla_p F(x, \nu^E(x)) \quad \text{a.e. on } \Lambda,$$

where $\nu^E(x)$ is the normal to the plane-like minimizer passing through x . In particular, if v_p is a minimizer of (6), then z is prescribed almost everywhere in $\text{Spt}(|Dv_p + p|)$.

4.2 Heteroclinic surfaces

We consider now $p \in \mathbb{R}^d \setminus \{0\}$ a non totally irrational vector and let us assume that there exists $x \in \mathbb{R}^d$ such that no plane-like minimizer in the direction p , with the strong Birkhoff property, passes through x . Let

$$E^+ = \bigcap \{E : E \in \mathcal{CA}(p), x \in E\}, \quad E^- = \bigcup \{E : E \in \mathcal{CA}(p), x \notin E\}. \quad (32)$$

Then, there exists an open set $G = \text{int}(E^+) \setminus \overline{E^-}$, called a gap, which contains x , and through which no plane-like minimizer with the strong Birkhoff property passes.




Figure 1: Heteroclinic surface inside a gap.

We show that for any $q \in V^r(p)$, one can build inside G a heteroclinic surface which is “growing in the direction q ”, see Fig. 1. Let us remind that $\Gamma(p) = V^r(p) \cap \mathbb{Z}^d = \{\alpha \in \mathbb{Z}^d : \alpha \cdot p = 0\}$.

Definition 4.24. Let $p \in \mathbb{R}^d \setminus \{0\}$ be a non totally irrational vector and let $q \in V^r(p)$. If H is a plane-like minimizer in the direction p satisfying the Birkhoff property we will say that H is a heterocline in the direction q if

- for every $\alpha \in \Gamma(p)$ with $\alpha \cdot q = 0$, we have $H + \alpha = H$;
- for every $\alpha \in \Gamma(p)$ with $\alpha \cdot q > 0$, there holds $H + \alpha \subseteq H$.

Proposition 4.25. Let $p \in \mathbb{R}^d \setminus \{0\}$ be a non totally irrational vector, and assume that there exist E^\pm, G as above. Then, there exists a compact set $K \subseteq G$ such that, for every $q \in V^r(p)$, there exists a heterocline H_q in the direction q whose boundary intersects K . Moreover, letting

$$H_q^+ := \bigcup_{\alpha \in \Gamma(p)} (H_q + \alpha) \quad H_q^- := \bigcap_{\alpha \in \Gamma(p)} (H_q + \alpha) \quad (33)$$

we have

$$H_q^+ = E^+ \quad \text{and} \quad H_q^- = E^-. \quad (34)$$

Proof. Let $(\bar{q}_1, \dots, \bar{q}_k) \in \mathbb{Z}^d$ be an orthogonal basis of $V^r(p)$. We choose a (very) small $\eta > 0$ such that $\{x \in G : \text{dist}(x, \partial G) \geq \eta\} \neq \emptyset$, and let

$$K := \bigcap_{i=1}^k \{0 \leq x \cdot (\bar{q}_i / |\bar{q}_i|) \leq |\bar{q}_i|\} \cap \{x \in G : \text{dist}(x, \partial G) \geq \eta\}.$$

Following [13, 25, 35], the idea consists in letting $p_n = p + \frac{1}{n}q$ and considering a plane-like minimizer $E_{p_n} \in \mathcal{CA}(p_n)$ calibrated by a vector field z_n . Since E_{p_n} is included in a strip with normal p_n , up to an integer translation, we can assume that $\partial^* E_{p_n} \cap K \neq \emptyset$ for every n . Thus there exists a subsequence of E_{p_n} converging to a plane-like set H_q which is calibrated by $z := \lim z_n$, and such that $\partial H_q \cap K \neq \emptyset$ (by Hausdorff convergence in K of the boundaries). Moreover, z is a calibration in the direction p , since by weak-* convergence we also have

$$\left(\int_Q z \right) \cdot p = \varphi(p).$$

By construction H_q satisfies the Birkhoff property, is periodic in all the rational directions which are orthogonal to p and q , and $H + \alpha \subseteq H$ for all $\alpha \in \Gamma(p)$ with $\alpha \cdot q > 0$.

Define now H_q^\pm as in (33). It is clear that $H_q^\pm \in \mathcal{CA}(p)$ and thus, by definition of G , $H_q^- \subseteq E^- \subseteq E^+ \subseteq H_q^+$. If z is the calibration of H_q given above, z calibrates also H_q^\pm and E^\pm and thus, by the maximum principle, $H_q \subseteq E^+$. Therefore, $H_q + \alpha \subseteq E^+$ for every $\alpha \in \Gamma(p)$, from which we get $H_q^+ \subseteq E^+$. Similarly we have $E^- \subseteq H_q^-$, so that (34) is true. \square

Remark 4.26. The previous proposition asserts that the set H_q is really a heteroclinic solution, in the sense that it is a plane-like minimizer in the direction p which connects two periodic plane-like minimizers in the same direction.

Remark 4.27. When investigating the differentiability properties of φ at a point p in the direction q , it is natural to consider the heteroclinic minimizers constructed as above, which amounts to studying the asymptotic behavior of $\varphi(p + \frac{1}{n}q)$ as $n \rightarrow +\infty$.

We can prove the following uniform energy estimate for the heteroclinic surfaces:

Proposition 4.28. *There exists $\delta > 0$ and $R > 0$ such that, for every $q \in V^r(p)$ and every H_q, H_{-q} heteroclinic surfaces in the direction $q, -q$ respectively, such that $\partial^* H_q \cap \partial^* H_{-q} \cap K \neq \emptyset$, where K is the compact set given in Proposition 4.25, there holds*

$$\int_{B_R \cap \partial^* H_q} F(x, \nu^{H_q}) - [z, \nu^{H_q}] \geq \delta$$

for every $z \in X$ calibrating H_{-q} .

Proof. Assume on the contrary that there exists $R_n \rightarrow +\infty$ and $\delta_n \rightarrow 0$ such that there exists $q_n \in V^r(p)$, with $|q_n| = 1$, H_{q_n} heteroclinic in the direction q_n , and H_{-q_n} heteroclinic in the direction $-q_n$, calibrated by $z_n \in X$ and such that

$$\int_{B_{R_n} \cap \partial^* H_{q_n}} F(x, \nu^{H_{q_n}}) - [z_n, \nu^{H_{q_n}}] \leq \delta_n. \quad (35)$$

Since for every n , $\partial^* H_{\pm q_n}$ intersects K , there exists a subsequence such that $q_n \rightarrow q$, $H_{\pm q_n} \rightarrow H_{\pm}$, with $\partial H_+ \cap \partial H_- \cap K \neq \emptyset$ and z_n converge weakly-* to $z \in X$, where z calibrates H_- . By Proposition 3.2, Proposition 4.3 and Proposition 4.12, H_{\pm} are plane-like minimizers with the Birkhoff property. Even if it is not clear that H_{\pm} are heteroclinic, for any $\alpha \in \Gamma(p)$ one of the following properties hold

1. $H_+ + \alpha \subseteq H_+$ and $H_- \subseteq H_- + \alpha$
2. $H_+ \subseteq H_+ + \alpha$ and $H_- + \alpha \subseteq H_-$.

Arguing as in the proof of Proposition 4.3, we easily deduce from (35) that z is a calibration also for H_+ . By Proposition 4.7 we deduce that

$$H_- \subseteq H_+ \quad \text{or} \quad H_+ \subseteq H_-. \quad (36)$$

Assume for instance that $H_- \subseteq H_+$, and let $x \in \partial H_- \cap \partial H_+ \cap K$. Let also $(\alpha_1, \dots, \alpha_k) \in \Gamma(p)$ be an integer basis of vectors such that 1. holds: $H_+ + \alpha_i \subseteq H_+$ and $H_- \subseteq H_- + \alpha_i$ for $i = 1, \dots, k$. Observe that for any integer $n \geq 0$,

$$n\alpha_i + H_- \subseteq (n+1)\alpha_i + H_- \subseteq (n+1)\alpha_i + H_+ \subseteq H_+.$$

Hence, letting $\tilde{H} = \bigcup_{n_i \geq 0} (H_- + \sum_{i=1}^k n_i \alpha_i)$, we obtain a plane-like minimizer such that $\tilde{H} \subseteq H_+ \subseteq E_+$, $x \in \tilde{H}$, and \tilde{H} satisfies the strong Birkhoff property: hence $\tilde{H} = E^+$. It follows that $H_+ = E^+$, in contradiction with the fact that $K \cap \partial H_+ \neq \emptyset$. \square

5 Differentiability and strict convexity of φ

We are now ready to prove our main result.

Theorem 5.1. *Let $F \in C^{2,\alpha}(\mathbb{R}^d \times (\mathbb{R}^d \setminus \{0\}))$ be a convex, one-homogeneous and elliptic integrand. Then, the associated stable norm φ has the following properties:*

- φ^2 is strictly convex;
- if p is totally irrational then $\nabla\varphi(p)$ exists;
- the same occurs for any p such that the plane-like minimizers satisfying the strong Birkhoff property give rise to a foliation of \mathbb{R}^d ;
- if, on the other hand, these minimizers form a lamination with a gap, then $\partial\varphi(p)$ is a convex set of dimension $\dim(V^r(p))$, φ is differentiable in the directions of $(V^r(p))^\perp$ and is non-differentiable in the directions of $\mathbb{R}^d \setminus (V^r(p))^\perp$.

5.1 Strict convexity

Theorem 5.2. *The function φ^2 is strictly convex.*

Proof. Let p_1, p_2 , with $p_1 \neq p_2$, and let $p = p_1 + p_2$. We want to show that, if $\varphi(p) = \varphi(p_1) + \varphi(p_2)$, then p_1 is proportional to p_2 , which gives the thesis.

Indeed, we have

$$\begin{aligned} \varphi(p) &= \int_{\mathbb{T}} [z_p, p + Dv_p] \\ &= \int_{\mathbb{T}} F(x, p + Dv_p) \\ &\leq \int_{\mathbb{T}} F(x, p + Dv_{p_1} + Dv_{p_2}) \\ &\leq \int_{\mathbb{T}} F(x, p_1 + Dv_{p_1}) + F(x, p_2 + Dv_{p_2}) \\ &= \varphi(p_1) + \varphi(p_2). \end{aligned}$$

Since $\varphi(p) = \varphi(p_1) + \varphi(p_2)$, it follows that $v_{p_1} + v_{p_2}$ is also a minimizer of (6) and, in particular, z_p satisfies

$$[z_p, p_1 + Dv_{p_1}] + [z_p, p_2 + Dv_{p_2}] = F(x, p_1 + Dv_{p_1}) + F(x, p_2 + Dv_{p_2})$$

$(|p_1 + Dv_{p_1}| + |p_2 + Dv_{p_2}|)$ -a.e., so that

$$[z_p, p_i + Dv_{p_i}] = F(x, p_i + Dv_{p_i}) \quad i \in \{1, 2\}.$$

This means that z_p is a calibration for the plane-like minimizers

$$\{v_p + p \cdot x \geq s\}, \{v_{p_1} + p_1 \cdot x \geq s\} \text{ and } \{v_{p_2} + p_2 \cdot x \geq s\}$$

for all $s \in \mathbb{R}$. By Proposition 4.7, it follows that they are included in one another which is possible only if p_1 is proportional to p_2 . \square

Remark 5.3. Observe that $\varphi^2/2$ may fail to be uniformly convex. The thesis of the theorem is equivalent to the strict convexity of the level sets of φ .

5.2 Differentiability of φ

We now turn to the study of the differentiability of φ . As already noticed, the differentiability of φ at a point $p \in \mathbb{R}^d$ is equivalent to the fact that $\partial\varphi(p)$ is a singleton, that is, for every calibration $z \in X$ in the direction p the integral $\int_Q z \, dx$ has the same value.

Let us first show that φ must be differentiable in the totally irrational directions.

Proposition 5.4. *Assume p is totally irrational. Then for any two calibrations z, z' in the direction p , $\int_Q z \, dx = \int_Q z' \, dx$. As a consequence, $\partial\varphi(p)$ is a singleton and φ is differentiable at p .*

Proof. Consider z, z' two calibrations and a solution v_p of (6), and let $\xi = \int_Q z(x) \, dx$, $\xi' = \int_Q z'(x) \, dx$.

Let us show that, for any s ,

$$\int_{\{v_p + p \cdot x = s\}} z(x) - z'(x) \, dx = 0. \quad (37)$$

Thanks to the density estimate, the level sets $\{x : v_p(x) + p \cdot x = s\}$ are equivalent (up to a negligible set) to an open set C_s with $\partial C_s = \partial\{v_p + p \cdot x > s\} \cup \partial\{v_p + p \cdot x \geq s\}$. Moreover, all the C_s are empty except for a countable number of values. Consider such a value s . Since z and z' calibrate C_s which is a plane-like minimizer we have $[z, \nu^{C_s}] = [z', \nu^{C_s}]$ on $\partial^* C_s$.

Then, we observe that the sets $C_s^q = Q \cap (C_s - q)$, $q \in \mathbb{Z}^d$, are all disjoint, so that their measures sum up to less than 1. Indeed, if there is a point $y \in (C_s - q_1) \cap (C_s - q_2)$ with $q_1, q_2 \in \mathbb{Z}^d$, then we have $v_p(y + q_1) + p \cdot y + p \cdot q_1 = s = v_p(y + q_2) + p \cdot y + p \cdot q_2 = s$, but since $v_p(y + q_1) = v_p(y + q_2)$ it follows that $p \cdot (q_2 - q_1) = 0$, hence $q_1 = q_2$, since p is totally irrational.

For $R \in \mathbb{N}^*$, let Ψ_R be a Lipschitz cutoff function equal to 1 on $[-R, R]^d$, to 0 out of $[-(R+1), R+1]^d$, and with $|\nabla \Psi_R| = 1$ a.e. in $K_R = [-(R+1), R+1]^d \setminus [-R, R]^d$. Recalling that $\operatorname{div}(z - z') = 0$, we compute

$$\int_{C_s} \Psi_R(x)(z(x) - z'(x)) \cdot e_i \, dx = - \int_{C_s} x_i(z(x) - z'(x)) \cdot \nabla \Psi_R(x) \, dx, \quad (38)$$

which is bounded by

$$L_R = 2c_0^{-1}(R+1)|K_R \cap C_s|, \quad (39)$$

where c_0 is the constant in (3). Since $\sum_{R \geq 1} |K_R \cap C_s| \leq 1$, we get $\liminf_{R \rightarrow \infty} L_R = 0$, otherwise we would have $|K_R \cap C_s| \gtrsim c/(R+1)$ for large R and for some constant $c > 0$,

which would imply $\sum_R |K_R \cap C_s| = +\infty$. Hence, there exists a subsequence $R_k \rightarrow +\infty$ with $L_{R_k} \rightarrow 0$, but then, passing to the limit in (38) along this subsequence, we get

$$\int_{C_s} (z(x) - z'(x)) \cdot e_i dx = 0, \quad (40)$$

which gives our claim.

We deduce that $\xi = \xi'$. Indeed, from Theorem 4.23 it follows that

$$\int_{\mathbb{R}^d} (z - z') dx = \sum_s \int_{C_s} z - z' dx = 0,$$

where the sum is on all s such that C_s is an open, nonempty set. In particular, we obtain that $\int_Q (z - z') dx = \xi - \xi' = 0$. \square

If p is not totally irrational, by taking the quotient of \mathbb{R}^d with respect to all rational directions orthogonal to p , by the same argument we get the following result.

Corollary 5.5. *For every $q \in (V^r(p))^\perp$, the function φ is differentiable at p in the direction q . This amounts to say that, for every $\xi_1, \xi_2 \in \partial\varphi(p)$ and every $q \in (V^r(p))^\perp$,*

$$\xi_1 \cdot q = \xi_2 \cdot q.$$

As a direct consequence of Theorem 4.23 we also have

Proposition 5.6. *If the plane-like minimizers of $\mathcal{CA}(p)$ fibrate \mathbb{R}^d then φ is differentiable at p .*

We finally investigate the non-differentiability of φ at points p which are not totally irrational, and such that there is $x \in \mathbb{R}^d$ such that no minimizer passes through x . In that case let G be a gap containing x , bounded by two plane-like minimizers E^\pm .

We start by investigating what happens for a rational vector $q \in \Gamma(p) = V^r(p) \cap \mathbb{Z}^d$. For such a q , let H_q be a heteroclinic solution in the direction q with z_1 an associated calibration. Let also z_2 be a calibration associated to H_{-q} , the heteroclinic solution in the direction $-q$. We will prove that

$$\int_Q z_1 \cdot q \neq \int_Q z_2 \cdot q.$$

Let us notice that by Theorem 4.17 z_1 and z_2 have the same normal component on the boundaries of every gap, moreover by Theorem 4.5 they agree outside the gaps. Therefore, we can assume that z_1 and z_2 differ only inside the projection on the torus of the gap G , and thus we are reduced to prove that

$$\int_{Q \cap \Pi(G)} (z_1 - z_2) \cdot q \neq 0,$$

where $\Pi : \mathbb{R}^d \rightarrow Q$ denotes the projection on Q . We consider a further decomposition of the space $V^r(p)$. By a Gram-Schmidt procedure, we see that $V^r(p)$ is spanned by a family (q, q_2, \dots, q_k) of orthogonal vectors in \mathbb{Z}^d . Let $\tilde{\Gamma} := \text{Span}_{\mathbb{Z}}(q_2, \dots, q_k)$, and let $\tilde{\mathbb{T}}_r := \mathbb{R}^d / \tilde{\Gamma}$ be the cylinder quotient of \mathbb{R}^d and $\tilde{\Gamma}$. Since the gap G , the sets in $\mathcal{CA}(p)$, and the heteroclinic plane-like minimizers in direction q are periodic with respect to vectors in $\tilde{\Gamma}$, we identify them with their quotient with respect to the group action of $\tilde{\Gamma}$. Furthermore, since the q_i are orthogonal, the cylinder $\tilde{\mathbb{T}}_r$ can be identified with $\bigcap_{i=2}^k \left\{ x \cdot \frac{q_i}{|q_i|^2} \in [0, 1[\right\}$. Let

$$\mathbb{S}_s^t := \left\{ x \in \tilde{\mathbb{T}}_r : s < x \cdot \frac{q}{|q|^2} < t \right\}$$

and let $\mathbb{S} := \mathbb{S}_0^1$ be the unit slab in the direction q . We will show that

$$\int_{\mathbb{S} \cap G} (z_1 - z_2) \cdot q \neq 0. \quad (41)$$

Notice that there exist vectors $\alpha \in \tilde{\mathbb{T}}^r$ such that $\alpha \cdot q = 0$. However, there is only a finite number of vectors $\alpha \in \Gamma(p)$ such that $\alpha \in \bigcap_{i=2}^k \left\{ x \cdot \frac{q_i}{|q_i|^2} \in [0, 1[\right\}$ and $\alpha \cdot \frac{q}{|q|^2} \in [s, t]$ which implies that $|\mathbb{S}_s^t \cap G|$ is finite for every $(s, t) \in \mathbb{R}$. We also let

$$S_t := \left\{ x \in \tilde{\mathbb{T}}^r : x \cdot \frac{q}{|q|^2} = t \right\}$$

and $S := S_0$. Since

$$|\mathbb{S}_s^t \cap G| = |q| \int_s^t \mathcal{H}^{d-1}(G \cap S_\tau) d\tau,$$

the measure $\mathcal{H}^{d-1}(G \cap S_t)$ is finite for almost every $t \in \mathbb{R}$.

In particular, without loss of generality, we can assume that $\mathcal{H}^{d-1}(G \cap S)$ is finite.

Proposition 5.7.

$$\int_{\mathbb{S} \cap G} (z_1 - z_2) \cdot q = N \int_{Q \cap \Pi(G)} (z_1 - z_2) \cdot q$$

for some $N \in \mathbb{N}$, with $N \leq C|q| \prod_{i=2}^k |q_i|$ for some $C > 0$.

Proof. We will prove that $\int_{\mathbb{S} \cap G} (z_1 - z_2) \cdot q$ is an entire multiple of $\int_{Q \cap \Pi(G)} (z_1 - z_2) \cdot q$. Notice first that

$$\mathbb{S} = \left\{ x \cdot \frac{q}{|q|^2} \in [0, 1[\right\} \cap \bigcap_{i=2}^k \left\{ x \cdot \frac{q_i}{|q_i|^2} \in [0, 1[\right\}.$$

Moreover, we have

$$\mathbb{S} \cap G = \bigcup_{\alpha \in \mathbb{Z}^d : (Q + \alpha) \cap \mathbb{S} \neq \emptyset} (\mathbb{S} \cap G) \cap (Q + \alpha).$$

Let $\Gamma_{\mathbb{S}} := \{ \alpha \in \Gamma(p) : Q + \alpha \cap \mathbb{S} \neq \emptyset \}$. By the strong Birkhoff property, for every $\alpha_1, \alpha_2 \in \mathbb{Z}^d$, if $\alpha_1 \cdot p \neq \alpha_2 \cdot p$ then $(G + \alpha_1) \cap (G + \alpha_2) = \emptyset$, and for every $\alpha_1, \alpha_2 \in \Gamma_{\mathbb{S}}$, $G + \alpha_1 = G + \alpha_2$.

It is therefore sufficient to prove that, for some $N \in \mathbb{N}$, we have

$$\sum_{\alpha \in \Gamma_{\mathbb{S}}} \int_{\mathbb{S} \cap G \cap (Q + \alpha)} (z_1 - z_2) \cdot q = N \int_{Q \cap G} (z_1 - z_2) \cdot q.$$

Let $\alpha \in \Gamma_{\mathbb{S}}$ and $Q_\alpha := Q + \alpha$. If $Q_\alpha \cap \partial\mathbb{S} = \emptyset$, since $G + \alpha = G$, we have $G \cap \mathbb{S} \cap Q_\alpha = (G \cap Q) + \alpha$ and thus

$$\int_{G \cap \mathbb{S} \cap Q_\alpha} (z_1 - z_2) \cdot q = \int_{G \cap Q} (z_1 - z_2) \cdot q.$$

If $Q_\alpha \cap \partial\mathbb{S} \neq \emptyset$, we assume first that Q_α intersects $\partial\mathbb{S}$ only on one of the facets of \mathbb{S} . Then, there exists $\tilde{q} \in (q, q_2, \dots, q_k)$ such that

$$Q_\alpha \cap \mathbb{S} = Q_\alpha \cap \left\{ x \cdot \frac{\tilde{q}}{|\tilde{q}|^2} \geq 0 \right\} \quad \text{or} \quad Q_\alpha \cap \mathbb{S} = Q_\alpha \cap \left\{ x \cdot \frac{\tilde{q}}{|\tilde{q}|^2} < 1 \right\}.$$

Assume that the first possibility holds (see Figure 2), then \tilde{q} is such that

$$(Q_\alpha + \tilde{q}) \cap \mathbb{S} = (Q_\alpha + \tilde{q}) \cap \left\{ (x - \tilde{q}) \cdot \frac{\tilde{q}}{|\tilde{q}|^2} < 1 \right\} = (Q_\alpha + \tilde{q}) \cap \left(\left\{ x \cdot \frac{\tilde{q}}{|\tilde{q}|^2} < 1 \right\} + \tilde{q} \right).$$




Figure 2: The set \mathbb{S} of Proposition 5.7.

This shows that

$$(Q_\alpha \cap \mathbb{S}) \cap ((Q_\alpha + \tilde{q}) \cap \mathbb{S}) - q = \emptyset \quad \text{and} \quad (Q_\alpha \cap \mathbb{S}) \cup ((Q_\alpha + \tilde{q}) \cap \mathbb{S}) - q = Q_\alpha$$

whence

$$\int_{Q_\alpha \cap \mathbb{S} \cap G} (z_1 - z_2) \cdot q + \int_{(Q_\alpha + \tilde{q}) \cap \mathbb{S} \cap G} (z_1 - z_2) \cdot q = \int_{Q \cap G} (z_1 - z_2) \cdot q.$$

If Q_α intersects m facets of \mathbb{S} , then we can assume that, for some $\tilde{q}_1, \dots, \tilde{q}_m \in (q, q_2, \dots, q_k)$,

$$Q_\alpha \cap \mathbb{S} = \bigcap_{j=1}^m \left\{ x \cdot \frac{\tilde{q}_j}{|\tilde{q}_j|^2} \geq 0 \right\}.$$

We can then repeat the above argument by pairing the cube Q_α with the 2^m cubes of the form $Q_\alpha + \sum_{j=1}^m \delta_j \tilde{q}_j$, where δ_j takes only values 0 or 1. This proves that, for some $N \in \mathbb{N}$,

$$\sum_{\alpha \in \Gamma_s} \int_{S \cap G \cap (Q + \alpha)} (z_1 - z_2) \cdot q = N \int_{Q \cap G} (z_1 - z_2) \cdot q.$$

Moreover $N \leq \#\Gamma_s \leq C|q| \prod_{i=2}^k |q_i|$. \square

Proposition 5.8. *For almost every $s, t \in \mathbb{R}$, we have*

$$\int_{S_t \cap G} \left[z_1 - z_2, \frac{q}{|q|} \right] = \int_{S_s \cap G} \left[z_1 - z_2, \frac{q}{|q|} \right]. \quad (42)$$

In particular,

$$\int_{S \cap G} (z_1 - z_2) \cdot q = |q|^2 \int_{S \cap G} \left[z_1 - z_2, \frac{q}{|q|} \right]. \quad (43)$$

Proof. Fix $s < t \in \mathbb{R}$ such that $\mathcal{H}^{d-1}(G \cap S_t)$ and $\mathcal{H}^{d-1}(G \cap S_s)$ are finite. Let ψ_R be the same cutoff function as in Proposition 5.4, then

$$\begin{aligned} \int_{S_s^t \cap G} \operatorname{div}(\psi_R(z_1 - z_2)) &= \int_{S_s^t \cap G} \nabla \psi_R \cdot (z_1 - z_2) \\ &= \int_{S_t \cap G} \psi_R[z_1 - z_2, q] - \int_{S_s \cap G} \psi_R[z_1 - z_2, q]. \end{aligned}$$

As in Proposition 5.4, $\int_{S_s^t \cap G} \nabla \psi_R \cdot (z_1 - z_2) \rightarrow 0$ and thus letting $R \rightarrow \infty$ we find (42). By [7, Prop. 2.7] applied to $u(x) = q \cdot x$, we have

$$\int_{S \cap G} (z_1 - z_2) \cdot q = \int_0^{|q|^2} \left(\int_{S_t \cap G} \left[z_1 - z_2, \frac{q}{|q|} \right] \right) dt$$

which gives (43). \square

Proposition 5.9. *Let ν be the inward normal to H_q , then*

$$\int_{S \cap G} \left[z_1 - z_2, \frac{q}{|q|} \right] = \int_{\partial^* H_q \cap \tilde{\mathbb{T}}_r} [z_1 - z_2, \nu]. \quad (44)$$

Proof. We first introduce some additional notation (see Figure 3): let

$$\Sigma^+ := \partial^* H_q \cap \{x \cdot q > 0\} \cap \tilde{\mathbb{T}}_r \quad G^+ := G \cap \{x \cdot q > 0\} \cap H_q^c \cap \tilde{\mathbb{T}}_r$$

and

$$S^+ := S \cap G \cap H_q^c.$$

Then, letting ψ_R be the usual cutoff function, we have

$$\int_{G^+} \operatorname{div}(\psi_R(z_1 - z_2)) = \int_{S^+} \psi_R \left[z_1 - z_2, \frac{q}{|q|} \right] - \int_{\Sigma^+} \psi_R[z_1 - z_2, \nu]$$




Figure 3: Heteroclinic solution in the direction q .

As usual

$$\int_{G^+} \operatorname{div}(\psi_R(z_1 - z_2)) \rightarrow 0 \quad \text{and} \quad \int_{S^+} \psi_R \left[z_1 - z_2, \frac{q}{|q|} \right] \rightarrow \int_{S^+} (z_1 - z_2) \cdot \frac{q}{|q|}$$

On Σ^+ , $[z_1, \nu] = F(x, \nu) \geq [z_2, \nu]$ hence $\psi_R[z_1 - z_2, \nu]$ converges monotonically to $[z_1 - z_2, \nu]$ and thus passing to the limit when $R \rightarrow +\infty$ we get by the Monotone Convergence Theorem,

$$\int_{\Sigma^+} [z_1 - z_2, \nu] = \int_{S^+} \left[z_1 - z_2, \frac{q}{|q|} \right].$$

Similarly we define Σ^- and S^- and get

$$\int_{\Sigma^-} [z_1 - z_2, \nu] = \int_{S^-} \left[z_1 - z_2, \frac{q}{|q|} \right].$$

Summing these two equalities we find (44). \square

We then have :

Proposition 5.10. *There exists a constant C such that, if we take $q \in \Gamma(p)$ and we let $(q, q_2, \dots, q_k) \in \mathbb{Z}^q$ be an orthogonal basis of $V^r(p)$ constructed as above, then for every heteroclinic surface H_q in the direction q and every $z \in X$ which calibrates a heteroclinic surface H_{-q} in the direction $-q$ there holds*

$$\int_{\partial^* H_q \cap \tilde{\mathbb{T}}_r} F(x, \nu^{H_q}) - [z, \nu^{H_q}] \geq C \delta \prod_{i=2}^k |q_i|. \quad (45)$$

Proof. Let K , R and δ be as in Proposition 4.28, so that whenever $\partial^* H_q \cap \partial^* H_{-q} \cap K \neq \emptyset$ we have

$$\int_{\partial^* H_q \cap B_R} F(x, \nu^{H_q}) - [z, \nu^{H_q}] \geq \delta.$$

For every $\alpha \in \Gamma(p)$, the same estimate holds for the set $B_R + \alpha$. If $(\bar{q}_1, \dots, \bar{q}_k)$ is the basis of $V^r(p)$ that we have fixed from the beginning, letting $\Gamma_R := \operatorname{Span}_{\mathbb{Z}}(2[R]\bar{q}_1, \dots, 2[R]\bar{q}_k)$ we have $(B_R + \alpha) \cap (B_R + \beta) = \emptyset$ for $\alpha, \beta \in \Gamma_R$ and thus

$$\int_{\partial^* H_q \cap \tilde{\mathbb{T}}_r} F(x, \nu^{H_q}) - [z, \nu^{H_q}] \geq \sum_{\substack{\alpha \in \Gamma_R \\ B_R + \alpha \subseteq \tilde{\mathbb{T}}_r}} \int_{\partial^* H_q \cap (B_R + \alpha)} F(x, \nu^{H_q}) - [z, \nu^{H_q}].$$

The number of vectors $\alpha \in \Gamma_R$ such that $\partial^* H_q \cap \partial^* H_{-q} \cap (B_R + \alpha) \neq \emptyset$ and $B_R + \alpha \subseteq \tilde{\mathbb{T}}_r$, is proportional to $\prod_{i=2}^k |q_i|$. Since for every such α , there holds

$$\int_{\partial^* H_q \cap (B_R + \alpha)} F(x, \nu^{H_q}) - [z, \nu^{H_q}] \geq \delta,$$

we get (45). □

Proposition 5.11. *Let $q \in V^r(p) \setminus \{0\}$ then there exists ξ_1 and ξ_2 in $\partial\varphi(p)$ such that $\xi_1 \cdot q \neq \xi_2 \cdot q$. As a consequence, $\partial\varphi(p)$ is a convex set of dimension $\dim(V^r(p))$.*

Proof. Let us start with $q \in \Gamma(p) \setminus \{0\}$. Take $\xi_1 := \int_Q z_1$ and $\xi_2 := \int_Q z_2$. If $(q_2, \dots, q_k) \in \mathbb{Z}^d$ is such that (q, q_2, \dots, q_k) is an orthogonal basis of $V^r(p)$ then using the same notation as before, by Proposition 5.7,

$$\int_Q (z_1 - z_2) \cdot q = \int_{Q \cap \Pi(G)} (z_1 - z_2) \cdot q = \frac{1}{N} \int_{S \cap G} (z_1 - z_2) \cdot q$$

with $N \leq C|q| \prod_{i=2}^k |q_i|$. By (43) we get

$$\int_Q (z_1 - z_2) \cdot q = \frac{|q|^2}{N} \int_{S \cap G} \left[z_1 - z_2, \frac{q}{|q|} \right].$$

Equation (44) then yields

$$\int_Q (z_1 - z_2) \cdot q = \frac{|q|^2}{N} \int_{\partial^* H_q \cap \tilde{\mathbb{T}}_r} [z_1 - z_2, \nu^{H_q}] \geq \frac{1}{C \prod_{i=2}^k |q_i|} |q| \int_{\partial^* H_q \cap \tilde{\mathbb{T}}_r} [z_1 - z_2, \nu^{H_q}].$$

Using finally (45) we find

$$\int_Q (z_1 - z_2) \cdot q \geq C\delta|q|, \tag{46}$$

for some $C > 0$, which gives the desired result for $q \in \mathbb{Z}^d$. If now $q = \alpha/\beta \in \mathbb{Q}^d$ with $\alpha \in \mathbb{Z}^d$ and $\beta \in \mathbb{N}^*$ then applying (46) to βq , by homogeneity (46) holds also for q . Finally, every vector of $V^r(p) \setminus \{0\}$ can be approximated by rational vectors and since the lower bound of (46) is uniform, it passes to the limit. □

Theorem 5.1 now follows from Theorem 5.2, Proposition 5.4, Proposition 5.6, Proposition 5.11, recalling the fact that the directions of differentiability of φ at a point p form a linear subspace of \mathbb{R}^d .

6 Some examples

Whether or not gaps do occur in laminations by plane-like minimizers is a delicate question. In [11], Bangert proved that for every Riemannian metrics (i.e. if $F(x, p) = a(x)|p|$ and $g = 0$), for every periodic open set $V \subseteq \mathbb{R}^d$, it is possible to modify the function $a(x)$ inside

V in such a way that for every direction there always exists a gap. A simple example of a functional for which there are gaps in every direction can be constructed as follows. In dimension two, let $a(x) := c_1$ in a square D strictly contained in Q , and $a(x) := c_2$ outside D , with $c_1 > \sqrt{2}c_2$. Then, any plane-like minimizer must contain or avoid D , so that there are gaps in every directions. Notice that by making a little regularization, it is possible to have the same behavior for a functional satisfying the hypotheses of Theorem 5.1. In [38] the author constructs an example of a functional with gaps in every lamination by plane-like minimizers, by considering the prescribed mean curvature functional (i.e. $F(x, p) = |p|$ and $g \neq 0$) with g equal to some λ in a small ball and $-\lambda$ in another ball. In the next example, the function φ is differentiable in every direction except from one.

Let $d = 2$ and $F(x, y, \nu) = a(x)|\nu|$, with $a(x)$ a periodic function of the first variable (for example $a(x) = \sin(\frac{x}{2\pi}) + 2$). Then, for $p = \pm(1, 0)$, the only plane-like minimizers in the direction p are the planes orthogonal to p passing through the minima of $a(x)$. Thus φ is not differentiable at $\pm(1, 0)$ whereas for every other $p \in \mathbb{S}^1$, if E is a plane-like minimizer in the direction p , by invariance of the functional by translation along the y component, the set $E + t(0, 1)$ is also a plane-like minimizer for every $t \in \mathbb{R}$, therefore there is no gap in the direction p , and φ is differentiable at p (see Figure 4). Clearly, this construction can be done in any dimension.




Figure 4: Example with a gap for a translation invariant functional.

One can wonder how non differentiability of φ is related to the invariance by translation of the functional. The relation is not clear at all as shown by our last example where the energy is not invariant in any direction but for which the associated φ is differentiable in some rational direction.

Indeed, let ψ be a periodic smooth function on \mathbb{R}^d with $\|\nabla\psi\|_\infty \leq \frac{1}{2}$, let $u(x) := x_1 + \psi(x)$, where $x = (x_1, x')$, and let $z := \frac{e_1 + \nabla\psi(x)}{|e_1 + \nabla\psi(x)|}$ (the condition $\|\nabla\psi\|_\infty \leq \frac{1}{2}$ ensures that z is well defined). The vector field z is then normal to all level-sets of u , which give a fibration

of \mathbb{R}^d . Letting $g = \operatorname{div} z$, we see that z is a calibration of the level-sets of u for the energy

$$P(E) + \int_E g \, dx$$

and thus there level-sets give fibration of the space by plane-like minimizers in the direction e_1 , so that the corresponding φ is differentiable along e_1 .

7 G-closure of isotropic perimeter functionals

A natural question is to determine the set of anisotropies that one can obtain by homogenization of the isotropic interfacial energies

$$\mathcal{E}(E, A) = \int_{\partial^* E \cap A} a(x) d\mathcal{H}^{d-1}(x)$$

through the formula (6). If no bound is imposed on $a(x)$ (that is no restriction is made on the c_0 of (3)), we can easily see that the set of such interfacial energies is dense in the set of all symmetric anisotropies φ , i.e convex one-homogenous functions φ with $\varphi(-p) = \varphi(p)$. Indeed, it is sufficient to prove that we can obtain any crystalline energy with rational vertices (these are functions φ for which the unit ball $\{\varphi \leq 1\}$ is a polytope whose vertices are rational points). Let p_1, \dots, p_n be the rational vertices of a given convex symmetric polytope K . Let Π be the projection from \mathbb{R}^d on the torus. By approximation we can consider a function $a(x)$ defined by

$$a(x) = \begin{cases} \lambda_i & \text{if } x \in \Pi(\{p_i \cdot y = 0\}) \text{ for some } i, \\ +\infty & \text{otherwise} \end{cases}$$

where $\lambda_i = |p_i|^{\frac{1}{d-1}}$, so that for $p = p_i$ the plane-like minimizers are given by the half-spaces $\{p_i \cdot x > 0\} + q$ with $q \in \mathbb{Z}^d$, hence $\varphi(p_i) = 1$ and it follows that $\{\varphi \leq 1\} = K$. Notice that, when the hyperplanes $\{p_i \cdot x = 0\}$ cross, $a(x)$ is not well defined but, since the intersection is of dimension $d - 2$, it does not contribute to the energy. In Figure 5, we show the simplest example where φ is the L^1 norm, so that K is the cube with vertices $\pm e_i$ for e_i the canonical basis of \mathbb{R}^d .

By slightly modifying this construction, it is possible to approximate any even anisotropy φ with $c_0|p| \leq \varphi(p) \leq \frac{1}{c_0}|p|$ by isotropic interfacial energies \mathcal{E} with Lipschitz functions $a(x)$ satisfying the same bounds, $c_0 \leq a(x) \leq \frac{1}{c_0}$ (see [6]).

This kind of issues is very related to the famous G-closure problem for composite materials (see the recent paper [10] and the references therein) or, in a setting closer to ours, to the density with respect to Γ -convergence of Riemannian metrics in the set of all Finsler metrics [18].




Figure 5: Example of the construction of $\varphi = |\cdot|_1$. Left: the metric $a(x)$. Right: the unit ball K

A A discrete “separation” result

We show here the following result (see [12] for a slightly more complex proof):

Lemma A.1. *Assume E is a set which satisfies the Birkhoff property, that is, for any $q \in \mathbb{Z}^d$, either $q + E \subseteq E$ or $q + E \supseteq E$. Then there exists $p \in \mathbb{R}^d$, $|p| = 1$, such that for any $q \in \mathbb{Z}^d$, $q \cdot p > 0 \Rightarrow q + E \subseteq E$ and (obviously) $q \cdot p < 0 \Rightarrow q + E \supseteq E$. Moreover, p is unique, unless $E + q = E$ for all $q \in \mathbb{Z}^d$.*

Proof. Let $Z = \{q \in \mathbb{Z}^d : q + E \subseteq E\}$, clearly, $0 \in Z$, $Z + Z = Z$ (so that $nZ \subseteq Z$ for any $n \geq 0$), and the Birkhoff property states that for any $z \in \mathbb{Z}^d$, either $z \in Z$ or $-z \in Z$. Without loss of generality we may therefore assume that $e_i \in Z$ for $i = 1, \dots, d$, where (e_i) is the canonical basis of \mathbb{R}^d .

We claim that either $Z = \mathbb{Z}^d$, or the closed convex envelope of Z is not \mathbb{R}^d . In the latter case, since this envelope is a convex cone, it must be contained in a semispace, hence the result.

Assume, thus, that $Z \neq \mathbb{Z}^d$ but any point in \mathbb{Z}^d is in the convexification of Z . In particular, it must be that $-\sum_i e_i \notin Z$, otherwise, for any $p = (p_1, \dots, p_d) \in \mathbb{Z}^d$ we would obtain that given a nonnegative integer $n \geq -\min_i p_i$, $p = -n \sum_i e_i + \sum_i (p_i + n)e_i$ is also in Z .

A consequence is that as soon as $p_i < 0$ for all i , then $p \notin Z$, otherwise $p + \sum_i (-p_i - 1)e_i = -\sum_i e_i \in Z$, which gives another contradiction.

Now, by assumption, for any $\varepsilon > 0$ there exist $(p^k)_{k=1}^K, (\theta^k)_{k=1}^K$ with $p^k \in Z$, $\theta^k \geq 0$, $\sum_k \theta^k = 1$ such that

$$\left| \sum_{k=1}^K \theta^k p^k + \sum_{i=1}^d e_i \right| < \varepsilon.$$

Possibly changing infinitesimally the θ^k 's we can assume that they are rational, hence, $\theta^k = n^k/m$ for some integers $n^k \geq 0$, $m > 0$, $\sum_k n^k = m$.

It follows, letting $p = \sum_k n^k p^k \in Z$,

$$\left| p + m \sum_{i=1}^d e_i \right| < m\varepsilon.$$

As a consequence, for each i , $p_i < -m(1 - \varepsilon) \leq 0$ as soon as $\varepsilon < 1$, which implies that $p \notin Z$, a contradiction. Hence, the closed convex envelope of Z is strictly contained in \mathbb{R}^d . \square

B A generic uniqueness result

In Theorem 4.21 we have shown that the minimizer of (6) is unique up to an additive constant, if the direction p is irrational. In addition, we show here that it is generically unique when p is rational, that is, we prove a geometric counterpart of Mañé's conjecture [27].

For this we follow and adapt the proof of [14]. Contrary to the non-parametric case, it is no longer true that, if F is an admissible anisotropy and $f \in C^\infty(\mathbb{T})$, then $F(x, p) + f(x)|p|$ is also an admissible anisotropy. Indeed, if $\inf f \leq -c_0$, the function $F(x, p) + f(x)|p|$ is not coercive anymore. We will thus restrict ourselves to positive perturbations. For this reason we cannot directly use [14, Thm. 5] as in [16].

We will try to stay as close as possible to the notation of [14]. A set \mathcal{O} is called a residual set if it is a countable intersection of dense open sets. In a complete metric space, by Baire's Theorem, this implies that \mathcal{O} is itself dense.

Theorem B.1. *For every rational vector p , there exists a residual set $\mathcal{O}(p)$ of $E := C^\infty(\mathbb{T}) \cap \{f \geq 0\}$ such that for every $f \in \mathcal{O}(p)$, the minimizer of (6) for $F + f(x)|p|$ is unique up to an additive constant.*

Following an idea of Mather, we first rewrite (6) as a linear problem. Notice that, for $u \in BV(\mathbb{T})$, if we set $\mu_u := |Du + p| \otimes \delta_{\frac{Du+p}{|Du+p|}}$ (which is a Radon measure on $\mathbb{T} \times \mathbb{S}^{d-1}$) then

$$\int_{\mathbb{T}} F(x, Du + p) = \int_{\mathbb{T} \times \mathbb{S}^{d-1}} F(x, \nu) d\mu_u$$

is linear in μ_u . Let \tilde{H}_r be the set of measures μ_u for $u \in BV(\mathbb{T})$ with total variation less than r . Let H_r be the weak-* closure of the convex hull of \tilde{H}_r . By Banach-Alaoglu's Theorem, H_r is compact, convex and metrizable. Let F be the space of Borel measures on $\mathbb{T} \times \mathbb{S}^{d-1}$, G be the space of Borel measures on \mathbb{T} and $K_r \subseteq G$ be the metrizable compact convex set of Radon measures on \mathbb{T} with total variation less than r . Then, if $\pi : F \rightarrow G$ is the projection on the first marginal, for every $\mu_u \in \tilde{H}_r$, $\pi(\mu_u) = |Du + p| \in K_r$ and thus $\pi(H_r) \subseteq K_r$. Letting

$$MA(F, \mu) := \int_{\mathbb{T} \times \mathbb{S}^{d-1}} F(x, \nu) d\mu \quad \mu \in H_r,$$

we have that $MA(L, \cdot)$ is linear and continuous for the weak-* topology on H_r . If u is a minimizer of (6) for $F + f$ with $f \in E$, we have

$$\left(\frac{1}{c_0} + |f|_\infty\right) |p| \geq \int_{\mathbb{T}} F(x, Du + p) + f(x)|Du + p| \geq c_0 |Du + p|(\mathbb{T})$$

and thus for $r \geq (c_0^{-1} + |f|_\infty)|p|/c_0$ the measure μ_u is the minimizer of $MA(F + f, \cdot)$ in \tilde{H}_r . Since $MA(F + f, \cdot)$ is linear, the minimum over H_r and over \tilde{H}_r coincide. Hence, for every minimizing u , the measure μ_u is also a minimizer of $MA(F + f, \cdot)$ in H_r . We are thus going to prove that such minimizers in H_r are generically unique. Following [14], let $M_{H_r}(F) := \operatorname{argmin}_{\mu \in H_r} MA(F, \mu)$ and $M_{K_r}(F) := \pi(M_{H_r}(F))$.

Proposition B.2. *For every $r > 0$ there exists a residual set $\mathcal{O}_r \subseteq E$ such that for every $f \in \mathcal{O}_r$, the set $M_{K_r}(F + f)$ is reduced to a single element.*

Proof. Let now $\mathcal{O}(\varepsilon)$ be the set of points $f \in E$ such that $M_{K_r}(F + f)$ is contained in a ball of radius ε in K_r . We will prove that the Proposition holds for

$$\mathcal{O}_r := \bigcap_{\varepsilon > 0} \mathcal{O}(\varepsilon).$$

Indeed, if $f \in \mathcal{O}_r$, and if $M_{K_r}(F + f)$ is not a singleton then it is a convex set of positive dimension which would not be included in a ball of radius ε for ε small enough, contradicting the hypothesis $f \in \mathcal{O}_r$. It is thus enough to prove that for every $\varepsilon > 0$, the sets $\mathcal{O}(\varepsilon)$ are open and dense. \square

The fact that $\mathcal{O}(\varepsilon)$ is open is a direct consequence of the continuity of the map

$$(f, \mu) \rightarrow \int_{\mathbb{T} \times \mathbb{S}^{d-1}} F(x, \nu) + f(x) |\nu| d\mu$$

which implies that for every open subset U of H_r , the set $\{f \in E : M_{H_r}(F + f) \subseteq U\}$ is an open set of E and similarly for $M_{K_r}(F + f)$. The density argument is more involved. Let $w \in E$, we want to prove that w is in the closure of $\mathcal{O}(\varepsilon)$. Repeating verbatim the proof of [14, Lemma 7] there holds:

Lemma B.3. *There exists an integer m and a continuous map $T_m : K_r \rightarrow \mathbb{R}^m$*

$$T_m(\eta) := \left(\int_{\mathbb{T}} w_1 d\eta, \dots, \int_{\mathbb{T}} w_m d\eta \right)$$

with $w_i \in E$ and such that

$$\forall x \in \mathbb{R}^m \quad \operatorname{diam} T_m^{-1}(x) < \varepsilon$$

where the diameter is taken for the distance on K_r .

Define the function $\Lambda_m : \mathbb{R}^m \rightarrow \mathbb{R} \cup \{+\infty\}$ as

$$\Lambda_m(x) := \min_{\substack{\mu \in H_r \\ T_m \circ \pi(\mu) = x}} MA(F + w, \mu)$$

if $x \in T_m(\pi(H_r))$ and $+\infty$ otherwise. For $y = (y_1, \dots, y_m) \in \mathbb{R}^m$, let

$$M_m(y) := \operatorname{argmin}_{x \in \mathbb{R}^m} \Lambda_m(x) + y \cdot x$$

then for $y \in \mathbb{R}_+^m$

$$M_{K_r}(F + w + \sum_{i=1}^m y_i w_i) \subseteq T_m^{-1}(M_m(y)).$$

Letting $\mathcal{O}_m := \{y \in \mathbb{R}^m : M_m(y) \text{ is reduced to a point}\}$, we have from Lemma B.3

$$y \in \mathcal{O}_m \text{ and } y \in \mathbb{R}_+^m \implies w + \sum_{i=1}^m y_i w_i \in \mathcal{O}(\varepsilon)$$

it is thus enough to prove that 0 can be approximated by positive vectors of \mathcal{O}_m .

For this consider the convex conjugate of Λ_m ,

$$\begin{aligned} G_m(y) &:= \sup_{x \in \mathbb{R}^m} y \cdot x - \Lambda_m(x) \\ &= \sup_{\mu \in H_r} MA \left(\sum_{i=1}^m y_i w_i - w - F, \mu \right). \end{aligned}$$

Since H_r is compact, it is a convex and finite valued function which is then continuous on \mathbb{R}^m . Letting $\Sigma := \{y \in \mathbb{R}^m : \dim \partial G_m(y) \geq 1\}$ we have that $\dim \Sigma \leq m - 1$ (see [14, App. A] or [1]) and thus the complement of Σ is dense in \mathbb{R}^m . Since for every $y \in \mathbb{R}_+^m$ we have $M_m(y) = \partial G_m(-y)$, it follows that for every $y \in \mathbb{R}_+^m \cap \Sigma^c$ the set $M_m(y)$ is reduced to a point, which proves the claim.

We can finally end the proof of Theorem B.1. Let $\mathcal{O}(p) := \bigcap_{r>0} \mathcal{O}_r$ then by Baire's Theorem, $\mathcal{O}(p)$ is still a residual set of E . If $f \in \mathcal{O}(p)$ then for $r \geq (c_0^{-1} + |f|_\infty)|p|/c_0$, the set $M_{K_r}(F+f)$ is reduced to a single element and if u and v are two different minimizers of (6) for $F+f$, it follows that $|Du+p| = |Dv+p|$. For $s \in \mathbb{R}$ and $E_s := \{u+p \cdot x > s\}$, we can construct as in Proposition 4.16, a minimizer \tilde{u} such that the levelsets of $\tilde{u} + p \cdot x$ corresponds exactly to the projection $\Pi(\partial E_s)$ of ∂E_s in the torus \mathbb{T} . Therefore, the measure $|Du+p|$ is concentrated on $\Pi(\partial E_s)$ and since on ∂E_s there holds $\frac{Du+p}{|Du+p|} = \nu^{E_s}$, we find that $Du+p = D\tilde{u}+p$ and hence Du is unique. \square

Remark B.4. From this uniqueness result, it can be easily proved that every plane-like minimizer with the Birkhoff property generically induces the same current. Since it is not the main focus of our work and it would consist in repeating the arguments in [16], we just sketch the proof. For $p \in \mathbb{S}^{d-1}$, u a minimizer of the cell problem, E a plane-like minimizer with the Birkhoff property and $v \in C^\infty(\mathbb{T})$ a periodic vectorfield, define the currents T_u and T_E by

$$T_u(v) := \int_{\mathbb{T}} v \cdot (Du+p) \quad \text{and} \quad T_E(v) := \lim_{R \rightarrow +\infty} \frac{1}{\mathcal{H}^{d-1}(\partial B_R)} \int_{\partial^* E \cap B_R} v \cdot \nu^E d\mathcal{H}^{d-1},$$

where the limit defining T_E exists arguing as in [15]. If Du is unique and E is recurrent then $T_u = T_E$. Now, for every plane-like minimizer E with the Birkhoff property and every $q \in \mathbb{Z}^d$, we have $T_{E+k} = T_E$ (see [16, Lem. 3.1]). Thus, letting $\tilde{E} := \bigcap_{q \cdot p > 0} (E+q)$, there holds $T_{\tilde{E}} = T_E$ (see [16, Lem. 4.4]). Since \tilde{E} is recurrent, this implies that $T_E = T_u$ and therefore every plane-like minimizer with the Birkhoff property induces the same current T_u .

References

- [1] G. ALBERTI AND L. AMBROSIO, *A geometrical approach to monotone functions in \mathbb{R}^n* , Math. Z. 230 (1999), no. 2, 259–316.
- [2] F.J. ALMGREN, R. SCHOEN AND L. SIMON, *Regularity and singularity estimates on hypersurfaces minimizing elliptic variational integrals*, Acta Math., 139, 217-265, 1977.
- [3] L. AMBROSIO, G. CRIPPA AND S. MANIGLIA, *Traces and fine properties of a BD class of vector fields and applications*, Ann. de la Fac. des Sc. de Toulouse Ser. 6, 14, 4, 527-561, 2005.
- [4] L. AMBROSIO, N. FUSCO AND D. PALLARA, *Functions of bounded variation and free discontinuity problems*, Oxford Science Publications, 2000.
- [5] F. ANDREU-VAILLO, V. CASELLES AND J.M. MAZÒN, *Parabolic Quasilinear Equations Minimizing Linear Growth Functionals*, Birkhäuser, “Progress in Mathematics”, 223, 2004.
- [6] N. ANSINI AND O. IOSIFESCU, *Approximation of anisotropic perimeter functionals by homogenization*, Boll. Un. Mat. Ital. , 3, 149-168 , 2010.
- [7] G. ANZELLOTTI, *Pairings between measures and bounded functions and compensated compactness*, Annali di Matematica Pura ed Applicata, 135, 1, 293-318, 1983.
- [8] S. AUBRY, *The Devil’s staircase Transformation in incommensurate Lattices*, Lecture Notes in Math. 925, 221-245, 1982.
- [9] F. AUER AND V. BANGERT, *Differentiability of the stable norm in codimension one*, Amer. J. Math., 128, 1, 215-238, 2006.
- [10] J.F. BABADJIAN AND M. BARCHESI, *A variational approach to the local character of G -closure: the convex case*, Ann. IHP Analyse non linéaire, 26, 2, 351-373, 2009.
- [11] V. BANGERT, *The existence of gaps in minimal foliations*, Aequationes Math., 34, 153–166, 1987.
- [12] V. BANGERT, *A uniqueness theorem for \mathbb{Z}^n -periodic variational problems*, Comment. Math. Helvetici, 62, 511-531, 1987.
- [13] V. BANGERT, *On minimal laminations of the torus*, Ann. IHP Analyse non linéaire, 2, 95–138, 1989.
- [14] P. BERNARD AND G. CONTRERAS, *A generic property of families of Lagrangian systems*, Ann. of Math. (2) 167 (2008), no. 3, 1099–1108.

- [15] U. BESSI, *Aubry sets and the differentiability of the minimal average action in codimension one*, ESAIM Control Optim. Calc. Var., 15, 1, 1–48, 2009.
- [16] U. BESSI AND D. MASSART, *Mañé’s conjectures in codimension 1*, Comm. Pure Appl. Math., 64, 7, 1008–1027, 2011.
- [17] J. BOURGAIN AND H. BRÉZIS, *On the Equation $\operatorname{div} Y = f$ and Application to Control of Phases*, Jour. of Amer. Math. Soc., 16, 2, 393–426, 2002.
- [18] A. BRAIDES, G. BUTTAZZO AND I. FRAGALÀ, *Riemannian approximation of Finsler metrics*, Asymptotic Anal. , 31, 2, 177–187, 2002.
- [19] D. BURAGO, S. IVANOV AND B. KLEINER, *On the structure of the stable norm of periodic metrics*, Math. Res. Lett., 4, 791–808, 1997.
- [20] L. CAFFARELLI AND R. DE LA LLAVE, *plane-like Minimizers in Periodic Media*, Comm. Pure Appl. Math., 54, 1403–1441, 2001.
- [21] A. CHAMBOLLE AND G. THOUROUDE, *Homogenization of interfacial energies and construction of plane-like minimizers in periodic media through a cell problem*, Netw. Heterog. Media, vol. 4 n. 1, 127–152, 2009.
- [22] G.Q. CHEN, M. TORRES AND M. ZIEMER, *Gauss-Green theorem for weakly differentiable vector fields, sets of finite perimeter, and balance laws*, Comm. Pure Appl. Math., 62, 2, 242–304, 2009.
- [23] D. GILBARG AND N. TRUDINGER, *Elliptic partial differential equations of second order*, Classics in Mathematics. Springer-Verlag, 2001.
- [24] E. GIUSTI, *Minimal surfaces and functions of bounded variation*, Monographs in Mathematics 80, Birkhäuser, 1984.
- [25] H. JUNGINGER-GESTRICH, *Minimizing hypersurfaces and differentiability properties of the stable norm*, PhD Thesis, Freiburg, 2007.
- [26] H. JUNGINGER-GESTRICH AND E. VALDINOCI, *Some connections between results and problems of De Giorgi, Moser and Bangert*, Z. angew. Math. Phys., 60, 2009.
- [27] R. MAÑÉ, *Generic properties and problems of minimizing measures of Lagrangian systems*, Nonlinearity, 9, 273–310, 1996.
- [28] D. MASSART, *Stable norms of surfaces: local structure of the unit ball at rational directions*, Geom. Funct. Anal., 7, 996–1010, 1997.
- [29] J.N. MATHER, *Differentiability of the Minimal Average Action as a Function of the Rotation Number*, Bol. Soc. Bras. Mat., 21, 59–70, 1990.

- [30] F. MORGAN, *The cone over the Clifford torus in \mathbb{R}^4 is Φ -minimizing*, Math. Annalen, 289, 341–354, 1991.
- [31] J. MOSER, *Minimal solutions of variational problems on a torus*, Ann. IHP Analyse non linéaire, 3, 229-272, 1986.
- [32] J. MOSER, *A stability theorem for minimal foliations on a torus*, Ergo. Th. Dynam. Syst., 8, 251-281, 1988.
- [33] Y.G. RESHETNYAK, *Weak Convergence of Completely Additive Vector Functions on a Set*, Siberian Math. J., 9, 1968, translated from Sibirskii Matematicheskii Zhurnal, 9, 1386-1394, 1968.
- [34] R. SCHNEIDER, *Convex Bodies: The Brunn-Minkowski Theory*, Encyclopedia of Mathematics and its Applications, Cambridge university Press, 1993.
- [35] W. SENN, *Differentiability properties of the minimal average action*, Calc. Var. Partial Differential Equations, 3, 343–384, 1995.
- [36] W. SENN, *Strikte Konvexität für Variationsprobleme auf dem n -dimensionalen Torus*, Manuscripta Math. 71, 45-65, 1991.
- [37] L. SIMON, *A strict maximum principle for area minimizing hypersurfaces*, J. Diff. Geometry, 26, 327-335, 1987.
- [38] G. THOUROUDE, PhD Thesis, in preparation.
- [39] B. WHITE, *Existence of smooth embedded surfaces of prescribed topological type that minimize parametric even elliptic functionals on three-manifolds*, J. Diff. Geometry, 33, 413–443, 1991.