

APPROXIMATION OF THE RELAXED PERIMETER FUNCTIONAL UNDER A CONNECTEDNESS CONSTRAINT BY PHASE-FIELDS

PATRICK W. DONDL, MATTEO NOVAGA, BENEDIKT WIRTH, AND STEPHAN WOJTOWYTSCH

ABSTRACT. We develop a phase-field approximation of the relaxation of the perimeter functional in the plane under a connectedness constraint based on the classical Modica-Mortola functional and the connectedness constraint of [8]. We prove convergence of the approximating energies and present numerical results and applications to image segmentation.

CONTENTS

1. Introduction	1
2. Approximation by Connected Sets	2
2.1. The sharp interface model	2
2.2. The phase-field model	5
2.3. The sharp interface limit	6
2.4. Extensions and further observations	11
3. Numerical Results	13
Acknowledgements	15
Appendix A. Proof of Lemma 2.3	15
References	18

1. INTRODUCTION

In this article, we consider a phase field approximation of the problem of finding a “connected perimeter” of a set $E \subset \mathbb{R}^2$. This connected perimeter is given as the limit of the perimeter of an optimal sequence of connected sets approximating E in an L^1 -sense. An application of the phase field energy developed here may be the segmentation of a given image to yield a connected (or simply connected) set. Our functional is based on the classical Modica-Mortola energy with an additional energy term that penalizes non-path-connectedness of the preimage of a given interval under the phase field function. Similar to the methods in [4, 3] we use a geodesic distance in order to detect path-(dis)connectedness.

In [8, 7], this topological functional was introduced in the context of diffuse curvature dependent energies. Here, we show that the Γ -limit of the sum of the usual Modica-Mortola-energy and our topological energy (2.1) is given by the L^1 -relaxation of the perimeter functional when only

Date: October 14, 2018.

2010 *Mathematics Subject Classification.* 49J45, 51E10.

Key words and phrases. Phase field, connectedness, Modica-Mortola, Relaxation, Topological Constraint, Steiner tree.

considering connected approximating sets. In the sharp-interface setting, this relaxation has been studied in [5].

The article is structured as follows. In section 2.1 we recall the results of [5] for the sharp interface problem and construct the connection between our result on finite domains and the sharp interface problem posed in the plane. Section 2.2 contains an intuitive explanation of how our phase-field energy incorporates the connectedness constraint. The main result on the approximation of the connected relaxation of the perimeter functional by phase-field energies is stated and proved in section 2.3. Some extensions of the result concerning approximation by simply connected sets and relative perimeters are collected in section 2.4. Finally, we present numerical evidence for the effectiveness of our approach to connectedness for diffuse sets in section 3. A technical result on the approximation of the closest point projection onto a closed convex sets by non-expansive diffeomorphisms is proved in the appendix.

2. APPROXIMATION BY CONNECTED SETS

2.1. The sharp interface model. For open sets $E \subset \mathbb{R}^2$ such that the characteristic function χ_E of E is in $BV(\mathbb{R}^2)$, the perimeter functional

$$P(E) = |D\chi_E|(\mathbb{R}^2) = \sup \left\{ \int_E \operatorname{div}(u) \, dx \mid u \in C_c^1(\mathbb{R}^2), |Du| \leq 1 \right\}$$

is a generalized measure of the size of the boundary of E which agrees with the \mathcal{H}^1 -measure of the boundary on Lipschitz sets due to the Gauss-Green theorem. It is well known that P is lower semi-continuous under the L^1 -convergence of characteristic functions [2, Proposition 3.38], and that the characteristic functions of C^∞ -smooth open sets lie L^1 -dense in the collection of characteristic functions of sets of finite perimeter [2, Theorem 3.42]. Therefore, the L^1 -lower semi-continuous envelope of the perimeter functional without any additional constraints agrees with the functional itself, independently of whether the approximating sets in the relaxation process are taken to be smooth or not. In this article, we wish to consider a similar relaxation process, but under additional topological constraints.

For an open set $E \subset \mathbb{R}^2$ we define two relaxations of the perimeter functional under a connectedness constraint

$$\begin{aligned} \overline{P}_C(E) &:= \inf \left\{ \liminf_{n \rightarrow \infty} P(E_n) \mid E_n \xrightarrow{L^1} E, E_n \text{ indecomposable} \right\} \\ \overline{P}_C^r(E) &:= \inf \left\{ \liminf_{n \rightarrow \infty} P(E_n) \mid E_n \xrightarrow{L^1} E, E_n \text{ connected and } C^\infty\text{-smooth} \right\} \end{aligned}$$

which differ in the degree of smoothness required of the approximating sets. Here ‘indecomposable’ is a measure-theoretic analogue of the notion of connectedness for open sets which are only defined in the L^1 -sense. As usual, we take the L^1 -topology on equivalence classes of bounded measurable sets as induced by the metric given by the L^1 -distance of their characteristic functions or equivalently the Lebesgue measure of their symmetric difference $E\Delta F = (E \cup F) \setminus E \cap F$

$$d_{L^1}(E, F) = \int_{\mathbb{R}^2} |\chi_E - \chi_F| \, dx = |E\Delta F|.$$

Definition 2.1. [1] An open set U is called decomposable if there exist open sets U_1, U_2 such that $U = U_1 \cup U_2$ (in the L^1 -sense) and $P(U) = P(U_1) + P(U_2)$. It is called indecomposable if it is not decomposable.

Heuristically, a set is decomposable if we need not create significant new boundaries when cutting it into pieces. It was shown recently [5] that for (essentially) bounded sets $E \subset \mathbb{R}^2$ such that $\partial E = \partial_* E$ modulo sets of zero \mathcal{H}^1 -measure the identity

$$\overline{P_C}(E) = \overline{P_C^r}(E) = P(E) + 2 St(E)$$

holds where $St(E)$ is the length of the Steiner tree of \overline{E} , i.e.

$$St(E) = \inf \{ \mathcal{H}^1(K) \mid E \cup K \text{ connected} \}.$$

Above, \mathcal{H}^1 denotes the 1-dimensional Hausdorff measure on \mathbb{R}^2 and $\partial_* E$ is the essential boundary of \overline{E} , see [2, Definition 3.60]. For the existence of Steiner trees, their properties and regularity see [11].

In this article, we develop a phase-field energy functional which approximates the connected relaxation of the perimeter functional in the sense of Γ -convergence. For technical reasons, we prefer to work on a bounded domain Ω , so we introduce a similar relaxation in this setting:

$$\overline{P_{C,cc,\Omega}^r}(E) := \inf \left\{ \liminf_{n \rightarrow \infty} P(E_n) \mid E_n \xrightarrow{L^1} E, E_n \Subset \Omega, E_n \text{ connected and } C^\infty\text{-smooth} \right\}.$$

The notation $E \Subset \Omega$ signifies that $E \subset \overline{E} \subset \Omega$ and that \overline{E} is compact. In general, forcing sets to remain within Ω may force us to make longer connections than the \mathbb{R}^2 -Steiner tree which leads to $\overline{P_{C,cc,\Omega}^r}(E) \neq \overline{P_C}(E)$. We do not provide an explicit characterization of the lsc envelope in this case, nor do we discuss the relationship of other possible relaxations. If Ω is convex, on the other hand, then a connected set approximating E ‘gains nothing’ by leaving Ω , and $\overline{P_{C,cc,\Omega}^r} = \overline{P_C}$. We prove a slightly more general statement.

Lemma 2.2. *Assume that the convex hull of E is contained in Ω . Then $\overline{P_{C,cc,\Omega}^r}(E) = \overline{P_C}(E)$.*

Before we prove Lemma 2.2, we introduce a separate useful statement. We assume this to be well-known, but have been unable to find a reference for it.

Lemma 2.3. *Let $U \subset \mathbb{R}^n$ be a convex open set and $K \subset U$ compact. Then there exists a C^∞ -diffeomorphism $\phi : \mathbb{R}^n \rightarrow U$ such that*

- (1) $\phi(x) = x$ for all $x \in K$ and
- (2) $|\phi(x) - \phi(y)| \leq |x - y|$ for all $x, y \in \mathbb{R}^n$.

Since the proof of Lemma 2.3 is unrelated to the main points of this article, we have moved it to the appendix. Now we can prove Lemma 2.2.

Proof of Lemma 2.2. It is clear that

$$\overline{P_{C,cc,\Omega}^r}(E) \geq \overline{P_C}(E)$$

since fewer sets are admissible in the approximation process, so it suffices to prove the inverse inequality. Without loss of generality, we may also assume that $E \neq \emptyset$ and $P(E) < \infty$.

Step 1. Suppose that $0 \in \text{conv}(E)$ and denote $E_n := \frac{n-1}{n}E$. Since $\text{conv}(E)$ is open, there exists $r > 0$ such that $B_r(0) \in \text{conv}(E)$, thus if $x \in E_n$ and $y \in \mathbb{R}^2$ with $|y - x| < \frac{r}{n}$, then $\frac{n}{n-1}x \in E$ and thus

$$y = x + (y - x) = \left(1 - \frac{1}{n}\right) \left[\frac{n}{n-1}x \right] + \frac{1}{n} [n \cdot (y - x)] \in \text{conv}(\text{conv}(E)) = \text{conv}(E)$$

which shows that $E_n \Subset \text{conv}(E) \subset \Omega$. Furthermore, $P(E_n) = \frac{n-1}{n}P(E) \leq P(E)$, so by BV-compactness there exists a set $E_\infty \subset \text{conv}(E)$ such that $E_n \rightarrow E_\infty$ in L^1 (up to a subsequence).

Now if $x \in E$, there exists $r > 0$ such that $B_r(x) \subset E$ and thus in particular there exists $N \in \mathbb{N}$ such that

$$\frac{n}{n-1} y \in B_r(x) \subset E \quad \forall y \in B_{r/2}(x), n \geq N$$

since $f_n(x) = \frac{n}{n-1}x$ converges to the identity map locally uniformly on \mathbb{R}^n . Hence $B_{r/2}(x) \subset E_n$ for all $n \geq N$ and therefore also $B_{r/2}(x) \subset E_\infty$. In total, this implies that $E \subset E_\infty$, and since $|E_n| = \frac{n-1}{n} |E| \leq |E|$, it follows that $|E_\infty| \leq |E|$ which combines to the statement that $E_\infty = E$. The uniqueness of the limit shows that in fact $E_n \rightarrow E$ also without choosing a subsequence.

Assuming for the moment that $\overline{P_C^r}(E_n) \equiv \overline{P_{C,cc,\Omega}^r}(E_n)$ for all $n \in \mathbb{N}$, we take a sequence of smooth connected sets $E_n^k \Subset \Omega$ such that

$$\lim_{k \rightarrow \infty} E_n^k = E_n, \quad \lim_{k \rightarrow \infty} P(E_n^k) = \overline{P_{C,cc,\Omega}^r}(E_n) = \overline{P_C^r}(E_n)$$

and a diagonal sequence $E'_n = E_n^{k_n}$ such that $E'_n \rightarrow E$ and

$$\lim_{n \rightarrow \infty} P(E'_n) = \lim_{n \rightarrow \infty} \lim_{k \rightarrow \infty} P(E_n^k) = \lim_{n \rightarrow \infty} \overline{P_C^r}(E_n) = \lim_{n \rightarrow \infty} \frac{n-1}{n} \overline{P_C^r}(E) = \overline{P_C^r}(E)$$

whence $\overline{P_{C,cc,\Omega}^r}(E) \leq \lim_{n \rightarrow \infty} P(E'_n) = \overline{P_C^r}(E)$ and thus $\overline{P_{C,cc,\Omega}^r}(E) = \overline{P_C^r}(E)$.

Step 2. It remains to prove that $\overline{P_{C,cc,\Omega}^r}(E_n) = \overline{P_C^r}(E_n)$ holds for all $n \in \mathbb{N}$. Fix any $n \in \mathbb{N}$ and let E_n^k be a sequence of smooth connected subsets of \mathbb{R}^2 such that

$$\lim_{k \rightarrow \infty} E_n^k = E_n, \quad \lim_{k \rightarrow \infty} P(E_n^k) = \overline{P_C^r}(E_n).$$

Now let ϕ be a diffeomorphism as in Lemma 2.3 with $U = \text{conv}(E)$ and $K = \overline{E_n}$. Since ϕ is a C^∞ -diffeomorphism, $\phi(E_n^k)$ is also a connected open set with C^∞ -boundary, but additionally $\phi(E_n^k) \Subset \text{conv}(E)$. Since ϕ is 1-Lipschitz, we find that

$$P(\phi(E_n^k)) = \mathcal{H}^1(\partial\phi(E_n^k)) = \mathcal{H}^1(\phi(\partial E_n^k)) \leq \mathcal{H}^1(\partial E_n^k) = P(E_n^k)$$

and also

$$|\phi(E_n^k) \Delta E_n| = |\phi(E_n^k) \Delta \phi(E_n)| = |\phi(E_n^k \Delta E_n)| \leq |E_n^k \Delta E_n|$$

since $\phi = \text{id}$ on E_n . Thus $\phi(E_n^k) \rightarrow E_n$ in L^1 and

$$\overline{P_C^r}(E_n) = \lim_{k \rightarrow \infty} P(E_n^k) \geq \liminf_{k \rightarrow \infty} P(\phi(E_n^k)) \geq \overline{P_{C,cc,\Omega}^r}(E_n)$$

by the definition of the lower semi-continuous envelope. Since the opposite inequality is obvious, we find that $\overline{P_C^r}(E_n) = \overline{P_{C,cc,\Omega}^r}(E_n)$ which concludes the proof. \square

This result remains true if we consider the relaxation of the perimeter functional under approximation by *simply connected* sets. Also for this functional, an explicit characterization is available due to [5] as

$$\overline{P_{sc}}(E) = \overline{P_{sc}^r}(E) = P(E) + 2 \text{St}(E) + 2 \text{St}(E^c)$$

with notation analogous to the connected relaxation. We will briefly come back to this problem in Theorem 2.9.

Remark 2.4. Let $E \subset \mathbb{R}^3$ be the set given by

$$E = B_1(-Re_1) \cup B_1(Re_1)$$

for some $R \gg 1$. The sets

$$E_n = E \cup \left\{ (x, y, z) \in \mathbb{R}^3 \mid -R < x < R, y^2 + z^2 < \frac{1}{n^2} \right\}$$

are connected, open, have a Lipschitz boundary and satisfy $P(E_n) \rightarrow P(E)$. A slightly modified sequence of sets can be constructed to have C^∞ -boundaries. Thus, since we can connect two components of an open set with a tube of small volume and perimeter, we do not expect the relaxation of the perimeter functional under a connectedness constraint to exhibit any interesting behaviour in ambient spaces of dimension ≥ 2 .

2.2. The phase-field model. We choose the classical Modica-Mortola approximation [10]

$$\mathcal{F}_\varepsilon(u) = \frac{1}{c_0} \int_\Omega \frac{\varepsilon}{2} |\nabla u|^2 + \frac{1}{\varepsilon} W(u) \, dx$$

of the perimeter functional where $W(u) = u^2(u-1)^2$ is a double-well potential and c_0 is a normalising constant given by

$$c_0 = \int_0^1 \sqrt{2W(s)} \, ds.$$

To incorporate a connectedness constraint, we follow an idea developed by two of the authors for a problem of surfaces in a three-dimensional ambient space [8] based on a similar model for the two-dimensional Steiner problem and related questions [4]. Due to its novelty, we include a heuristic motivation here.

Recall that an open set E in \mathbb{R}^n is connected if and only if it is path-connected, so if and only if for every $x, y \in \mathbb{R}^n$ there exists a continuous curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$ such that $\gamma(0) = x$, $\gamma(1) = y$ and $\gamma(t) \in E$ for all $t \in [0, 1]$. It is well-known that we may assume that γ is smooth.

We introduce a quantitative notion of path-connectedness to generalize this concept. Let $F \in C^0(\mathbb{R}^n)$ be a function such that

$$F(x) = 0 \text{ if } x \in \overline{E}, \quad F(x) > 0 \text{ if } x \in \mathbb{R}^n \setminus \overline{E}$$

and define the geodesic distance

$$d^F(x, y) = \inf \left\{ \int_\gamma F \, d\mathcal{H}^1 \mid \gamma(0) = x, \gamma(1) = y, \gamma : [0, 1] \rightarrow \mathbb{R}^n \text{ Lipschitz} \right\}.$$

Then, if x, y lie in the same connected component of E (or rather, \overline{E}), then $d^F(x, y) = 0$, while if they lie in different connected components, we would expect them to be separated by a positive d^F -distance (at least in ‘nice’ cases). So we think of $d^F(x, y)$ as a quantitative measure of the path-disconnectedness of the set E at the points $x, y \in E$. To obtain a single number to measure the total path-disconnectedness of E , we can consider a double-integral

$$\int_E \int_E d^F(x, y) \, dx \, dy \quad \text{or} \quad \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \beta(x) \beta(y) d^F(x, y) \, dx \, dy$$

where $\beta : \mathbb{R}^n \rightarrow \mathbb{R}$ is a measurable function such that $\beta(x) = 0$ if $x \in \mathbb{R}^n \setminus E$ and $\beta(x) > 0$ if $x \in E$.

This can be adapted to a phase-field setting as follows. We want to approximate a set E by connected open sets E_n . Letting u denote the phase-field function approximating the characteristic function of E for some phase-field parameter ε , this corresponds to keeping the set $\{u \approx 1\}$ connected. More precisely, we choose $0 < s < 1/2$ and penalize the quantitative total disconnectedness of the set $\{1 - \varepsilon^s < u_\varepsilon\}$. So take Lipschitz-functions $\beta_\varepsilon, F_\varepsilon$ which are monotone increasing/decreasing respectively such that

$$\beta_\varepsilon(z) = \begin{cases} 0 & z \leq 1 - 2\varepsilon^s \\ 1 & z \geq 1 - \varepsilon^s \end{cases}, \quad F_\varepsilon(z) = \begin{cases} 1 & z \leq 1 - 2\varepsilon^s \\ 0 & z \geq 1 - \varepsilon^s \end{cases}.$$

Note that if γ is a Lipschitz-curve, we can take the trace of a $W^{1,2}$ -function u on γ , so that a geodesic distance with weight $F_\varepsilon(u)$ can be defined in the same way as above, albeit with a weight which is only non-negative, bounded and measurable on the curve. We introduce the ‘diffuse connectedness functional’

$$(2.1) \quad \mathcal{C}_\varepsilon(u) = \int_{\Omega} \int_{\Omega} \beta_\varepsilon(u(x)) \beta_\varepsilon(u(y)) d^{F_\varepsilon(u)}(x, y) dx dy$$

and the total energy of a phase-field

$$\mathcal{E}_\varepsilon : W_0^{1,2}(\Omega) \rightarrow \mathbb{R}, \quad \mathcal{E}_\varepsilon(u) = \mathcal{F}_\varepsilon(u) + \varepsilon^{-\kappa} \mathcal{C}_\varepsilon(u)$$

for some $\kappa > 0$ which measures the perimeter and penalizes disconnectedness.

Remark 2.5. We can allow different double-well potentials W , but we need to couple the parameter s in the choice $\beta_\varepsilon, F_\varepsilon$ to the order at which W vanishes at the potential wells.

2.3. The sharp interface limit. In this section we prove our main result, which essentially states that the functionals \mathcal{E}_ε approximate the relaxed connected perimeter.

Theorem 2.6. *Let $\Omega \subset \mathbb{R}^2$ be open and bounded. Then*

$$\left[\Gamma(L^1) - \lim_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon \right] (u) = \begin{cases} \overline{P_{C,cc,\Omega}^r}(\{u = 1\}) & \text{if } u \in BV(\overline{\Omega}, \{0, 1\}) \\ +\infty & \text{else.} \end{cases}$$

In particular, if Ω is convex and $P(E) = \mathcal{H}^1(\partial E)$ for $E = \{u = 1\}$, the Γ -limit is known to be

$$\overline{P_C}(E) = P(E) + 2 St(E)$$

due to Lemma 2.2 and [5, Theorems 3.4 and 3.7].

Proof of the lim sup-inequality. This construction is classical and thus we only sketch the proof. For more detailed arguments concerning the Modica-Mortola functional, see [10]. Let $u \in BV(\overline{\Omega}, \{0, 1\})$ and denote $E := \{u = 1\}$. We want to construct a sequence of phase-fields u_ε such that

$$\lim_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon) = \overline{P_{C,cc,\Omega}^r}(E).$$

Take a sequence of connected sets E_n such that

$$E_n \Subset \Omega, \quad \partial E_n \in C^\infty, \quad E_n \xrightarrow{L^1} E, \quad P(E_n) \rightarrow \overline{P_{C,cc,\Omega}^r}(E).$$

For every $n \in \mathbb{N}$, we may pick r_n such that the tubular neighbourhood

$$U_n = \{x \in \mathbb{R}^n \mid \text{dist}(x, \partial E_n) < r_n\}$$

is diffeomorphic to $\partial E_n \times (-r_n, r_n)$ via the map

$$\Phi : \partial E_n \times (-r_n, r_n) \rightarrow U_n, \quad \Phi(x, t) = x + t \nu_x.$$

Without loss of generality, we assume that the sequence r_n is strictly monotone decreasing to zero. For $r_{n+1}^2 \leq \varepsilon < r_n^2$, we insert the usual recovery sequence for E_n ,

$$u_\varepsilon(x) = q \left(\frac{\text{sdist}(x, \partial E_n)}{\varepsilon} \right) \cdot \chi_\varepsilon,$$

where q solves the 1-dimensional cell problem

$$q'' - W'(q) = 0, \quad q(-\infty) = 0, \quad q(0) = \frac{1}{2}, \quad q(+\infty) = 1,$$

$\text{sdist}(x, \partial E_n) = \text{dist}(x, E_n^c) - \text{dist}(x, E_n)$ is the signed distance function from ∂E_n chosen to be positive inside E_n and χ_ε is a cut-off function to ensure that $u_\varepsilon = 0$ close to $\partial\Omega$. Then also the set $\{u_\varepsilon > 1 - \varepsilon^s\}$ is diffeomorphic to E_n , thus connected, and $\mathcal{C}_\varepsilon(u_\varepsilon) \equiv 0$. It is well-known that

$$\mathcal{F}_\varepsilon(u_\varepsilon) - P(E_n) \rightarrow 0$$

as $\varepsilon \rightarrow 0$ (where $n = n_\varepsilon$ is the corresponding index), so in total we have shown that $\mathcal{E}_\varepsilon(u_\varepsilon) \rightarrow \overline{P_{C,\Omega}}(E)$ as required. \square

Proof of the lim inf-inequality. Preliminaries and heuristics. Let $u_\varepsilon \in W_0^{1,2}(\Omega)$ be a sequence of functions such that $u_\varepsilon \rightarrow u$ in $L^1(\Omega)$. Without loss of generality we may assume that $\liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon) < \infty$. As for the Modica-Mortola functional, this implies that u is the characteristic function of a set of finite perimeter in $\overline{\Omega}$, so we need to show that

$$\liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon) \geq \overline{P_{C,cc,\Omega}^r}(\{u = 1\}).$$

We denote $E := \{u = 1\}$. Since the energy $\mathcal{E}_\varepsilon(u_\varepsilon)$ decreases when we truncate u_ε from above at 1 and from below at 0, we may assume that $u_\varepsilon \in W_0^{1,2}(\Omega, [0, 1])$, and by the density of smooth functions even that $u_\varepsilon \in C_c^\infty(\Omega, [0, 1])$.

Let $0 < \delta < \frac{c_0}{2}$ and consider the primitive function $G(z) = \int_0^z \sqrt{2W(s)} \, ds$. Using the co-area formula for BV -functions we obtain that

$$\begin{aligned} \int_\Omega \frac{\varepsilon}{2} |\nabla u_\varepsilon|^2 + \frac{W(u_\varepsilon)}{\varepsilon} \, dx &\geq \int_\Omega \sqrt{2W(u_\varepsilon)} |\nabla u_\varepsilon| \, dx \\ &= \int_\Omega |\nabla G(u_\varepsilon)| \, dx \\ &= \int_0^{c_0} \int_\Omega d|\nabla \chi_{\{G(u_\varepsilon) > t\}}| \, dt \end{aligned}$$

for χ_A the characteristic function of a set A , so there exists $t_\varepsilon \in (\delta, c_0 - \delta)$ such that

$$\int_\Omega d|\nabla \chi_{\{G(u_\varepsilon) > t_\varepsilon\}}| \leq \frac{1}{c_0 - 2\delta} \int_\Omega \frac{\varepsilon}{2} |\nabla u_\varepsilon|^2 + \frac{W(u_\varepsilon)}{\varepsilon} \, dx.$$

Since almost all t are regular values of $G(u_\varepsilon)$, we can even pick t_ε such that $\{G(u_\varepsilon) = t_\varepsilon\}$ is a C^∞ -submanifold of \mathbb{R}^2 . Since the level set is additionally closed and bounded, we see that $\{G(u_\varepsilon) = t_\varepsilon\}$ is a compact manifold – in particular, $\{G(u_\varepsilon) = t_\varepsilon\}$ has only a finite number M_ε of connected components. Since $\nabla G(u_\varepsilon)$ does not vanish on $\{G(u_\varepsilon) = t_\varepsilon\}$ by assumption and since $u_\varepsilon = 0 < t_\varepsilon$ on $\partial\Omega$, we see that

$$\{G(u_\varepsilon) = t_\varepsilon\} = \partial\{G(u_\varepsilon) > t_\varepsilon\}.$$

Now, let us go through the heuristic of the proof: If we could show that $\{G(u_\varepsilon) > t_\varepsilon\}$ were connected, we would be done, arguing that

$$\{G(u_\varepsilon) > t_\varepsilon\} = \{u_\varepsilon > G^{-1}(t_\varepsilon)\} \xrightarrow{L^1} \{u = 1\}$$

and then concluding that

$$\begin{aligned} \overline{P_{C,cc,\Omega}^r}(E) &\leq \liminf_{\varepsilon \rightarrow 0} P(\{G(u_\varepsilon) > t_\varepsilon\}) \\ &\leq \liminf_{\varepsilon \rightarrow 0} \frac{1}{c_0 - 2\delta} \int_\Omega \frac{\varepsilon}{2} |\nabla u_\varepsilon|^2 + \frac{W(u_\varepsilon)}{\varepsilon} \, dx \\ &\leq \frac{c_0}{c_0 - 2\delta} \liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon) \end{aligned}$$

for all $0 < \delta < \frac{c_0}{2}$. Then, taking $\delta \rightarrow 0$, we would obtain the Γ – lim inf-inequality

$$\overline{P_{C,cc,\Omega}^r}(E) \leq \liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon).$$

In general, there is no good reason for $\{G(u_\varepsilon) > t_\varepsilon\}$ to be connected since the super-level set is highly sensitive to very slight perturbations which are barely visible in the Modica-Mortola energy – however, the energy contribution of C_ε prevents the set $\{u_\varepsilon > 1 - \varepsilon^s\}$ from being ‘too disconnected’, so we can take a slight modification of the set which barely changes area or perimeter, but makes it connected.

Step 1. In this step, we show that

$$\lim_{\varepsilon \rightarrow 0} E_\varepsilon = E$$

in the L^1 -topology of open sets for all sets E_ε such that

$$\{u_\varepsilon > 1 - \varepsilon^s\} \subset E_\varepsilon \subset \{u_\varepsilon > \varepsilon^s\}.$$

In particular we note that

$$\{u_\varepsilon > 1 - \varepsilon^s\} \subset \{u_\varepsilon > G^{-1}(t_\varepsilon)\} \subset \{u_\varepsilon > \varepsilon^s\}$$

since $\delta < t_\varepsilon < c_0 - \delta$, so $G^{-1}(t_\varepsilon)$ is bounded away from 0 and 1. Second, we note that

$$|\{\varepsilon^s < u_\varepsilon < 1 - \varepsilon^s\}| \leq \frac{\varepsilon}{W(\varepsilon^s)} \int_{\Omega} \frac{W(u_\varepsilon)}{\varepsilon} dx \leq \frac{C\varepsilon}{\varepsilon^{2s}} = C\varepsilon^{1-2s} \rightarrow 0,$$

where C is an energy bound uniform in ε , so that any set containing $\{u_\varepsilon > 1 - \varepsilon^s\}$ and contained in $\{u_\varepsilon > \varepsilon^s\}$ has the same L^1 -limit (if one of them exists). Here we use that W vanishes quadratically at the potential wells, for other double-well potentials, other s may be admissible. Now observe that

$$\begin{aligned} \|u_\varepsilon - \chi_{\{u_\varepsilon > 1 - \varepsilon^s\}}\|_{L^1} &= \int_{\{u_\varepsilon > 1 - \varepsilon^s\}} |u_\varepsilon - 1| dx + \int_{\{\varepsilon^s < u_\varepsilon < 1 - \varepsilon^s\}} |u_\varepsilon| dx + \int_{\{u_\varepsilon < \varepsilon^s\}} |u_\varepsilon| dx \\ &\leq \varepsilon^s \cdot |\Omega| + C\varepsilon^{1-2s} + \varepsilon^s \cdot |\Omega| \rightarrow 0 \end{aligned}$$

so that u_ε and $\chi_{\{u_\varepsilon > 1 - \varepsilon^s\}}$ have the same L^1 -limit $u = \chi_E$, in other words

$$\lim_{\varepsilon \rightarrow 0} \{u_\varepsilon > 1 - \varepsilon^s\} = E$$

in the L^1 -topology.

Step 2. In this step, we eliminate the connected components of the approximating set which we deem too small to matter. Denote the connected components of $\{G(u_\varepsilon) > t_\varepsilon\}$ by $U_{i,\varepsilon}$, $i = 1, \dots, M_\varepsilon$, where the components are ordered by volume:

$$|U_{1,\varepsilon}| \geq \dots \geq |U_{M_\varepsilon,\varepsilon}|.$$

Denote

$$E_\varepsilon = \{G(u_\varepsilon) > t_\varepsilon\} = \bigcup_{i=1}^{M_\varepsilon} U_{i,\varepsilon}.$$

Applying the iso-perimetric inequality, we observe that for $N < M_\varepsilon$ we have

$$\begin{aligned} \left| \bigcup_{i=N}^{M_\varepsilon} U_{i,\varepsilon} \right| &= \sum_{i=N}^{M_\varepsilon} |U_{i,\varepsilon}| \leq \sqrt{|U_{N,\varepsilon}|} \sum_{i=N}^{M_\varepsilon} \sqrt{|U_{i,\varepsilon}|} \leq \sqrt{|U_{N,\varepsilon}|} \sum_{i=N}^{M_\varepsilon} \sqrt{\frac{1}{4\pi} \text{Per}(U_{i,\varepsilon})^2} \\ &= \sqrt{\frac{|U_{N,\varepsilon}|}{4\pi}} \sum_{i=N}^{M_\varepsilon} \text{Per}(U_{i,\varepsilon}) = \sqrt{\frac{|U_{N,\varepsilon}|}{4\pi}} \text{Per} \left(\bigcup_{i=N}^{M_\varepsilon} U_{i,\varepsilon} \right) \leq \sqrt{|U_{N,\varepsilon}|} \frac{\text{Per}(E_\varepsilon)}{\sqrt{4\pi}}, \end{aligned}$$

so if $U_{N,\varepsilon}$ carries little mass, then all the remaining components together have little mass as well. The identity

$$\sum_{i=N}^{M_\varepsilon} \text{Per}(U_{i,\varepsilon}) = \text{Per} \left(\bigcup_{i=N}^{M_\varepsilon} U_{i,\varepsilon} \right)$$

holds easily since E_ε is a smooth set whose boundary has finitely many connected components. Choose $N_\varepsilon \leq M_\varepsilon$ such that

$$|U_{i,\varepsilon}| \geq \frac{1}{|\log \varepsilon|} \quad \forall i = 1, \dots, N_\varepsilon, \quad |U_{N_\varepsilon+1,\varepsilon}| < \frac{1}{|\log \varepsilon|}.$$

It may happen that $N_\varepsilon = M_\varepsilon$ – this will rather simplify the proof, so we do not consider that case. We take

$$E'_\varepsilon = \bigcup_{i=1}^{N_\varepsilon} U_{i,\varepsilon}$$

and note that $P(E'_\varepsilon) \leq P(E_\varepsilon)$ (since we only remove boundary components) and still $E'_\varepsilon \rightarrow E$ in L^1 .

Step 3. In this step, we show that it is possible to connect all the components $U_{i,\varepsilon}$ of E'_ε without changing the L^1 -limit or increasing the perimeter by much. We note that the number of large connected components cannot increase too quickly in ε since

$$\frac{N_\varepsilon}{|\log \varepsilon|} \leq N_\varepsilon |U_{N_\varepsilon,\varepsilon}| \leq \sum_{i=1}^{N_\varepsilon} |U_{i,\varepsilon}| = |E'_\varepsilon| \leq |\Omega| \quad \Rightarrow \quad N_\varepsilon \leq |\Omega| |\log \varepsilon|.$$

Furthermore, we note that for $i \leq N_\varepsilon$

$$|U_{i,\varepsilon} \cap \{u_\varepsilon > 1 - \varepsilon^s\}| \geq |U_{i,\varepsilon}| - |\{x \in \Omega \mid \varepsilon^s < u_\varepsilon < 1 - \varepsilon^s\}| \geq |\log \varepsilon|^{-1} - C \varepsilon^{1-2s} \geq \frac{1}{2|\log \varepsilon|}$$

for all small enough $\varepsilon > 0$, so that within each connected component $U_{i,\varepsilon}$ we have a large volume on which u_ε is between $1 - \varepsilon^s$ and 1. Now let $2 \leq i \leq N_\varepsilon$. Then we know that

$$\begin{aligned} C &\geq \varepsilon^{-\kappa} \int_{\Omega} \int_{\Omega} \beta_\varepsilon(u_\varepsilon(x)) \beta_\varepsilon(u_\varepsilon(y)) d^{F_\varepsilon(u_\varepsilon)}(x, y) dx dy \\ &\geq \varepsilon^{-\kappa} \int_{U_{i,\varepsilon} \cap \{u_\varepsilon > 1 - \varepsilon^s\}} \int_{U_{1,\varepsilon} \cap \{u_\varepsilon > 1 - \varepsilon^s\}} d^{F_\varepsilon(u_\varepsilon)}(x, y) dx dy \\ &= \varepsilon^{-\kappa} |U_{i,\varepsilon} \cap \{u_\varepsilon > 1 - \varepsilon^s\}| |U_{1,\varepsilon} \cap \{u_\varepsilon > 1 - \varepsilon^s\}| \text{dist}^{F_\varepsilon(u_\varepsilon)}(U_{1,\varepsilon}, U_{i,\varepsilon}) \\ &\geq \frac{1}{4|\log \varepsilon|^2 \varepsilon^\kappa} \text{dist}^{F_\varepsilon(u_\varepsilon)}(U_{1,\varepsilon}, U_{i,\varepsilon}) \end{aligned}$$

from the energy bound, so

$$\text{dist}^{F_\varepsilon(u_\varepsilon)}(U_{1,\varepsilon}, U_{i,\varepsilon}) \leq 4C |\log \varepsilon|^2 \varepsilon^\kappa.$$




FIGURE 1. Schematic illustration of the modification process. From top to bottom E_ε (all components), E'_ε (only the large components), E''_ε (also the small components on connecting curves), E'''_ε (same components, connected with tubes), E''''_ε (smoothed out version of E'''_ε).

Here we denoted as usual

$$\text{dist}^{F(u_\varepsilon)}(A, B) = \inf \left\{ d^{F(u_\varepsilon)}(x, y) \mid x \in A, y \in B \right\}.$$

This means that there exist two points $x \in U_{1,\varepsilon}$, $y \in U_{i,\varepsilon}$ and a Lipschitz curve γ_i from x to y inside Ω such that

$$\int_{\gamma_i} F_\varepsilon(u_\varepsilon) \, d\mathcal{H}^1 \leq 8C |\log \varepsilon|^2 \varepsilon^\kappa.$$

Without loss of generality, we may assume that the curve is C^∞ -smooth, and we observe that in particular

$$\mathcal{H}^1(\gamma_i \setminus \{G(u_\varepsilon) > t_\varepsilon\}) = \int_{\gamma_i \setminus \{G(u_\varepsilon) > t_\varepsilon\}} F_\varepsilon(u_\varepsilon) \, d\mathcal{H}^1 \leq 8C |\log \varepsilon|^2 \varepsilon^\kappa.$$

Making the curve potentially shorter, we may assume that it has a unique point of entry and point of exit from every connected component of $E_\varepsilon = \{G(u_\varepsilon) > t_\varepsilon\}$ since every connected open subset of \mathbb{R}^n is also path-connected. If γ_i happens to pass through a connected component of E_ε which we had eliminated before, we need to add it back:

$$E''_\varepsilon = \bigcup_{i=1}^{N_\varepsilon} \left(U_{i,\varepsilon} \cup \bigcup_{U_{j,\varepsilon} \cap \gamma_i \neq \emptyset} U_{j,\varepsilon} \right)$$

and note that still $E''_\varepsilon \rightarrow E$ and $P(E''_\varepsilon) \leq P(E_\varepsilon)$. Now we know that

$$E''_\varepsilon \cup \bigcup_{i=1}^{N_\varepsilon} (\gamma_i \setminus E''_\varepsilon)$$

is connected (but not open). Since γ_i is C^∞ -smooth and only meets finitely many connected components, there exists $\rho_\varepsilon > 0$ such that the tubular neighbourhood

$$B_{\rho_\varepsilon}(\gamma_i) = \{x \in \mathbb{R}^2 \mid \text{dist}(x, \gamma_i) < \rho_\varepsilon\}$$

is compactly contained in Ω , such that

$$\mathcal{H}^1((\partial B_{\rho_\varepsilon}(\gamma_i)) \setminus E''_\varepsilon) \leq 3 \mathcal{H}^1(\gamma_i \setminus E''_\varepsilon)$$

and such that the tubes only add a negligible amount of area. Now, using that we only had at most $O(|\log \varepsilon|)$ tubes to add which were all small compared to $\frac{1}{|\log \varepsilon|}$, we observe that

$$E'''_\varepsilon = E''_\varepsilon \cup \bigcup_{i=1}^{N_\varepsilon} B_{\rho_\varepsilon}(\gamma_i)$$

is open, connected, converges to E as $\varepsilon \rightarrow 0$ and satisfies

$$\liminf_{\varepsilon \rightarrow 0} \left(\frac{c_0}{c_0 - 2\delta} \mathcal{E}_\varepsilon(u_\varepsilon) - P(E'''_\varepsilon) \right) \geq 0.$$

We also note that E'''_ε has a smooth boundary (if we choose ρ_ε small enough) except at the finitely many points where the tubular neighbourhoods hit the connected components. We can smooth those corners out locally to a set E''''_ε with the exact same properties otherwise, but a boundary which is actually C^∞ -smooth. This proves the theorem. \square

2.4. Extensions and further observations. As for the pure Modica-Mortola functional, a compactness result holds.

Remark 2.7. If $u_\varepsilon \in W_0^{1,2}(\Omega)$ is a sequence such that $\liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon) < \infty$, then there exists a subsequence $\varepsilon \rightarrow 0$ and a set of finite perimeter $E \subset \Omega$ such that

$$u_\varepsilon \xrightarrow{L^1} \chi_E, \quad P(E) \leq \overline{P_C(E)} \leq \overline{P_{C,cc,\Omega}^r(E)} \leq \liminf_{\varepsilon \rightarrow 0} \mathcal{E}_\varepsilon(u_\varepsilon).$$

In fact, the convergence holds in $L^p(\Omega)$ for all $p < \infty$.

Let us quickly collect a few thoughts on how similar ideas may be used in related problems. If we define the energy functional \mathcal{E}'_ε formally given by the same formula on $W^{1,2}(\Omega)$ instead of $W_0^{1,2}(\Omega)$, i.e.

$$\mathcal{E}'_\varepsilon : W^{1,2}(\Omega) \rightarrow \mathbb{R}, \quad \mathcal{E}'_\varepsilon(u) = \frac{1}{c_0} \int_\Omega \frac{\varepsilon}{2} |\nabla u|^2 + \frac{W(u)}{\varepsilon} dx + \varepsilon^{-\kappa} \mathcal{C}_\varepsilon(u)$$

for some $\kappa > 0$, we get a connected relaxation of the relative perimeter.

Theorem 2.8. *Assume that $\Omega \subset \mathbb{R}^2$ is a bounded Lipschitz domain. Then*

$$\left[\Gamma(L^1) - \lim_{\varepsilon \rightarrow 0} \mathcal{E}'_\varepsilon \right] (u) = \begin{cases} \overline{P_{C,rel,\Omega}^r}(\{u = 1\}) & \text{if } u \in BV(\overline{\Omega}, \{0, 1\}) \\ +\infty & \text{else} \end{cases}$$

where

$$\overline{P_{C,rel,\Omega}^r}(E) = \inf \left\{ \liminf_{n \rightarrow \infty} P_\Omega(E_n) \mid E_n \subseteq \Omega, \quad (\partial E_n) \cap \Omega \in C^\infty, \quad E_n \text{ connected}, \quad E_n \xrightarrow{L^1} E \right\}.$$

The relative perimeter $P_\Omega(E)$ of $E \subseteq \Omega$ is defined as $P_\Omega(E) = |D\chi_E|(\Omega)$ while the full perimeter of a set $E \subset \Omega$ is $P(E) = P_{\mathbb{R}^2}(E) = |D\chi_E|(\overline{\Omega}) = |D\chi_E|(\mathbb{R}^2)$, i.e. the relative perimeter does not count the part of the boundary of E that lies inside the boundary of Ω , see e.g. [2, Definition 3.35].

The proof of the lim inf-inequality is the same as that of Theorem 2.6, using the famous result that $H = W$ (i.e. that smooth functions on \mathbb{R}^n lie dense in $W^{1,2}(\Omega)$) and the relative iso-perimetric inequality which holds in all domains where the Sobolev inequality holds (in particular, Lipschitz domains). The construction of the recovery sequence goes through as before, assuming that the boundary of Ω is not too wild.

Note that a bounded set $E \subset \mathbb{R}^2$ is simply connected if and only if both E and $\mathbb{R}^2 \setminus E$ are connected. This leads us to investigate a modified functional

$$\mathcal{E}''_\varepsilon : W_0^{1,2}(\Omega) \rightarrow \mathbb{R}, \quad \mathcal{E}''_\varepsilon(u) = \mathcal{F}_\varepsilon(u) + \varepsilon^{-\kappa} [\mathcal{C}_\varepsilon^{(1)}(u) + \mathcal{C}_\varepsilon^{(2)}(u)]$$

where

$$\begin{aligned} \mathcal{C}_\varepsilon^{(1)}(u) &= \int_\Omega \int_\Omega \beta_\varepsilon(u(x)) \beta_\varepsilon(u(y)) d^{F_\varepsilon(u)}(x, y) dx dy \\ \mathcal{C}_\varepsilon^{(2)}(u) &= \int_\Omega \int_\Omega \beta_\varepsilon(1 - u(x)) \beta_\varepsilon(1 - u(y)) d^{F_\varepsilon(1-u)}(x, y) dx dy \end{aligned}$$

i.e. $\mathcal{C}_\varepsilon^{(1)}$ as before serves to keep the phase $\{u \approx 1\}$ approximately connected whereas $\mathcal{C}_\varepsilon^{(2)}$ keeps the phase $\{1 - u \approx 1\} = \{u \approx 0\}$ connected. We have the following result.

Theorem 2.9.

$$\left[\Gamma(L^1) - \lim_{\varepsilon \rightarrow 0} \mathcal{E}''_\varepsilon \right] (u) = \begin{cases} \overline{P_{sc,cc,\Omega}^r}(\{u = 1\}) & \text{if } u \in BV(\overline{\Omega}, \{0, 1\}) \\ +\infty & \text{else} \end{cases}$$

where

$$\overline{P_{sc,cc,\Omega}^r}(E) = \inf \left\{ \liminf_{n \rightarrow \infty} P_\Omega(E_n) \mid E_n \subset \Omega, \quad E_n \in C^\infty, \quad E_n \text{ simply connected}, \quad E_n \xrightarrow{L^1} E \right\}.$$

Proof. The proof of the lim sup-inequality proceeds in the usual way, so we will only look at the necessary modifications for the lim inf-inequality. The boundary of the approximating set $E_\varepsilon = \{G(u_\varepsilon) > t_\varepsilon\}$ is a compact embedded C^∞ -submanifold of \mathbb{R}^2 , so the union of finitely many circles which do not touch each other. In particular, if $U_{i,\varepsilon}$ is a connected component of E_ε , it is only in contact with *one* connected component of $\overline{E_\varepsilon}^c$.

This means that if we add a C^∞ -curve $\gamma_{i,j,\varepsilon}$ to E_ε which connects two connected components $U_{i,\varepsilon}$ and $U_{j,\varepsilon}$ in such a way that it has one entry and one exit point to $\overline{E_\varepsilon}^c$ and no loops, then every connected component of $\overline{E_\varepsilon}^c$ will still be connected after the modification. The same is true after slightly fattening the curve. A simple proof of this fact can be constructed using path-connectedness and the regularity of the approximating sets to look at tubular neighbourhoods of the boundaries and the connecting curve.

Thus we may carefully construct connecting curves $\gamma_{i,j,\varepsilon}$ between components which have no loops and connect components in such a way that there always is only one entry and one exit point for a component, also proceeding iteratively and merging components to be the same after they have been connected before connecting the next one.

After constructing E_ε'''' in such a way, we can modify the complement $\overline{E_\varepsilon''''}^c$ in the same way to make it connected without changing the fact that E_ε'''' is connected, creating a new set \widehat{E}_ε . By construction, we again barely changed the perimeter and know that both \widehat{E}_ε and the complement of its closure are connected. Since \widehat{E}_ε is also C^∞ -smooth, it follows that also $\widehat{E}_\varepsilon^c$ is connected, which means that \widehat{E}_ε is simply connected.

As before, this concludes the proof. \square

Of course it is possible to combine the previous two extensions. We conclude this section with two notes on possible applications of our approximation results.

Remark 2.10. In order to make use of our functional for image segmentation applications, it is of course possible to add a fidelity term of the form

$$\mathcal{F}_{\text{fid}}(u) = \int_{\Omega} \frac{1}{2} \Phi(x) |u(x) - g(x)|^2 dx$$

for a given image $g: \Omega \rightarrow [0, 1]$ and local fidelity prefactor $\Phi: \Omega \rightarrow [0, 1]$. This term simply carries over to the Γ -limit proved above.

3. NUMERICAL RESULTS

As in [7], we consider a fully discrete gradient flow of the functional

$$\mathcal{E}_{\text{im}}(u) = \mathcal{F}_\varepsilon(u) + \eta \mathcal{C}_\varepsilon(u) + \delta \mathcal{F}_{\text{fid}}(u).$$

The two wells of the function W in \mathcal{F}_ε are at 0 and 1. We only consider a fixed $\varepsilon = 5 \cdot 10^{-3}$ for these numerical experiments and the functions β_ε and F_ε used to define \mathcal{C}_ε are given by

$$\beta_\varepsilon(s) = \begin{cases} 0 & s \leq 1 - \alpha \\ \frac{c_1}{2}(s - 1 + \alpha)^2 & s > 1 - \alpha \end{cases}$$

and

$$F_\varepsilon(s) = \begin{cases} \frac{1}{2}(s - 1 + \alpha)^2 & s < 1 - \alpha \\ 0 & s \leq 1 - \alpha, \end{cases}$$

respectively, with c_1 chosen such that $\int_{\alpha}^1 \beta_\varepsilon(s) ds = 1$. The value of α for all numerical examples is 0.35, the value for η is 300 for all experiments where the topological penalty is turned on. The value for δ varies somewhat from experiment to experiment.

For the finite element implementation of the discrete gradient flow, we use the algorithm described in detail in [7] and a time-step size of $\tau = 5 \cdot 10^{-8}$. The basic idea is to first separate the set $\{u > 1 - \alpha\}$ into connected components and then calculate their distances (and the respective variations, both modulo a mesh-dependent factor), by using Dijkstra's algorithm [6]. All computations are done on a unit square made up of approximately $2.3 \cdot 10^4$ P1 triangle elements. Some numerical experiments are already presented in [7], we chose to not repeat those here.

The first numerical experiment, illustrated in Figures 2 and 3, shows that, indeed, the method produces a phase field approximation of the perimeter of a set plus (twice) its Steiner-tree. The double-layer introduced in order to maintain connectedness is clearly visible. In these simulations,




FIGURE 2. Results for a numerical example using an image with close disks on the unit square. From left to right: given image g (black corresponding to the value +1, pale green to 0, both with $\Phi = 1$), stationary state u without disconnectedness penalty, stationary state with disconnectedness penalty. We use $\delta = 140$.




FIGURE 3. Results for a numerical example using an image with disks of greater distance, see also Figure 2. We use $\delta = 50$.

the initial condition was given by $u = 1$, interpolated to zero on the boundary. The additional approximate perimeter introduced through the double-layer is $1.3 \cdot 10^{-2}$ for the nearby disks, and 0.87 for the further apart disks. We note that these values are somewhat below the value for twice the length of the connecting double layers ($8.4 \cdot 10^{-2}$ and 1.14, respectively) for the figures, however, our numerical examples were performed with a fairly large value for α .

The second numerical experiment shows the applicability in image segmentation. We would like to recover an a-priori known to be connected object (in this case, a simple rectangle) which has been partly occluded (in this case, by a vertical strip). In addition, there are some smaller artifacts in the image (small disks in our example) that should be ignored. The results of this experiment are displayed in Figure 4. It is evident, that the stationary points in the experiments without topological energy term do not yield the desired recovered image: starting with $u = 1$ (as in the first experiment), the mean curvature flow becomes pinned at the obstacles. Starting with $u = g$, however, the four artifacts remain visible in the segmentation and the rectangle is divided into two pieces due to the occlusion. Adding the topological term and starting with $u = g$, however, does yield an approximation of the single rectangle – the connectedness term creates a bridge between the two pieces which then, in the course of the gradient flow, expands. The artifacts in our experiment are small enough such that the energetically better solution is to pay the fidelity penalty as opposed to creating a connecting double layer.




FIGURE 4. Results for a numerical example to recover a rectangle that has been partly occluded. The two black rectangles form the image (with the occlusion being the yellow strip between), the four small disks are image artifacts that should be ignored in the segmentation. In the black regions we have $g = \Phi = 1$, in the pale yellow region $g = \Phi = 0$. The two middle images show the final stationary states without disconnectedness penalty, top for initial condition $u = g$, bottom for $u = 1$. The right image is the result of the simulation with disconnectedness penalty and initial condition $u = g$. We use $\delta = 140$.

ACKNOWLEDGEMENTS

PWD gratefully acknowledges partial support from the German Scholars Organization/Carl-Zeiss-Stiftung in the form of the Wissenschaftler-Rückkehrprogramm. MN was supported by INDAM-GNAMP and by the University of Pisa Project PRA 2017 ?Problemi di ottimizzazione e di evoluzione in ambito variazionale?.

APPENDIX A. PROOF OF LEMMA 2.3

Recall the statement we are showing here: *If $U \subset \mathbb{R}^n$ is a convex open set and $K \subset U$ is compact, then there exists a C^∞ -diffeomorphism $\phi : \mathbb{R}^n \rightarrow U$ such that*

- (1) $\phi(x) = x$ for all $x \in K$ and
- (2) $|\phi(x) - \phi(y)| \leq |x - y|$ for all $x, y \in \mathbb{R}^n$.

Proof of Lemma 2.3. Step 1. Since K is compact, also its convex hull is compact (easily proved using sequential compactness), and since U is convex, we find that $\text{conv}(K) \subset U$. In particular, $\delta := \text{dist}(\text{conv}(K), \partial U) > 0$. The distance function

$$d : \mathbb{R}^n \rightarrow [0, \infty), \quad d(x) = \text{dist}(x, \text{conv}(K))$$

is 1-Lipschitz and convex since for $x, y \in \mathbb{R}^n$ and $\lambda \in [0, 1]$ we have

$$\begin{aligned} \lambda d(x) + (1 - \lambda) d(y) &= \lambda |x - \pi(x)| + (1 - \lambda) |y - \pi(y)| \\ &= |\lambda x - \lambda \pi(x)| + |(1 - \lambda)y - (1 - \lambda)\pi(y)| \\ &\geq |\lambda x - \lambda \pi(x) + (1 - \lambda)y - (1 - \lambda)\pi(y)| \\ &= |(\lambda x + (1 - \lambda)y) - (\lambda \pi(x) + (1 - \lambda)\pi(y))| \\ &\geq d(\lambda x + (1 - \lambda)y) \end{aligned}$$

where $\pi : \mathbb{R}^n \rightarrow \text{conv}(K)$ denotes the closest point projection onto a closed convex set $\text{conv}(K)$. Since d is convex, also its convolution d_ε with a standard mollifier of scale ε is convex as the convexity property is preserved due to the linearity of the operation. Now, when we choose ε so small that $\|d - d_\varepsilon\|_{L^\infty} < \frac{\delta}{3}$, we can use Sard's theorem and the regular value theorem together with the convexity of d_ε to find $\frac{\delta}{3} < t < \frac{2\delta}{3}$ such that $E := \{d_\varepsilon < t\}$ satisfies the following:

- (1) $\text{conv}(K) \Subset E \Subset U$,
- (2) E is convex and
- (3) $\partial E \in C^\infty$.

We can now forget K, π and δ and only work with E .

Step 2. Now denote $\delta := \text{dist}(\partial E, \partial U)$, $d_x := \text{dist}(x, \overline{E})$ and let π be the closest point projection onto \overline{E} . π is a 1-Lipschitz map which is the identity on \overline{E} and compresses the exterior space $\mathbb{R}^n \setminus \overline{E}$ into $\partial E \subset \overline{E} \Subset U$. It is C^∞ -smooth on E and $\mathbb{R}^n \setminus \overline{E}$, but only continuous at ∂E , and definitely not a diffeomorphism. We can, however, use the little space between \overline{E} and ∂U to make it smooth and a diffeomorphism.

Let $f : [0, \infty) \rightarrow [0, \frac{\delta}{2})$ be a C^∞ -function such that

$$f(t) = t \quad \forall t < \delta/4, \quad 0 < f' \leq 1.$$

Any point $x \in \mathbb{R}^n \setminus \overline{E}$ can be written as $x = \pi(x) + d(x)\nu_{\pi(x)}$ where ν is the exterior normal field to ∂E – it is well-known that the map

$$\tilde{\phi} : \partial E \times (-\varepsilon, \varepsilon) \rightarrow \{\text{dist}(\cdot, \partial E) < \varepsilon\}, \quad \tilde{\phi}(x, t) = x + t\nu_x$$

is always a diffeomorphism for small enough ε (see e.g. [9, Section 14.6]), and since E is convex, it is easy to show that the map is bijective on the whole exterior domain using the uniqueness of the closest point projection. It is a diffeomorphism since when τ is a tangent vector to ∂E we have

$$\nabla_\tau \tilde{\phi}(x, t) = \tau + tA(\tau), \quad \partial_t \tilde{\phi}(x, t) = \nu_x$$

where A denotes the shape operator of ∂E , and because E is convex, $\langle \tau, A(\tau) \rangle \geq 0$ and thus the derivative map is injective. Abbreviating $d_x = d(x)$, we define the new diffeomorphism

$$\phi : \mathbb{R}^n \rightarrow U, \quad \phi(x) = \begin{cases} x & x \in E \\ \pi(x) + f(d_x)\nu_{\pi(x)} & x \notin E. \end{cases}$$

Since $\phi(x) = x$ for all x in a neighbourhood of \overline{E} , the function ϕ is C^∞ -smooth. It remains to show that it is a non-expansive diffeomorphism.

Step 3. For diffeomorphic smoothness, we only need to show that

$$\phi : \mathbb{R}^n \setminus \overline{E} \rightarrow \left\{ \text{dist}(\cdot, E) < \frac{\delta}{2} \right\} \setminus \overline{E}$$

is a diffeomorphism since $\phi = \text{id}$ on a neighbourhood of \overline{E} . But this is obvious since ϕ is given as

$$\phi(x) = \tilde{\phi} \circ \alpha \circ \tilde{\phi}^{-1}(x) \quad \text{where } \alpha : \partial E \times (0, \infty) \rightarrow \partial E \times \left(0, \frac{\delta}{2}\right), \quad \alpha(x, t) = (x, f(t))$$

Step 4. To see that ϕ is non-expansive, we need to check this in the cases that $x, y \in \mathbb{R}^n \setminus \overline{E}$ and $x \in \overline{E}, y \in \mathbb{R}^n \setminus \overline{E}$. Let us look at the simpler second case first. Then

$$\begin{aligned} |\phi(x) - \phi(y)|^2 &= |x - [\pi(y) + f(d_y)\nu_y]|^2 \\ &= |x - \pi(y)|^2 + 2f(d_y)\langle x - \pi(y), -\nu_y \rangle + f(d_y)^2 \\ &\leq |x - \pi(y)|^2 + 2d_y\langle x - \pi(y), -\nu_y \rangle + d_y^2 \\ &= |x - y|^2 \end{aligned}$$

since $f(d_y) \leq d_y$ and since $x \in \overline{E}$, so by a common characterization of the closest point projection

$$\langle -\nu_y, x - \pi(y) \rangle = \frac{1}{|y - \pi(y)|} \langle y - \pi(y), \pi(y) - x \rangle \geq 0.$$

In the first case, we have

$$\begin{aligned} |\phi(x) - \phi(y)|^2 &= |\pi(x) + f(d_x)\nu_{\pi(x)} - [\pi(y) + f(d_y)\nu_{\pi(y)}]|^2 \\ &= |\pi(x) - \pi(y)|^2 + 2\langle \pi(x) - \pi(y), f(d_x)\nu_{\pi(x)} - f(d_y)\nu_{\pi(y)} \rangle \\ &\quad + |f(d_x)\nu_{\pi(x)} - f(d_y)\nu_{\pi(y)}|^2 \\ &= |\pi(x) - \pi(y)|^2 + 2f(d_x)\langle \nu_{\pi(x)}, \pi(x) - \pi(y) \rangle + 2f(d_y)\langle \nu_{\pi(y)}, \pi(y) - \pi(x) \rangle \\ &\quad + |f(d_x)\nu_{\pi(x)} - f(d_y)\nu_{\pi(y)}|^2 \\ &\leq |\pi(x) - \pi(y)|^2 + 2d_x\langle \nu_{\pi(x)}, \pi(x) - \pi(y) \rangle + 2d_y\langle \nu_{\pi(y)}, \pi(y) - \pi(x) \rangle \\ &\quad + |f(d_x)\nu_{\pi(x)} - f(d_y)\nu_{\pi(y)}|^2 \end{aligned}$$

as before. To treat the last term, consider $0 < s \leq t$ and $\nu, e \in S^{n-1}$ and observe that

$$\begin{aligned} |f(t)\nu - f(s)e|^2 &= |[f(t) - f(s)]\nu + f(s)[\nu - e]|^2 \\ &= [f(t) - f(s)]^2 + 2[f(t) - f(s)]f(s)\langle \nu, \nu - e \rangle + f(s)^2|\nu - e|^2 \\ &\leq |t - s|^2 + 2[t - s]s\langle \nu, \nu - e \rangle + s^2|\nu - e|^2 \\ &= |t\nu - se|^2. \end{aligned}$$

Applying this in the above inequality, we find that in total

$$\begin{aligned} |\phi(x) - \phi(y)|^2 &\leq |\pi(x) - \pi(y)|^2 + 2d_x\langle \nu_{\pi(x)}, \pi(x) - \pi(y) \rangle + 2d_y\langle \nu_{\pi(y)}, \pi(y) - \pi(x) \rangle \\ &\quad + |d_x\nu_{\pi(x)} - d_y\nu_{\pi(y)}|^2 \\ &= |x - y|^2. \end{aligned}$$

This concludes the proof. \square

REFERENCES

- [1] L. Ambrosio, V. Caselles, S. Masnou, and J.-M. Morel. Connected components of sets of finite perimeter and applications to image processing. *Journal of the European Mathematical Society*, 3(1):39–92, 2001.
- [2] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*, volume 254. Clarendon Press Oxford, 2000.
- [3] F. Benmansour, G. Carlier, G. Peyre, and F. Santambrogio. Derivatives with respect to metrics and applications: subgradient marching algorithm. *Numerische Mathematik*, 116(3):357–381, 2010.
- [4] M. Bonnivard, A. Lemenant, and F. Santambrogio. Approximation of length minimization problems among compact connected sets. *SIAM J. Math. Anal.*, 47(2):1489–1529, 2015.
- [5] F. Dayrens, S. Masnou, and M. Novaga. Relaxation of the perimeter under connectedness constraints in the plane. In preparation.
- [6] E. W. Dijkstra. A note on two problems in connexion with graphs. *Numerische Mathematik*, 1(1):269–271, 1959.
- [7] P. Dondl and S. Wojtowytsch. Keeping it together: a phase field version of path-connectedness and its implementation. 2018, arXiv:1806.04767.
- [8] P. W. Dondl, A. Lemenant, and S. Wojtowytsch. Phase Field Models for Thin Elastic Structures with Topological Constraint. *Arch. Ration. Mech. Anal.*, 223(2):693–736, 2017.
- [9] D. Gilbarg and N. S. Trudinger. *Elliptic Partial Differential Equations of Second Order*. Springer, 2001.
- [10] L. Modica. The gradient theory of phase transitions and the minimal interface criterion. *Arch Ration Mech Anal*, 98(2):123–142, 1987.
- [11] E. Paolini and E. Stepanov. Existence and regularity results for the Steiner problem. *Calculus of Variations and Partial Differential Equations*, 46(3-4):837–860, 2013.

PATRICK W. DONDL, ABTEILUNG FÜR ANGEWANDTE MATHEMATIK, ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, HERMANN-HERDER-STR. 10, 79104 FREIBURG I. BR., GERMANY
E-mail address: `patrick.dondl@mathematik.uni-freiburg.de`

MATTEO NOVAGA, DIPARTIMENTO DI MATEMATICA UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, 56127 PISA, ITALY
E-mail address: `matteo.novaga@unipi.it`

BENEDIKT WIRTH, INSTITUT FÜR ANALYSIS UND NUMERIK, WESTFÄLISCHE WILHELMS-UNIVERSITÄT MÜNSTER, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY
E-mail address: `Benedikt.Wirth@uni-muenster.de`

STEPHAN WOJTOWYTSCH, DEPARTMENT OF MATHEMATICAL SCIENCES, CARNEGIE-MELLON UNIVERSITY, 5000 FORBES AVENUE, PITTSBURGH, PA 15213, USA
E-mail address: `swojtowy@andrew.cmu.edu`