

STABILITY RESULTS FOR NONLOCAL GEOMETRIC EVOLUTIONS AND LIMIT CASES FOR FRACTIONAL MEAN CURVATURE FLOWS

A. CESARONI, L. DE LUCA, M. NOVAGA, AND M. PONSIGLIONE

ABSTRACT. We introduce a notion of uniform convergence for local and nonlocal curvatures. Then, we propose an abstract method to prove the convergence of the corresponding geometric flows, within the level set formulation. We apply such a general theory to characterize the limits of s -fractional mean curvature flows as $s \rightarrow 0^+$ and $s \rightarrow 1^-$. In analogy with the s -fractional mean curvature flows, we introduce the notion of s -Riesz curvature flows and characterize its limit as $s \rightarrow 0^-$. Eventually, we discuss the limit behavior as $r \rightarrow 0^+$ of the flow generated by a regularization of the r -Minkowski content.

KEYWORDS: Local and nonlocal geometric evolutions; Viscosity solutions; Level set formulation; Fractional perimeter; Fractional mean curvature flow; Riesz energy; Minkowski content.

AMS SUBJECT CLASSIFICATIONS: 53C44 35D40 35K93 35R11

CONTENTS

| | |
|---|----|
| 1. Introduction | 1 |
| 2. Level set formulation | 3 |
| 2.1. Axioms of nonlocal curvature | 4 |
| 2.2. Notion of viscosity solution | 5 |
| 2.3. Existence and uniqueness of a viscosity solution | 6 |
| 3. Convergence of nonlocal curvature flows | 10 |
| 4. Convergence of fractional mean curvature flows | 11 |
| 4.1. Fractional perimeters and curvatures | 11 |
| 4.2. Convergence of the s -fractional mean curvature flow as $s \rightarrow 0^+$ | 14 |
| 4.3. Convergence of the s -fractional mean curvature flows as $s \rightarrow 1^-$ | 17 |
| 5. The Riesz curvature flow | 19 |
| 6. The flow generated by the regularized r -Minkowski content | 22 |
| References | 25 |

1. INTRODUCTION

In recent years, several examples of geometric flows arising in different models from Physics, Biology and Materials science, have been considered and studied. Without trying to be exhaustive, some relevant examples of these evolutions are the classical mean curvature flow, together with its anisotropic versions (see [26] and the references therein), the fractional mean curvature flow [22, 24, 7, 12], nonlocal evolutions driven by suitable interaction kernels emerging as models for dislocation dynamics in crystals [4, 15].

It is well known that geometric evolutions may exhibit singularities even starting from smooth initial data, so that, together with classical solutions, much effort has been devoted to develop weak formulations taking into account topology changes and providing global solutions.

Here we focus on the so-called *level set* formulation [28, 21, 13]. Specifically, we follow the approach in [11], which provides a unified framework to deal with a wide class of local and nonlocal geometric flows. There, the authors introduced a class of generalized curvatures \mathcal{H} defined on pairs (x, E) where E is a smooth set and $x \in \partial E$, satisfying suitable structural assumptions, accounting continuity properties, monotonicity and translational invariance. For such a class of curvatures, the authors provided global existence and uniqueness of generalized geometric evolutions of sets with compact boundary, i.e., of continuous viscosity solutions of the corresponding level set equation

$$(1.1) \quad \begin{cases} \partial_t u(x, t) + |Du(x, t)| \mathcal{H}(x, \{y : u(y, t) \geq u(x, t)\}) = 0, \\ u(\cdot, 0) = u_0(\cdot), \end{cases}$$

where $u_0 : \mathbb{R}^d \rightarrow \mathbb{R}$ is a continuous function, which is constant outside a compact set. We shall revisit this theory in Section 2.

In this paper we propose an abstract method to prove stability of geometric flows with respect to curvature variations, within the same general framework introduced in [11]. We present the method in Section 3. First of all, in Definition 3.1, we introduce a notion of uniform convergence of a family $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ of curvatures to a limit curvature \mathcal{H}^∞ , where $\mathcal{H}^n, \mathcal{H}^\infty$ are generalized curvatures in the sense of [11]. By enforcing uniform bounds on the velocity of evolving balls, we recover uniform continuity estimates of the corresponding level set solutions, which provide the desired compactness properties for the viscosity solutions u^n to (1.1) with \mathcal{H} replaced by \mathcal{H}^n . Moreover, using the uniform convergence of the curvatures, and the appropriate notion of test functions for (1.1), suitable to deal with the singularities of the operators at points where $Du = 0$, we recover the uniform convergence of the differential operators appearing in (1.1); we conclude that the (whole) sequence $\{u^n\}_{n \in \mathbb{N}}$ locally uniformly converges to the unique solution u^∞ to (1.1) with $\mathcal{H} = \mathcal{H}^\infty$. This is the main abstract tool of the paper, precisely stated in Theorem 3.2.

We apply such a result to characterize the limit cases of several parametrized families of geometric flows. First, in Section 4 we analyze the asymptotics of the (reparametrized in time) s -fractional mean curvature flow, as $s \rightarrow 1$ and $s \rightarrow 0$. The limit case $s \rightarrow 1$ is actually well understood: as pointed out in [22], suitably reparametrized in time solutions of the s -fractional mean curvature flow converge to the classical mean curvature flow. The case $s \rightarrow 0$ is, to our knowledge, completely new and, in some respects, more intriguing. Indeed, the s -fractional perimeters, multiplied by s , converge (up to a prefactor) to the Lebesgue measure [27, 18], suggesting, at a first glance, that the corresponding reparametrized flows converge to a trivial geometric evolution where sets move with constant normal velocity. Such a trivial motion reflects the degeneracy of the limit behaviour of the rescaled fractional perimeters as $s \rightarrow 0$. Our methods provide a rigorous proof of these facts.

Furthermore, very recently the next order expansion of s -fractional perimeters as $s \rightarrow 0$ has been obtained in [17] in terms of Γ -convergence (see [8] for the limit $s \rightarrow 1$). The limit perimeter, referred to as *0-fractional perimeter*, is now much less degenerate, enjoying for instance the fractional isoperimetric inequality, which establishes that balls are the only minimizers of the 0-fractional perimeter under a volume constraint. In this paper we compute

the first variation of such a 0-fractional perimeter, referred to as *0-fractional curvature* (see Definition 4.3), we prove that the 0-fractional curvature is a generalized curvature in the sense of [11], so that the general results about existence and uniqueness of the level set flow obtained in [11] apply. Remarkably, we show that this is the relevant object to describe the limit of s -fractional mean curvature flows as $s \rightarrow 0$. In fact, we first show that the next order asymptotics of the s -fractional curvature as $s \rightarrow 0$ is exactly the 0-fractional curvature; then, as a byproduct of our general stability result, we prove that solutions of suitably forced s -fractional mean curvature flows converge to the solution of the 0-fractional mean curvature flow. Here the appropriate forcing term is nothing but $1/s$ times the constant normal velocity appearing in the leading term of the Taylor expansion of the s -fractional curvature.

Our method is, as a matter of fact, very robust, and applies to several different contexts. To illustrate that, in analogy with the s -fractional curvature, we introduce in Section 5 the notion of *s -Riesz curvature* \mathcal{K}^s , for $s < 0$, as the first variation of Riesz-type energies, see (5.1). We show that \mathcal{K}^s is a generalized curvature in the sense of [11], and hence the geometric flow driven by \mathcal{K}^s is globally well-defined; moreover, we study the first and second order limits of the rescaled flow as $s \rightarrow 0$, obtaining the same limit flows of the rescaled s -fractional curvature flow.

Finally, in Section 6 we consider the flow generated by a suitable regularization of the r -Minkowski content, which is the measure of the r -neighborhood of the boundary of the set, divided by $2r$, see(6.1). The r -Minkowski content has been introduced in [3] in the context of image denoising and then applied to vessel segmentation [30]. In order to get well-posedness of the corresponding geometric flow, in [11, 10] the authors introduced a regularized version of the r -Minkowski content, see (6.3) for a precise definition, whose first variation is a generalized curvature in the sense of [11]. We prove that the limit of such a geometric flow as $r \rightarrow 0$ is the classical mean curvature flow.

As already mentioned, the problem of convergence of rescaled nonlocal curvature flows to local (isotropic and anisotropic) mean curvature flows has already been faced in the literature; see for instance [22, 15, 9]. One of the merits of our abstract approach is the capability to detect the asymptotic behavior of geometric evolutions even when the limit itself has a nonlocal character; this is actually the case of the 0-fractional mean curvature flow.

It would be interesting to apply our general method to recover also the results in [15, 9], and to extend our formalism to other weak notion of solutions; one could consider for instance the method of geometric barriers by De Giorgi [16] (adopted in [9]), which turns out to be equivalent in many cases to the level set formulation, at least for local evolutions (see [5]).

Finally, our method could be exploited to study limit cases of many other geometric evolutions. As a relevant example, we mention the limit of geometric flows driven by the p -capacity as $p \rightarrow 1$. Incidentally, the viscosity approach proposed in this paper could turn out to be convenient also to deal with non-geometric equations such as nonlinear versions of s -fractional heat equations.

Acknowledgments: The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

2. LEVEL SET FORMULATION

In this section we revisit the level set formulation of generalized nonlocal curvature flows introduced in [11]. We start by providing the appropriate notion of nonlocal curvature.

2.1. Axioms of nonlocal curvature. Let \mathfrak{C} be the class of subsets of \mathbb{R}^d , which can be obtained as the closure of an open set with compact $C^{\ell,\beta}$ boundary. Throughout the paper $\ell \geq 2$ and $\beta \in [0, 1]$ will be fixed; the reader may simply assume $\ell = 2$, $\beta = 0$.

We will deal with “curvature” functions $x \mapsto \mathcal{H}(x, E) \in \mathbb{R}$ defined for $E \in \mathfrak{C}$ and $x \in \partial E$.

Definition 2.1. We say that \mathcal{H} is a nonlocal curvature if it satisfies the standing assumptions (M), (T) and (C) below.

- (M) Monotonicity: If $E, F \in \mathfrak{C}$ with $E \subseteq F$, and if $x \in \partial F \cap \partial E$, then $\mathcal{H}(x, F) \leq \mathcal{H}(x, E)$;
- (T) Translational invariance: for any $E \in \mathfrak{C}$, $x \in \partial E$, $y \in \mathbb{R}^d$, $\mathcal{H}(x, E) = \mathcal{H}(x+y, E+y)$;
- (C) Continuity: If $\{E_n\}_{n \in \mathbb{N}} \subset \mathfrak{C}$, $E \in \mathfrak{C}$, and $E_n \rightarrow E$ in \mathfrak{C} , then $\mathcal{H}(x, E_n) \rightarrow \mathcal{H}(x, E)$ for every $x \in \partial E_n \cap \partial E$.

Here and throughout the paper, by $E_n \rightarrow E$ in \mathfrak{C} we mean that there exists a sequence of diffeomorphisms $\{\Phi_n\}_{n \in \mathbb{N}}$ converging to the identity in $C^{\ell,\beta}$, with $E_n = \Phi_n(E)$.

By assumption (C), for any $\rho > 0$ we can define the quantities

$$(2.1) \quad \bar{c}(\rho) := \max_{x \in \partial B_\rho} \max\{\mathcal{H}(x, \bar{B}_\rho), -\mathcal{H}(x, \mathbb{R}^d \setminus B_\rho)\},$$

$$(2.2) \quad \underline{c}(\rho) := \min_{x \in \partial B_\rho} \min\{\mathcal{H}(x, \bar{B}_\rho), -\mathcal{H}(x, \mathbb{R}^d \setminus B_\rho)\},$$

which are continuous functions of $\rho > 0$. Observe that thanks to the monotonicity axiom (M), the functions $\rho \mapsto \bar{c}(\rho)$ and $\rho \mapsto \underline{c}(\rho)$ are nonincreasing.

In [11] (see Theorem 2.9 below) it has been proved that the assumptions (M), (T), (C) above are enough to guarantee the existence of a viscosity level set solution of the nonlocal \mathcal{H} -curvature flow; as in [11], a stronger property than (C) will be required in order to have uniqueness for such a solution. Such a property is the following:

- (C') Uniform continuity: Given $\mathcal{R} > 0$, there exists a modulus of continuity $\omega_{\mathcal{R}}$ such that the following holds. For all $E \in \mathfrak{C}$, $x \in \partial E$, such that E has both an interior and exterior ball condition of radius \mathcal{R} at x , and for all diffeomorphisms $\Phi : \mathbb{R}^d \rightarrow \mathbb{R}^d$ of class $C^{\ell,\beta}$, with $\Phi(y) = y$ for $|y - x| \geq 1$, we have

$$|\mathcal{H}(x, E) - \mathcal{H}(\Phi(x), \Phi(E))| \leq \omega_{\mathcal{R}}(\|\Phi - \text{Id}\|_{C^{\ell,\beta}}).$$

Assumption (B) below will guarantee that the curvature flow starting from a bounded set remains bounded at all times, yielding existence of global solutions.

- (B) Lower bound on the curvature of the balls: There exists $K > 0$ such that

$$(2.3) \quad \underline{c}(\rho) \geq -K\rho \quad \text{for all } \rho \geq 1.$$

Such an assumption is useful when dealing with a sequence of curvatures, since it guarantees that the corresponding geometric evolutions are all defined on the whole $[0, +\infty)$, and hence, in particular, on the same time interval. Moreover, the assumption (B) could be weakened in several ways with minor changes in the proofs, by requiring for instance $\underline{c}(\rho) \geq -K\rho + C$ for some (possibly negative) $C \in \mathbb{R}$.

The following symmetry condition guarantees that geometric evolutions are preserved passing to the complementary sets, and it will play a role in proving (uniform in) time continuity estimates.

- (S) Symmetry: For all $E \in \mathfrak{C}$ and for every $x \in \partial E$, it holds $\mathcal{H}(x, E) = -\mathcal{H}(x, \mathbb{R}^d \setminus \overset{\circ}{E})$, where $\overset{\circ}{E}$ denotes the interior of E .

2.2. Notion of viscosity solution. Let \mathcal{H} be a nonlocal curvature in the sense of Definition 2.1. Given a continuous function $u_0 : \mathbb{R}^d \rightarrow \mathbb{R}$, constant out of a compact set, we provide the proper notion of solution to the following parabolic Cauchy problem

$$(2.4) \quad \begin{cases} \partial_t u(x, t) + |Du(x, t)| \mathcal{H}(x, \{y : u(y, t) \geq u(x, t)\}) = 0 \\ u(\cdot, 0) = u_0(\cdot). \end{cases}$$

Here and in the following, D and D^2 stand for the spatial gradient and the spatial Hessian matrix, respectively. Notice that if the superlevel sets of u are not smooth, the meaning of (2.4) is unclear. For this reason, it is necessary to use a definition based on appropriate smooth test functions whose level sets curvatures are well defined. Following [11] we will adopt the appropriate framework of viscosity solutions. We start by introducing the class of admissible test functions.

Let $\gamma : (0, +\infty) \rightarrow \mathbb{R}$ be a nonincreasing continuous function with $\gamma \geq \bar{c}$, where $\bar{c}(\rho)$ is the function introduced in (2.1). Let \mathcal{F} be the family of functions $f \in C^\infty([0, +\infty))$ such that $f(0) = f'(0) = f''(0) = 0$, $f''(r) > 0$ for all r in a neighborhood of 0, f is constant in $[M, +\infty)$ for some $M > 0$ (depending on f), and

$$(2.5) \quad \lim_{\rho \rightarrow 0^+} f'(\rho) \gamma(\rho) = 0.$$

In [23, p. 229] it has been proven that the family \mathcal{F} is not empty. Note that (2.5) implies

$$(2.6) \quad \lim_{\rho \rightarrow 0^+} f'(\rho) \gamma(f^{-1}(\rho)) = 0,$$

since $f^{-1}(\rho) > \rho$ for small values of ρ and γ is nonincreasing.

With a small abuse of language, we will say that a function $g : \mathbb{R}^d \times A \rightarrow \mathbb{R}$, with $A \subseteq [0, +\infty)$, is constant outside a compact set if for all $t \in A$ and for all $t' \in A \cap [0, t)$ we have $g(\cdot, t') \equiv C_{t'}$ on $\mathbb{R}^d \setminus \mathcal{K}_t$ for some $C_{t'} \in \mathbb{R}$ and some compact set $\mathcal{K}_t \subset \mathbb{R}^d$.

Definition 2.2. Let $\hat{z} = (\hat{x}, \hat{t}) \in \mathbb{R}^d \times (0, +\infty)$ and let $A \subset (0, +\infty)$ be any open bounded interval containing \hat{t} . We will say that $\varphi \in C^0(\mathbb{R}^d \times \bar{A})$ is *admissible at the point* $\hat{z} = (\hat{x}, \hat{t})$ if it is of class C^2 in a neighborhood of \hat{z} , if it is constant out of a compact set, and, in case $D\varphi(\hat{z}) = 0$, the following holds: there exists $f \in \mathcal{F}$ and $\omega \in C^\infty([0, +\infty))$ with $\omega(0) = \omega'(0) = 0$, $\omega(r) > 0$ for $r \neq 0$ such that

$$|\varphi(x, t) - \varphi(\hat{z}) - \varphi_t(\hat{z})(t - \hat{t})| \leq f(|x - \hat{x}|) + \omega(|t - \hat{t}|)$$

for all (x, t) in $\mathbb{R}^d \times A$.

In the following we will say that a level of a smooth function φ is noncritical if $D\varphi$ does not vanish on such a level. We are now ready to provide the definition of viscosity sub and supersolution as in [11].

Definition 2.3. An upper semicontinuous function $u : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ (in short $u \in USC(\mathbb{R}^d \times [0, +\infty))$), constant outside a compact set, is a viscosity subsolution of the Cauchy problem (2.4) if $u(\cdot, 0) \leq u_0(\cdot)$ and for all $z := (x, t) \in \mathbb{R}^d \times (0, +\infty)$ and all C^∞ -test functions φ such that φ is admissible at z and $u - \varphi$ has a maximum at z (in the domain of definition of φ) the following holds:

- (i) If $D\varphi(z) = 0$, then $\varphi_t(z) \leq 0$;

(ii) If the level set $\{\varphi(\cdot, t) = \varphi(z)\}$ is noncritical, then

$$\varphi_t(z) + |D\varphi(z)| \mathcal{H}(x, \{y : \varphi(y, t) \geq \varphi(z)\}) \leq 0.$$

A lower semicontinuous function u (in short $u \in LSC(\mathbb{R}^d \times [0, +\infty))$), constant outside a compact set, is a viscosity supersolution of the Cauchy problem (2.4) if $u(\cdot, 0) \geq u_0(\cdot)$ and for all $z = (x, t) \in \mathbb{R}^d \times (0, +\infty)$ and all C^∞ -test functions φ such that φ is admissible at z and $u - \varphi$ has a minimum at z (in the domain of definition of φ) the following holds:

- (i) If $D\varphi(z) = 0$, then $\varphi_t(z) \geq 0$;
- (ii) If the level set $\{\varphi(\cdot, t) = \varphi(z)\}$ is noncritical, then

$$\varphi_t(z) + |D\varphi(z)| \mathcal{H}(x, \{y : \varphi(y, t) \geq \varphi(z)\}) \geq 0.$$

Finally, a function u is a viscosity solution of the Cauchy problem (2.4) if its upper semicontinuous envelope is a subsolution and its lower semicontinuous envelope is a supersolution of (2.4).

Remark 2.4. As it is standard in the theory of viscosity solutions, the maximum in Definition 2.3 of subsolutions can be assumed to be strict (and similarly for supersolutions). Indeed, assume for instance that u is a subsolution, $u - \varphi$ has a maximum at some (\hat{x}, \hat{t}) , with φ as in Definition 2.3. We now replace φ by

$$\varphi^s(x, t) := \varphi(x, t) + sf(|x - \hat{x}|) + |t - \hat{t}|^2,$$

where $s > 0$ is sufficiently small and $f \in \mathcal{F}$. Then the maximum of $u - \varphi^s$ at (\hat{x}, \hat{t}) is strict and we recover the subsolution inequality for φ by letting $s \rightarrow 0$ and using the continuity of \mathcal{H} .

Remark 2.5. Property (S) in Subsection 2.1 implies that if u is a continuous viscosity solution to (2.4), then also $-u$ is a solution. Indeed, assume that u is a subsolution; if $u - \varphi$ attains a maximum at (x, t) , then setting $\psi := -\varphi$, we have that $-u - \psi$ attains a minimum at (x, t) . Moreover, $\{\psi(\cdot, t) \geq \psi(x, t)\}$ is the complementary set of the interior of $\{\varphi(\cdot, t) \geq \varphi(x, t)\}$. This fact together with (S) implies that $-u$ is a supersolution. Similarly, it follows that, if u is a supersolution, then $-u$ is a subsolution.

Throughout the paper, we will use (with a small abuse of terminology) the terms subsolutions and supersolutions (omitting the locution “of the Cauchy problem (2.4)”) also for functions which do not satisfy the corresponding inequalities at time zero.

2.3. Existence and uniqueness of a viscosity solution. The next lemma, proved in [11], establishes a comparison result between viscosity subsolutions and smooth supersolutions, and vice-versa.

Lemma 2.6. *Let \mathcal{H} be a nonlocal curvature. Let $u \in USC(\mathbb{R}^d \times [0, +\infty))$ be a subsolution of (2.4). Let $0 \leq t_0 \leq t_1 < +\infty$, and let $\varphi \in C^2(\mathbb{R}^d \times [t_0, t_1])$ be admissible at all points in the sense of Definition 2.2 and such that $\varphi(\cdot, t_0) \geq u(\cdot, t_0)$,*

$$(2.7) \quad \varphi_t(x, t) + |D\varphi(x, t)| \mathcal{H}(x, \{\varphi(\cdot, t) \geq \varphi(x, t)\}) \geq 0$$

for all $(x, t) \in \mathbb{R}^d \times (t_0, t_1)$, with $|D\varphi(x, t)| \neq 0$, and $\varphi_t(x, t) \geq 0$ if $|D\varphi(x, t)| = 0$. Then, $\varphi \geq u$ in $\mathbb{R}^d \times [t_0, t_1]$. An analogous comparison principle holds between viscosity supersolutions and classical subsolutions.

Let $\bar{\gamma} : (0, +\infty) \rightarrow \mathbb{R}$ be a nonincreasing C^1 function with $\bar{\gamma} \geq \bar{c}$ (recall (2.1)). Let $R_0 > 0$, $T > 0$. Let $\bar{R} : [0, T) \rightarrow (0, +\infty)$ be a solution to

$$(2.8) \quad \begin{cases} \dot{R}(t) = -\bar{\gamma}(R(t)) \\ R(0) = R_0. \end{cases}$$

Notice that for $T > 0$ small enough such a solution always exists and is uniquely determined. In fact, either the unique solution exists and is strictly positive for all times or there could exist a maximal interval $[0, T^*)$ such that either $\lim_{t \rightarrow T^*} \bar{R}(t) = 0$ or $\lim_{t \rightarrow T^*} \bar{R}(t) = +\infty$.

The following result establishes that balls evolving with normal velocity given by $\bar{\gamma}(\rho)$ and $K\rho$ (with K given by property (B)) provide inner and outer barriers, respectively, for any viscosity solution. Its rigorous statement and its standard proof are slight variants of [11, Lemma 2.19], the difference being that here we deal with curvatures which are not bounded from below by a constant.

Lemma 2.7. *Let \mathcal{H} be a nonlocal curvature satisfying (B). Let $\bar{\gamma} : (0, +\infty) \rightarrow \mathbb{R}$ be a nonincreasing C^1 function with $\bar{c} \leq \bar{\gamma}$ (with \bar{c} defined in (2.1)). Let moreover $R_0 > 0$ and $t_0 \geq 0$. Then we have:*

(i) *If u is a subsolution to (2.4) with $u(\cdot, t_0) \leq \lambda + \mu\chi_{B_{R_0}}(\cdot)$ for some $\lambda \in \mathbb{R}$, $\mu > 0$, then*

$$u(\cdot, t) \leq \lambda + \mu\chi_{B_{\underline{R}(t-t_0)}}(\cdot) \quad \text{for every } t \in [t_0, +\infty),$$

where $\underline{R}(t) := R_0 e^{Kt}$ for every $t \geq 0$, with K given in (2.3).

(ii) *If u is a supersolution to (2.4) with $u(\cdot, t_0) \geq \lambda + \mu\chi_{B_{R_0}}(\cdot)$ for some $\lambda \in \mathbb{R}$, $\mu < 0$, then*

$$u(\cdot, t) \geq \lambda + \mu\chi_{B_{\underline{R}(t-t_0)}}(\cdot) \quad \text{for every } t \in [t_0, +\infty).$$

(iii) *If u is a supersolution to (2.4) with $u(\cdot, t_0) \geq \lambda + \mu\chi_{B_{R_0}}(\cdot)$ for some $\lambda \in \mathbb{R}$, $\mu > 0$, then*

$$u(\cdot, t) \geq \lambda + \mu\chi_{B_{\bar{R}(t-t_0)}}(\cdot) \quad \text{for every } t \in [t_0, t_0 + T),$$

where \bar{R} is a solution to (2.8) in $[0, T)$.

(iv) *If u is a subsolution to (2.4) with $u(\cdot, t_0) \leq \lambda + \mu\chi_{B_{R_0}}(\cdot)$ for some $\lambda \in \mathbb{R}$, $\mu < 0$, then*

$$u(\cdot, t) \leq \lambda + \mu\chi_{B_{\bar{R}(t-t_0)}}(\cdot) \quad \text{for every } t \in [t_0, t_0 + T).$$

Proof. We only prove (i) and (iii), the proof of (ii) and (iv) being fully analogous.

We start with the proof of (i); for every $\varepsilon > 0$ let $\psi^\varepsilon : \mathbb{R} \rightarrow [0, +\infty)$ be a smooth nonincreasing function, constant in $(-\infty, 0]$, with support in $(-\infty, R_0 + \varepsilon]$, $(\psi^\varepsilon)'(0) = (\psi^\varepsilon)''(0) = 0$, $\psi^\varepsilon \geq \chi_{(-\infty, R_0]}$ and $\psi^\varepsilon \rightarrow \chi_{(-\infty, R_0]}$ pointwise. Moreover, introduce the function $\underline{R}^\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$ defined by $\underline{R}^\varepsilon(\tau) = (R_0 + \varepsilon)e^{K\tau}$. For every $\varepsilon > 0$ we set

$$\varphi^\varepsilon(x, t) := \lambda + \mu\psi^\varepsilon(|x| + R_0 + \varepsilon - \underline{R}^\varepsilon(t - t_0)), \quad x \in \mathbb{R}^d, t \geq t_0.$$

One can always choose ψ^ε flat enough wherever $(\psi^\varepsilon)' = 0$, so that the functions φ^ε are admissible at all points. By assumption

$$\varphi^\varepsilon(x, t_0) = \lambda + \mu\psi^\varepsilon(|x|) \geq \lambda + \mu\chi_{B_{R_0}}(x) \geq u_0(x).$$

By a direct computation we will briefly show that φ^ε satisfies the assumptions of Lemma 2.6 above, i.e., it is a smooth supersolution to (2.4). On the one hand, recalling that φ^ε is constant whenever the argument of ψ^ε is nonpositive, we easily get that if $D\varphi^\varepsilon(x, t) = 0$, then

$\varphi_t^\varepsilon(x, t) = 0$. On the other hand, in the noncritical case all the superlevels of φ^ε expand with normal velocity equal to $K\underline{R}^\varepsilon(t - t_0)$, so that

$$\begin{aligned} \varphi_t^\varepsilon(x, t) &= |\mathrm{D}\varphi^\varepsilon(x, t)|K\underline{R}^\varepsilon(t - t_0) \geq -|\mathrm{D}\varphi^\varepsilon(x, t)|\mathcal{H}(x, \overline{B}_{\underline{R}^\varepsilon(t-t_0)}) \\ &\geq -|\mathrm{D}\varphi^\varepsilon(x, t)|\mathcal{H}(x, \{\varphi^\varepsilon(\cdot, t) \geq \varphi^\varepsilon(x, t)\}), \end{aligned}$$

where the first inequality is a consequence of (B) and the second one follows noticing that, by the very definition of φ^ε ,

$$\{\varphi^\varepsilon(\cdot, t) \geq \varphi^\varepsilon(x, t)\} \subset B_{\underline{R}^\varepsilon(t-t_0)}$$

and using property (M). By Lemma 2.6, we deduce that $u \leq \varphi^\varepsilon$, which, sending $\varepsilon \rightarrow 0$, yields $u \leq \lambda + \mu\chi_{B_{\underline{R}(t-t_0)}}$.

Let us pass to the proof of (iii). For every $\varepsilon > 0$ let now $\psi^\varepsilon : [0, +\infty) \rightarrow [0, +\infty)$ be a smooth nonincreasing function, with support in $(-\infty, R_0 - \varepsilon]$, $(\psi^\varepsilon)'(0) = (\psi^\varepsilon)''(0) = 0$, $\psi^\varepsilon \leq \chi_{[0, R_0]}$ and $\psi^\varepsilon \rightarrow \chi_{[0, R_0]}$ pointwise. Moreover, consider the Cauchy problem

$$(2.9) \quad \begin{cases} \dot{R}(t) = -\bar{\gamma}(R(t)) \\ R(0) = R_0 - \varepsilon. \end{cases}$$

Since $\bar{\gamma} \in C^1$, for $\varepsilon > 0$ small enough the (unique) solutions \overline{R}^ε to the problem (2.9) are all defined and C^2 regular on some interval $[0, T)$, with $T > 0$ independent of ε , and uniformly converge to the solution \overline{R} to (2.8).

Now, setting, for every $\varepsilon > 0$,

$$\varphi^\varepsilon(x, t) := \lambda + \mu\psi^\varepsilon(|x| + R_0 - \varepsilon - \overline{R}^\varepsilon(t - t_0)), \quad x \in \mathbb{R}^d, t \geq t_0,$$

and arguing as in the proof of (i), one can easily show the claim. \square

While Lemma 2.6 provides a comparison principle between viscosity solutions and regular evolving sets, the following result establishes the comparison principle between pairs of viscosity solutions.

Theorem 2.8 (Comparison Principle). *Assume that \mathcal{H} is a nonlocal curvature satisfying (C') and (B). Let $u \in USC(\mathbb{R}^d \times [0, +\infty))$ and $v \in LSC(\mathbb{R}^d \times [0, +\infty))$, both constant (spatially) out of a compact set, be a subsolution and a supersolution of (2.4), respectively. If $u(\cdot, 0) \leq v(\cdot, 0)$, then $u \leq v$ in $\mathbb{R}^d \times [0, +\infty)$.*

The existence and uniqueness result for viscosity solutions to (2.4) is provided by the following theorem.

Theorem 2.9 (Existence and uniqueness). *Assume that \mathcal{H} is a nonlocal curvature satisfying (B). Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function with $u_0 = C_0$ in $\mathbb{R}^d \setminus B_{R_0}$, for some $C_0, R_0 \in \mathbb{R}$ with $R_0 > 0$.*

There exists a viscosity solution $u : \mathbb{R}^d \times [0, +\infty)$ to (2.4). Moreover, any viscosity solution satisfies $u(\cdot, t) = C_0$ in $\mathbb{R}^d \setminus B_{R(t)}$, where $R(t) := R_0 e^{Kt}$ with K introduced in (2.3).

Finally, if \mathcal{H} satisfies (C'), then the viscosity solution to (2.4) is unique.

Theorems 2.8 and 2.9 have been established in [11] under minor irrelevant differences in the assumptions and in the claims. In fact, in [11] the class of test functions has been defined for $\gamma = \bar{c}$. However, replacing \bar{c} with a larger γ does not affect any step in the proof. Moreover, the proof of the existence in Theorem 2.9 follows along the lines of the results [11, Subsection 2.6]; there, the existence of a viscosity solution in any given compact interval $[0, T]$ is provided,

under the assumption (B') that the curvature of balls B_R is bounded from below by a constant $-K$ independent of R . However, the results in [11] can be easily extended to obtain global solutions also replacing (B') with our weaker assumption (B) and using Lemma 2.7 in place of [11, Lemma 2.19].

Remark 2.10. The notion of viscosity solutions apparently depends on the choice of the function γ in Definition 2.2. In fact, such a dependence is fictitious, since if $\gamma_2 \geq \gamma_1$, then the class of admissible test functions corresponding to γ_2 are also admissible replacing γ_2 with γ_1 . Therefore, the unique solution provided by Theorem 2.9 does not depend on the specific choice of γ .

Given a continuous function $v : \mathbb{R}^d \times A \rightarrow \mathbb{R}$, with $A \subseteq [0, +\infty)$, we will say that a function $\omega^{\text{sp}} : [0, +\infty) \rightarrow [0, +\infty)$ is a spatial modulus of continuity of v if it is strictly increasing and continuous in $[0, R)$ for some $R > 0$, it satisfies $\omega^{\text{sp}}(0) = 0$ and

$$|v(x_1, t) - v(x_2, t)| \leq \omega^{\text{sp}}(|x_1 - x_2|) \quad \text{for all } x_1, x_2 \in \mathbb{R}^d, t \in A.$$

The time moduli of continuity are defined analogously and denoted by ω^{ti} .

Proposition 2.11 (Uniform continuity). *Let ω^{sp} be a spatial modulus of continuity and let $\gamma : (0, +\infty) \rightarrow \mathbb{R}$ be a nonincreasing C^1 function.*

Then, there exists a time modulus of continuity ω^{ti} such that, for all nonlocal curvatures \mathcal{H} satisfying (B), (C'), (S), and such that $\bar{c} \leq \gamma$ (with \bar{c} defined in (2.1)) and for any initial datum $u_0 \in C(\mathbb{R}^d)$ constant out of a compact set and with spatial modulus of continuity ω^{sp} , we have that ω^{sp} and ω^{ti} are spatial and time moduli of continuity, respectively, of the viscosity solution to (2.4) with initial datum u_0 .

Proof. The fact that ω^{sp} is a spatial modulus of continuity for u is a well known consequence of the comparison principle and the invariance by translations of the curvature \mathcal{H} . For the reader's convenience we provide such a proof. Let $\eta \in \mathbb{R}^d$ and set $v_0^\eta(x) := u_0(x + \eta) + \omega^{\text{sp}}(|\eta|)$ for every $x \in \mathbb{R}^d$. Clearly, v_0 is constant outside a compact set, and, since ω^{sp} is a modulus of continuity of u_0 , we have

$$u_0(x) \leq u_0(x + \eta) + \omega^{\text{sp}}(|\eta|) = v_0^\eta(x) \quad \text{for every } x \in \mathbb{R}^d.$$

By Theorem 2.9 and by property (T), the function $v^\eta : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ defined by $v^\eta(x, t) := u(x + \eta, t) + \omega^{\text{sp}}(|\eta|)$ is the viscosity solution of (2.4) with initial datum v_0^η . By Theorem 2.8 we have

$$u(x, t) \leq u(x + \eta, t) + \omega^{\text{sp}}(|\eta|) \quad \text{for every } x, \eta \in \mathbb{R}^d, t \in [0, +\infty).$$

We now prove the existence of the modulus of continuity ω^{ti} for u , i.e., we show that for every $\varepsilon > 0$ there exists $\tau_\varepsilon > 0$ such that

$$(2.10) \quad u(x, t_0) - \varepsilon \leq u(x, t) \leq u(x, t_0) + \varepsilon \quad \text{for every } x \in \mathbb{R}^d, 0 \leq t_0 \leq t \leq t_0 + \tau_\varepsilon.$$

We start by proving the first inequality in (2.10). Let $\varepsilon > 0$ and set $R_0^\varepsilon := (\omega^{\text{sp}})^{-1}(\varepsilon)$. Then, for every $x \in \mathbb{R}^d$

$$B_{R_0^\varepsilon}(x) \subseteq \{y : u(y, t_0) > u(x, t_0) - \varepsilon\}.$$

Therefore, $(u(x, t_0) - \varepsilon)\chi_{B_{R_0^\varepsilon}(x)}(\cdot) \leq u(\cdot, t_0)$. Let $\bar{R}(t)$ be the solution to (2.8) with $\bar{\gamma} = \gamma$ and $R_0 = R_0^\varepsilon$; notice that $\bar{R}(t) \geq \frac{R_0^\varepsilon}{2}$ for every $t_0 \leq t \leq t_0 + \tau_\varepsilon$ where $\tau_\varepsilon := \frac{R_0^\varepsilon}{2\gamma(R_0^\varepsilon/2)}$ if $\gamma(R_0^\varepsilon/2) > 0$ and $\tau_\varepsilon = 1$ otherwise.

By Lemma 2.7, we deduce that

$$u(x, t) \geq (u(x, t_0) - \varepsilon)\chi_{B_{\frac{R_0\varepsilon}{2}}(x)}(x) = u(x, t_0) - \varepsilon \quad \text{for every } t_0 \leq t \leq t_0 + \tau_\varepsilon.$$

The second inequality in (2.10) is a direct consequence of the first one, applied to the function $-u$, which, in view of Remark 2.5, is a viscosity solution to (2.4) with initial datum $-u_0$. \square

3. CONVERGENCE OF NONLOCAL CURVATURE FLOWS

In this section we consider a sequence $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ of nonlocal curvatures and introduce a notion of convergence of \mathcal{H}^n to some limit curvature \mathcal{H}^∞ . In order to have uniform continuity estimates of the corresponding level set solutions, we enforce uniform bound on the velocity of blowing up and blowing down evolving balls; more precisely, let \underline{c}^n and \bar{c}^n be defined as in (2.1) and (2.2), with \mathcal{H} replaced by \mathcal{H}^n , we set

$$(3.1) \quad \underline{c}^{\text{inf}}(\rho) := \inf_{n \in \mathbb{N}} \underline{c}^n(\rho), \quad \bar{c}^{\text{sup}}(\rho) := \sup_{n \in \mathbb{N}} \bar{c}^n(\rho).$$

We assume that

(UB) There exists $K \geq 0$ such that $\underline{c}^{\text{inf}}(\rho) \geq -K\rho$ for all $\rho > 1$, and $\bar{c}^{\text{sup}}(\rho) < +\infty$ for all $\rho > 0$.

Definition 3.1. Let $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ be a sequence of nonlocal curvatures and let \mathcal{H}^∞ be a nonlocal curvature. We say that $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ converges to \mathcal{H}^∞ and we write $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ whenever for every $\{E_n\}_{n \in \mathbb{N}} \subset \mathfrak{C}$, $E \in \mathfrak{C}$ with $E_n \rightarrow E$ in \mathfrak{C} , and for every $x \in \partial E \cap \partial E_n$, it holds $\mathcal{H}^n(x, E_n) \rightarrow \mathcal{H}^\infty(x, E)$.

Notice that, if $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies (UB) and $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$, then \mathcal{H}^∞ satisfies (B). Moreover, if \mathcal{H}^n satisfy (S) for every $n \in \mathbb{N}$ and $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$, then \mathcal{H}^∞ satisfies (S).

Theorem 3.2. *Let $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ be a sequence of nonlocal curvatures satisfying (C'), (S) and (UB) such that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ for some nonlocal curvature \mathcal{H}^∞ satisfying (C').*

Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function with $u_0 = C_0$ in $\mathbb{R}^d \setminus B_{R_0}$, for some $C_0, R_0 \in \mathbb{R}$ with $R_0 > 0$. For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^n . Then $u^n \rightarrow u^\infty$ where $u^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^∞ .

Proof. Notice that, under the assumption (UB), the function \bar{c}^{sup} is finite and nonincreasing; therefore, there exists a nonincreasing C^1 function $\gamma : (0, +\infty) \rightarrow \mathbb{R}$ with $\gamma \geq \bar{c}^{\text{sup}}$.

Let $\bar{R}(t)$ be a solution to (2.8) with $\bar{\gamma} = \gamma$, and set $\underline{R}(t) := R_0 e^{Kt}$ for every $t \geq 0$, with K given by (UB).

By Theorem 2.9, given $T > 0$, the functions $u^n(x, t) \equiv C_0$ for all $0 \leq t \leq T$ and for all $x \in \mathbb{R}^d \setminus B_{\underline{R}(T)}$. By Proposition 2.11 and by Ascoli-Arzelá Theorem, we have that, up to a (not- relabeled) subsequence, $u^n \rightarrow u^\infty$ locally uniformly on $\mathbb{R}^d \times [0, +\infty)$, for some function u^∞ with $u^\infty(\cdot, t) \equiv C_0$ in $\mathbb{R}^d \setminus B_{\underline{R}(t)}$ for every $t \geq 0$, and such that u^∞ has the same spatial modulus of continuity ω^{sp} of u_0 and time modulus of continuity ω^{ti} given by Proposition 2.11.

It remains to show that u^∞ is a viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^∞ . We only prove that u is a subsolution, since the proof that u is a supersolution is identical, and we will verify Definition 2.3 employing test functions φ which are admissible, according to Definition 2.2, with the choice of γ done at the beginning of this proof.

Let $\bar{z} = (\bar{x}, \bar{t}) \in \mathbb{R}^d \times (0, +\infty)$ and let φ be a test function which is admissible at \bar{z} , such that $u^\infty - \varphi$ has a maximum at \bar{z} .

First we consider the case that the level $\{\varphi(\cdot, \bar{t}) = \varphi(\bar{z})\}$ is noncritical. In view of Remark 2.4, we can assume that the maximum at \bar{z} is strict. Since $u^n \rightarrow u^\infty$ locally uniformly, for n large enough, $u^n - \varphi$ admits a maximum at a point $\bar{z}^n = (\bar{x}^n, \bar{t}^n)$ for some $\bar{z}^n \rightarrow \bar{z}$. Moreover, for n large enough, the levels $\{\varphi(\cdot, \bar{t}^n) = \varphi(\bar{z}^n)\}$ are noncritical. Since u^n are subsolutions, we have

$$\partial_t \varphi(\bar{z}^n) + |\mathrm{D}\varphi(\bar{z}^n)| \mathcal{H}^n(\bar{x}_n, \{\varphi(\cdot, \bar{t}^n) \geq \varphi(\bar{z}^n)\}) \leq 0,$$

which, letting $n \rightarrow +\infty$, in view of properties (C) and (T), yields (ii) of Definition 2.3.

Assume now that $\mathrm{D}\varphi(\bar{z}) = 0$; then

$$|\varphi(x, t) - \varphi(\bar{z}) - \varphi_t(\bar{z})(t - \bar{t})| \leq f(|x - \bar{x}|) + \omega(|t - \bar{t}|),$$

with the functions f and ω as in Definition 2.2. Let us consider the admissible test function

$$\psi(x, t) := \varphi_t(\bar{z})(t - \bar{t}) + 2f(|x - \bar{x}|) + 2\omega(|t - \bar{t}|).$$

It is easy to see that $u^n - \psi$ admits a strict maximum at some $\bar{z}^n = (\bar{x}^n, \bar{t}^n)$ with $\bar{z}^n \rightarrow \bar{z}$. If $\mathrm{D}\psi(\bar{z}^n) = 0$, then

$$(3.2) \quad 0 \geq \partial_t \psi(\bar{z}^n) = \varphi_t(\bar{z}) + 2 \operatorname{sgn}(\bar{t}_n - \bar{t}) \omega'(|\bar{t}_n - \bar{t}|).$$

If $\mathrm{D}\psi(\bar{z}^n) \neq 0$, then, since ψ is radial, the level $\{\psi(\cdot, \bar{t}^n) = \psi(\bar{z}^n)\}$ is noncritical, and

$$(3.3) \quad \begin{aligned} 0 &\geq \partial_t \psi(\bar{z}^n) + |\mathrm{D}\psi(\bar{z}^n)| \mathcal{H}^n(\bar{x}^n, \{\psi(\cdot, \bar{t}^n) \geq \psi(\bar{z}^n)\}) \\ &= \varphi_t(\bar{z}) + 2 \operatorname{sgn}(\bar{t}_n - \bar{t}) \omega'(|\bar{t}_n - \bar{t}|) + 2|f'(|\bar{x}^n - \bar{x}|)| \mathcal{H}^n(\bar{x}^n, \chi_{\mathbb{R}^d \setminus B_{|\bar{x} - \bar{x}^n|}(\bar{x})}) \\ &\geq \varphi_t(\bar{z}) + 2 \operatorname{sgn}(\bar{t}_n - \bar{t}) \omega'(|\bar{t}_n - \bar{t}|) - 2|f'(|\bar{x}^n - \bar{x}|)| \gamma(|\bar{x} - \bar{x}^n|). \end{aligned}$$

By (3.2) and (3.3), letting $n \rightarrow +\infty$, recalling also (2.6) and the fact that $\omega'(0) = 0$, we have that u^∞ satisfies condition (i) of Definition 2.3. \square

4. CONVERGENCE OF FRACTIONAL MEAN CURVATURE FLOWS

Here we apply the general theory of Section 3 to study the limit cases of s -fractional mean curvature flows as $s \rightarrow 0$ and $s \rightarrow 1$.

4.1. Fractional perimeters and curvatures. Let $s \in (0, 1)$. For every $E \in \mathfrak{C}$ and for every $x \in \partial E$, the s -fractional curvature of E at x is defined [22] by

$$(4.1) \quad \mathrm{H}^s(x, E) = \lim_{r \rightarrow 0^+} \int_{\mathbb{R}^d \setminus B_r(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^{d+s}} \, dy.$$

It is well known that the curvature \mathcal{H}^s is the first variation of the s -fractional perimeter \mathcal{P}^s defined by

$$(4.2) \quad \mathcal{P}^s(E) := \int_E \int_{\mathbb{R}^d \setminus E} \frac{1}{|x - y|^{d+s}} \, dy \, dx.$$

The following definition introduces the notion of 0-fractional curvature.

Definition 4.1 (The zero curvature). Let $E \in \mathfrak{C}$. We define the 0-fractional curvature as

$$(4.3) \quad \mathbb{H}^0(x, E) := \begin{cases} \lim_{R \rightarrow +\infty} \lim_{r \rightarrow 0^+} \int_{B_R(x) \setminus B_r(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy - d\omega_d \log R, & E \subset\subset \mathbb{R}^d, \\ \lim_{R \rightarrow +\infty} \lim_{r \rightarrow 0^+} \int_{B_R(x) \setminus B_r(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy + d\omega_d \log R, & \mathbb{R}^d \setminus E \subset\subset \mathbb{R}^d. \end{cases}$$

Notice that for every compact set $E \in \mathfrak{C}$ we have $\mathbb{H}^0(x, E) = \mathbb{H}^0(x, \mathbb{R}^d \setminus \overset{\circ}{E})$ and, for all $R > \text{diam}(E)$,

$$(4.4) \quad \mathbb{H}^0(x, E) = \lim_{r \rightarrow 0^+} \int_{B_R(x) \setminus B_r(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy - d\omega_d \log R.$$

Observe also that $\mathbb{H}^0(x, E)$ coincides with $L_\Delta \chi_E(x)$, up to a dimensional constant, where L_Δ is the logarithmic Laplacian introduced in [14].

Remark 4.2. One can prove that, for E compact, $\mathbb{H}^0(x, E)$ is the first variation of the 0-fractional perimeter \mathcal{P}^0 introduced in [17], and defined for all measurable sets E with finite volume as

$$(4.5) \quad \mathcal{P}^0(E) := \int_E \left[\int_{B_R(x) \setminus E} \frac{1}{|x - y|^d} dy - \int_{E \setminus B_R(x)} \frac{1}{|x - y|^d} dy - d\omega_d \log R \right] dx.$$

It is proved in [17] that (4.5) is independent of R . We also notice that for all $s \in [0, 1)$ the s -fractional curvatures introduced above are well defined for all sets with compact boundary of class $C^{1,1}$. This is a consequence of the fact that if B^\pm are interior and exterior tangent balls to ∂E at x , then, due to the symmetry of the kernel, the integral contributions of $\mathbb{H}^s(x, E)$ on B^\pm cancel each other, while outside such balls the kernel is integrable. This well known fact is the content of the two lemmas below.

Lemma 4.3. *Let $\delta > 0$ and let $\bar{s} \in [0, 1)$. Then, there exists an increasing continuous function $\omega_\delta^{\bar{s}} : [0, +\infty) \rightarrow \mathbb{R}$ with $\omega_\delta^{\bar{s}}(0) = 0$ such that for every $s \in [0, \bar{s}]$ and for every $\eta \geq 0$*

$$(4.6) \quad \int_{B_\eta \setminus (B_\delta(\delta e_d) \cup B_\delta(-\delta e_d))} \frac{1}{|y|^{d+s}} dy \leq \omega_\delta^{\bar{s}}(\eta),$$

where e_d denotes the d -th vector of the canonical basis of \mathbb{R}^d .

Lemma 4.4. *Let $E \in \mathfrak{C}$ and let $\delta > 0$ be such that $B_\delta(-\delta e_d) \subseteq E$ and $B_\delta(\delta e_d) \subseteq \mathbb{R}^d \setminus E$. Then, for every $s \in [0, 1)$ and for every $\eta > 0$, it holds*

$$(4.7) \quad \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^{d+s}} dy = \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^{d+s}} dy.$$

Proposition 4.5. *For every $s \in [0, 1)$, the functionals \mathbb{H}^s satisfy the properties (M), (T), (C'), (B), and (S) in Subsection 2.1.*

Proof. The properties (M), (T) and (S) are easy consequences of the very definitions of the fractional curvatures. Concerning property (B), it is trivially satisfied by \mathbb{H}^s for positive s , since in this case the fractional curvature of any ball is always positive. For $s = 0$, we first notice that \mathbb{H}^0 satisfies the following scaling property: for every compact set $E \in \mathfrak{C}$ we have

$$(4.8) \quad \mathbb{H}^0(\lambda x, \lambda E) = \mathbb{H}^0(x, E) - d\omega_d \log \lambda \quad \text{for all } \lambda > 0, x \in \partial E;$$

therefore, we deduce that

$$(4.9) \quad \mathbf{H}^0(\rho x, \overline{B}_\rho) = \mathbf{H}^0(x, \overline{B}_1) - d\omega_d \log \rho \quad \text{for all } \rho > 0, x \in \partial B_1,$$

which yields property (B) also for $s = 0$.

It remains to show that property (C') holds. We prove it only for $s = 0$, being the case $s \in (0, 1)$ fully analogous and claimed in [11, Remark 5.2]. In view of property (S) it is enough to prove the claim only for compact sets $E \in \mathfrak{C}$. Let $\mathcal{R} > 0$, and let $E \in \mathfrak{C}$, $x \in \partial E$ be such that E has both an interior and exterior ball condition of radius \mathcal{R} at x . We denote by $B_{\mathcal{R}}^{\text{in}}$ and $B_{\mathcal{R}}^{\text{ou}}$ the interior and exterior tangent balls to ∂E at x , respectively. Let $\Phi : \mathbb{R}^d \rightarrow \mathbb{R}^d$ be a diffeomorphism of class $C^{\ell, \beta}$, with $\Phi(y) = y$ for $|y - x| \geq 1$. We set $\tilde{E} := \Phi(E)$ and we notice that for $\|\Phi - \text{Id}\|_{C^{\ell, \beta}}$ small enough \tilde{E} satisfies interior and exterior ball condition at $\tilde{x} := \Phi(x)$ with radius $\frac{\mathcal{R}}{2}$. We denote such tangent balls with $\tilde{B}_{\frac{\mathcal{R}}{2}}^{\text{in}}$ and $\tilde{B}_{\frac{\mathcal{R}}{2}}^{\text{ou}}$.

Let $R > \max\{\text{diam}(E), 1\}$. Notice that, if $\|\Phi - \text{Id}\|_{C^{\ell, \beta}}$ is small enough, then, also $\text{diam}(\tilde{E}) < R$. Moreover, by applying Lemma 4.3 and Lemma 4.4 with $\delta = \frac{\mathcal{R}}{2}$, for every $\eta > 0$ we have that

$$(4.10) \quad \left| \lim_{r \rightarrow 0^+} \int_{B_\eta(\tilde{x}) \setminus B_r(\tilde{x})} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| + \left| \lim_{r \rightarrow 0^+} \int_{B_\eta(x) \setminus B_r(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy \right| \leq 4\omega_{\frac{\mathcal{R}}{2}}^0(\eta).$$

Let $\varepsilon > 0$ and let η_ε be such that $4\omega_{\frac{\mathcal{R}}{2}}^0(\eta_\varepsilon) \leq \frac{\varepsilon}{3}$ and $\omega_{\frac{\mathcal{R}}{2}}^0(2\eta_\varepsilon) \leq \frac{\varepsilon}{3}$.

By (4.10), we have

$$(4.11) \quad \begin{aligned} & |\mathbf{H}^0(x, E) - \mathbf{H}^0(\Phi(x), \Phi(E))| \\ & \leq \left| \int_{B_R(x) \setminus B_{\eta_\varepsilon}(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy - \int_{B_R(\tilde{x}) \setminus B_{\eta_\varepsilon}(\tilde{x})} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| + \frac{\varepsilon}{3} \\ & \leq \left| \int_{B_R(x) \setminus B_{\eta_\varepsilon}(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy - \int_{\Phi(B_R(x) \setminus B_{\eta_\varepsilon}(x))} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| \\ & \quad + \left| \int_{\Phi(B_R(x) \setminus B_{\eta_\varepsilon}(x))} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy - \int_{B_R(\tilde{x}) \setminus B_{\eta_\varepsilon}(\tilde{x})} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| + \frac{\varepsilon}{3}, \end{aligned}$$

where in the last step we have used triangular inequality.

On the one hand, using that $\Phi = \text{Id}$ outside from $B_1(x)$, we have

$$(4.12) \quad \begin{aligned} & \left| \int_{B_R(x) \setminus B_{\eta_\varepsilon}(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy - \int_{\Phi(B_R(x) \setminus B_{\eta_\varepsilon}(x))} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| \\ & = \left| \int_{B_1(x) \setminus B_{\eta_\varepsilon}(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|^d} dy - \int_{\Phi(B_1(x) \setminus B_{\eta_\varepsilon}(x))} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| \\ & \leq \frac{\omega_d}{\eta_\varepsilon^d} \|\Phi - \text{Id}\|_{C^1}; \end{aligned}$$

on the other hand, for $\|\Phi - \text{Id}\|_{C^1}$ small enough,

$$\left(\Phi(B_R(x) \setminus B_{\eta_\varepsilon}(x))\right) \Delta \left(B_R(\tilde{x}) \setminus B_{\eta_\varepsilon}(\tilde{x})\right) \subset B_{2\eta_\varepsilon}(\tilde{x}) \setminus \left(\tilde{B}_{\frac{R}{2}}^{\text{in}} \cup \tilde{B}_{\frac{R}{2}}^{\text{ou}}\right),$$

so that, by Lemma 4.3, we have

$$(4.13) \quad \left| \int_{\Phi(B_R(x) \setminus B_{\eta_\varepsilon}(x))} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy - \int_{B_R(\tilde{x}) \setminus B_{\eta_\varepsilon}(\tilde{x})} \frac{\chi_{\mathbb{R}^d \setminus \tilde{E}}(y) - \chi_{\tilde{E}}(y)}{|\tilde{x} - y|^d} dy \right| \\ \leq \int_{B_{2\eta_\varepsilon}(\tilde{x}) \setminus (\tilde{B}_{\frac{R}{2}}^{\text{in}} \cup \tilde{B}_{\frac{R}{2}}^{\text{ou}})} \frac{1}{|\tilde{x} - y|^d} dy \\ \leq \omega_{\frac{R}{2}}^0(2\eta_\varepsilon) \leq \frac{\varepsilon}{3}.$$

In view of (4.11), (4.12) and (4.13), there exists $\tau = \tau_\varepsilon > 0$ such that

$$|\mathbf{H}^0(x, E) - \mathbf{H}^0(\Phi(x), \Phi(E))| \leq \varepsilon \quad \text{for } \|\Phi - \text{Id}\|_{C^1} \leq \tau.$$

□

Existence and uniqueness of s -fractional mean curvature flows for $s \in (0, 1)$ has been established in [22] (see also [11]). The following result, which is a direct consequence of Theorem 2.9 and of Proposition 4.5, establishes existence and uniqueness also for the 0-fractional mean curvature flow.

Theorem 4.6. *For every $s \in [0, 1)$ and for every uniformly continuous function $u_0 \in C(\mathbb{R}^d)$ constant outside a compact set, there exists a unique viscosity solution to (2.4) with \mathcal{H} replaced by \mathbf{H}^s .*

4.2. Convergence of the s -fractional mean curvature flow as $s \rightarrow 0^+$.

Theorem 4.7. *Let $\{s_n\}_{n \in \mathbb{N}} \subset (0, 1)$ be such that $s_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $\{E_n\}_{n \in \mathbb{N}} \subset \mathfrak{C}$ be such that $E_n \rightarrow E$ in \mathfrak{C} for some $E \in \mathfrak{C}$. For every $x \in \partial E \cap \partial E_n$ it holds*

$$(4.14) \quad \lim_{n \rightarrow +\infty} \mathbf{H}^{s_n}(x, E_n) - \frac{d\omega_d}{s_n} = \mathbf{H}^0(x, E),$$

where \mathbf{H}^0 is defined in (4.3). In particular,

$$(4.15) \quad \lim_{n \rightarrow +\infty} s_n \mathbf{H}^{s_n}(x, E_n) = d\omega_d.$$

Proof. We start by proving (4.15) (that for $E_n \equiv E$ has already been proved in [29]).

Let $x \in \partial E_n \cap \partial E$ for all $n \in \mathbb{N}$. First, notice that all the curvatures we are dealing with satisfy assumption (S) and are invariant by rotations (and translations). In particular, we can assume without loss of generality that E_n and E are compact, that $x = 0$ and $\nu_E(0) = \nu_{E_n}(0) = e_d$, where for all $F \in \mathfrak{C}$ and $y \in \partial F$ we denote by $\nu_F(y)$ the outer normal to ∂F at y .

Setting $\eta := 2 \text{diam}(E)$, we have $E, E_n \subset B_\eta$ for n large enough. We get

$$(4.16) \quad \mathbf{H}^{s_n}(0, E_n) = \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy + \int_{\mathbb{R}^d \setminus B_\eta} \frac{1}{|y|^{d+s_n}} dy \\ = \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy + \frac{d\omega_d}{s_n} \eta^{-s_n}.$$

Since $E_n \rightarrow E$ in \mathfrak{C} , we deduce that E_n and E satisfy a uniform interior and exterior ball condition. More precisely, there exists $\delta > 0$ such that $B_\delta(-\delta e_d) \subseteq E_n$ and $B_\delta(\delta e_d) \subseteq \mathbb{R}^d \setminus E_n$ and the same for E .

In view of (4.16) and (4.7), we deduce that

$$\begin{aligned}
 & \lim_{n \rightarrow +\infty} s_n \mathbf{H}^{s_n}(x, E_n) \\
 (4.17) \quad &= \lim_{n \rightarrow +\infty} s_n \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy + \lim_{n \rightarrow +\infty} d\omega_d \eta^{-s_n} \\
 &= d\omega_d,
 \end{aligned}$$

where the last equality follows from (4.6), noticing that $s_n \in (0, \frac{1}{2}]$ for n large enough.

Let us pass to the proof of (4.14). By arguing as in (4.16), we have

$$\begin{aligned}
 & \lim_{n \rightarrow +\infty} \mathbf{H}^{s_n}(0, E_n) - \frac{d\omega_d}{s_n} \\
 &= \lim_{n \rightarrow +\infty} \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy + d\omega_d \lim_{n \rightarrow +\infty} \frac{\eta^{-s_n} - 1}{s_n} \\
 (4.18) \quad &= \lim_{n \rightarrow +\infty} \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy - d\omega_d \log \eta.
 \end{aligned}$$

In view of (4.7) we have

$$\begin{aligned}
 & \lim_{n \rightarrow +\infty} \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy \\
 (4.19) \quad &= \lim_{n \rightarrow +\infty} \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy \\
 &= \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^d} dy,
 \end{aligned}$$

where the last equality follows by Lemma 4.3 and by the Dominated Convergence Theorem. By (4.18), (4.19) and by (4.7), we get

$$\begin{aligned}
 \lim_{n \rightarrow +\infty} \mathbf{H}^{s_n}(0, E_n) - \frac{d\omega_d}{s_n} &= \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^d} dy - d\omega_d \log \eta \\
 &= \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^d} dy - d\omega_d \log \eta,
 \end{aligned}$$

which, in view of (4.4), implies (4.14). \square

In the next two results we characterize the limit of s -fractional mean curvature flows as $s \rightarrow 0$.

Theorem 4.8. *Let $\{s_n\}_{n \in \mathbb{N}} \subset (0, 1)$ with $s_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function, constant outside a compact set.*

For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by \mathbf{H}^{s_n} , and set $v^n(x, t) := u^n(x, s_n t)$ for all $x \in \mathbb{R}^d$, $t \geq 0$. Then, $v^n \rightarrow v^\infty$ locally uniformly where $v^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by $d\omega_d$.

Proof. For every $n \in \mathbb{N}$ we set $\mathcal{H}^n := s_n \mathbf{H}^{s_n}$ and $\mathcal{H}^\infty := d\omega_d$. By Proposition 4.5, \mathcal{H}^n are nonlocal curvatures in the sense of Subsection 2.1 and satisfy (C') and (S). Trivially, also \mathcal{H}^∞ is a nonlocal curvature satisfying (C') and (S). Moreover, \mathcal{H}^n are positive on every ball of radius $\rho > 0$; furthermore, by the scaling property

$$(4.20) \quad \mathbf{H}^s(\lambda x, \lambda E) = \lambda^{-s} \mathbf{H}^s(x, E) \quad \text{for all } \lambda > 0, s \in (0, 1), E \in \mathfrak{C}, x \in \partial E,$$

we deduce that

$$\mathcal{H}^n(\rho x, \overline{B}_\rho) = \rho^{-s_n} \mathcal{H}^n(x, \overline{B}_1) \quad \text{for all } \rho > 0, n \in \mathbb{N}, x \in \partial B_1.$$

Therefore, $0 \leq \mathcal{H}^n(\rho x, \overline{B}_\rho) \leq \max(1, \rho^{-1}) \sup_n \mathcal{H}^n(x, \overline{B}_1)$ and so in view of Theorem 4.7, the sequence $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies also property (UB). Again by Theorem 4.7 (in particular, by (4.15)), we get that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ in the sense of Definition 3.1.

One can easily check that v^n are viscosity solutions to (2.4) with \mathcal{H} replaced by \mathcal{H}^n , so that, by Theorem 3.2 we can conclude that $v^n \rightarrow v^\infty$ locally uniformly, where v^∞ is the viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^∞ . \square

Theorem 4.9. *Let $\{s_n\}_{n \in \mathbb{N}} \subset (0, 1)$ with $s_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function, constant outside a compact set.*

For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by $\mathbf{H}^{s_n} - \frac{d\omega_d}{s_n}$. Then, $u^n \rightarrow u^\infty$ locally uniformly where $u^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by \mathbf{H}^0 .

Proof. For every $n \in \mathbb{N}$ we set $\mathcal{H}^n := \mathbf{H}^{s_n} - \frac{d\omega_d}{s_n}$ and $\mathcal{H}^\infty := \mathbf{H}^0$. By Proposition 4.5, \mathcal{H}^n and \mathcal{H}^∞ are nonlocal curvatures in the sense of Subsection 2.1 and satisfy (C') and (S). In view of (4.20), we have

$$(4.21) \quad \mathcal{H}^n(\rho x, \overline{B}_\rho) = \rho^{-s_n} \mathcal{H}^n(x, \overline{B}_1) + d\omega_d \frac{\rho^{-s_n} - 1}{s_n} \quad \text{for all } \rho > 0, x \in \partial B_1.$$

For $\rho > 1$, by Lagrange Theorem, for every $n \in \mathbb{N}$ there exists $\xi_n \in (0, s_n)$ such that

$$(4.22) \quad \frac{\rho^{-s_n} - 1}{s_n} = -\rho^{-\xi_n} \log \rho \geq -\log \rho$$

therefore, by (4.21), (4.22), and Theorem 4.7, we have that there exists a constant $K \geq 0$ such that

$$(4.23) \quad \underline{c}^{\text{inf}}(\rho) := \inf_{n \in \mathbb{N}} \underline{c}^n(\rho) \geq \inf_n \rho^{-s_n} \mathcal{H}^n(x, \overline{B}_1) - d\omega_d \log \rho \geq -K\rho \quad \text{for all } \rho > 1.$$

Moreover, again by (4.21), (4.22), and Theorem 4.7, it is easy to see that there exist two constants $C_1, C_2 > 0$ such that

$$(4.24) \quad \overline{c}^{\text{sup}}(\rho) := \sup_{n \in \mathbb{N}} \overline{c}^n(\rho) \leq C_1 \frac{1 + |\log \rho|}{\rho} + C_2 \quad \text{for all } \rho > 0.$$

Therefore, by (4.23) and (4.24) we deduce that $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies also property (UB), and again by Theorem 4.7 (in particular, by (4.14)), we get that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ in the sense of Definition 3.1. One can thus apply Theorem 3.2 in order to get the claim. \square

4.3. Convergence of the s -fractional mean curvature flows as $s \rightarrow 1^-$. For every $E \in \mathfrak{C}$, and for every $x \in \partial E$, we denote by $H^1(x, E)$ the scalar mean curvature of the set ∂E at x , i.e., the sum of the principal curvatures of ∂E at x . Notice that, since $E \in \mathfrak{C}$, near $x = (x', x_d)$ the boundary of E can be described, in suitable coordinates, as the graph of a function f in $C^{\ell, \beta}$, with $f(x') = 0$, $Df(x') = 0$; then, denoting by A_x the $(d-1) \times (d-1)$ Hessian matrix of f at a point $x = (x', f(x'))$, it is well known that

$$(4.25) \quad H^1(x, E) = -\frac{1}{d(d-1)\omega_d} \int_{\mathbb{S}^{d-2}} e^t A_x e \, d\mathcal{H}^{d-2}(e),$$

where \mathbb{S}^{d-2} stands for the unit sphere in \mathbb{R}^{d-1} .

Theorem 4.10. *Let $\{s_n\}_{n \in \mathbb{N}} \subset (0, 1)$ be such that $s_n \rightarrow 1$ as $n \rightarrow +\infty$. Let $\{E_n\}_{n \in \mathbb{N}} \subset \mathfrak{C}$ be such that $E_n \rightarrow E$ in \mathfrak{C} for some $E \in \mathfrak{C}$. For every $x \in \partial E \cap \partial E_n$*

$$(4.26) \quad \lim_{n \rightarrow +\infty} (1 - s_n) H^{s_n}(x, E_n) = d(d-1)\omega_d H^1(x, E).$$

Proof. The pointwise convergence (i.e., for $E_n \equiv E$) has been proved in [1, 6, 22]. By a direct inspection of the proof it is easy to check that actually the convergence is uniform. For the sake of completeness, we briefly sketch the computation. Arguing as in Theorem 4.7, we may assume that $x = 0$ and $\nu_E(0) = \nu_{E_n}(0) = e_d$, where for all $F \in \mathfrak{C}$ and $y \in \partial F$ the symbol $\nu_F(y)$ denotes the outer normal to ∂F at y . Since

$$\operatorname{div} \left(\frac{y}{|y|^{d+s}} \right) = -s \frac{1}{|y|^{d+s}},$$

we may rewrite

$$(4.27) \quad H^{s_n}(0, E_n) = \frac{2}{s_n} \int_{\partial E_n} \frac{\nu(y) \cdot y}{|y|^{d+s_n}} \, d\mathcal{H}^{d-1}(y).$$

For every $\delta > 0$ the symbol B'_δ denotes the ball in \mathbb{R}^{d-1} of center 0 and radius δ .

Since $E_n \rightarrow E$ in \mathfrak{C} there exist $\delta > 0$ and functions $f_n, f \in C^{\ell, \beta}(B'_\delta; [0, \delta])$ such that $f_n \rightarrow f$ in $C^{\ell, \beta}(B'_\delta; \mathbb{R})$ such that $f_n(0) = f(0) = 0$, $Df_n(0) = Df(0) = 0$ and

$$\begin{aligned} \partial E_n \cap B_\delta &= \{(y', f_n(y')) : y' \in B'_\delta\}, & \partial E \cap B_\delta &= \{(y', f(y')) : y' \in B'_\delta\} \\ E_n \cap B_\delta &= \{(y', y_d) : y' \in B'_\delta, y_d \leq f_n(y')\}, & E \cap B_\delta &= \{(y', y_d) : y' \in B'_\delta, y_d \leq f(y')\}. \end{aligned}$$

Let $\eta > \operatorname{diam}(\partial E)$, so that for n large enough $\partial E_n, \partial E \subset B_\eta$.

On the one hand, we notice that

$$(4.28) \quad \left| \frac{2}{s_n} \int_{\partial E_n \setminus B_\delta} \frac{\nu(y) \cdot y}{|y|^{d+s_n}} \, d\mathcal{H}^{d-1}(y) \right| \leq \frac{2}{s\delta^{d+s_n-1}} \operatorname{Per}(E_n) \leq \frac{C}{s\delta^{d+s_n-1}},$$

for some constant $C > 0$ independent of n . On the other hand we have

$$(4.29) \quad \frac{2}{s_n} \int_{\partial E_n \cap B_\delta} \frac{\nu_{E_n}(y) \cdot y}{|y|^{d+s_n}} \, d\mathcal{H}^{d-1}(y) = \frac{2}{s_n} \int_{B'_\delta} \frac{f_n(y') - Df_n(y') \cdot y'}{(f_n^2(y') + |y'|^2)^{\frac{d+s_n}{2}}} \, dy'.$$

Since $f_n(y') = \frac{1}{2}(y')^t D^2 f_n(0) y' + o(|y'|^2)$ and $Df_n(y') = D^2 f_n(0) y' + o(|y'|)$, letting $e = \frac{y'}{|y'|}$, we have that

$$(4.30) \quad f_n(y') - Df_n(y') \cdot y' = -\frac{1}{2}|y'|^2(e^t D^2 f_n(0) e) + o(|y'|^2)$$

$$(4.31) \quad \frac{1}{(f_n^2(y') + |y'|^2)^{\frac{d+s_n}{2}}} = \frac{1}{|y'|^{d+s_n} \left[1 + \frac{1}{4}|y'|^2(e^t D^2 f_n(0) e)^2 + o(|y'|^2)\right]^{\frac{d+s_n}{2}}} \\ = \frac{1}{|y'|^{d+s_n}} \left[1 - \frac{d+s_n}{8}|y'|^2(e^t D^2 f_n(0) e)^2 + o(|y'|^2)\right].$$

Replacing (4.30) and (4.31) in (4.29) we get

$$(4.32) \quad \frac{2}{s_n} \int_{\partial E_n \cap B_\delta} \frac{\nu_{E_n}(y) \cdot y}{|y|^{d+s_n}} d\mathcal{H}^{d-1}(y) \\ = -\frac{1}{s_n} \int_{B'_\delta} \frac{e^t D^2 f_n(0) e}{|y'|^{d+s_n-2}} \left[1 - \frac{d+s_n}{8}|y'|^2(e^t D^2 f_n(0) e)^2 + o(|y'|^2)\right] dy' \\ = -\frac{1}{s_n(1-s_n)} \delta^{1-s_n} \int_{\mathbb{S}^{d-2}} e^t D^2 f_n(0) e d\mathcal{H}^{d-2}(e) \\ + \frac{d+s_n}{8s_n(3-s_n)} \delta^{3-s_n} \int_{\mathbb{S}^{d-2}} (e^t D^2 f_n(0) e)^3 d\mathcal{H}^{d-2}(e) + o(1).$$

Therefore, (4.27), (4.28) and (4.32), together with the $C^{\ell, \beta}$ -convergence of f_n to f and (4.25), imply (4.26). \square

Theorem 4.11. *Let $\{s_n\}_{n \in \mathbb{N}} \subset (0, 1)$ with $s_n \rightarrow 1$ as $n \rightarrow +\infty$. Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function, constant outside a compact set.*

For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^{s_n} , and set $v^n(x, t) := u^n(x, (1-s_n)t)$ for all $x \in \mathbb{R}^d$, $t \geq 0$. Then, $v^n \rightarrow v^\infty$ locally uniformly where $v^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by $d(d-1)\omega_d \mathcal{H}^1$.

Proof. For every $n \in \mathbb{N}$ we set $\mathcal{H}^n := (1-s_n)\mathcal{H}^{s_n}$ and $\mathcal{H}^\infty := d(d-1)\omega_d \mathcal{H}^1$. By Proposition 4.5, \mathcal{H}^n are nonlocal curvatures in the sense of Subsection 2.1 and satisfy (C') and (S). Trivially, also \mathcal{H}^∞ is a nonlocal curvature satisfying (C') and (S). Moreover, \mathcal{H}^n are positive on all the balls of radius $\rho > 0$; furthermore, by the scaling property (4.20), we deduce that

$$\mathcal{H}^n(\rho x, \overline{B}_\rho) = \rho^{-s_n} \mathcal{H}^n(x, \overline{B}_1) \quad \text{for all } \rho > 0, n \in \mathbb{N}, x \in \partial B_1.$$

Therefore, in view of Theorem 4.10, the sequence $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies also property (UB) and again by Theorem 4.10, we get that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ in the sense of Definition 3.1.

One can easily check that v^n are viscosity solutions to (2.4) with \mathcal{H} replaced by \mathcal{H}^n , so that, by Theorem 3.2, we can conclude that $v^n \rightarrow v^\infty$ locally uniformly, where v^∞ is the viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^∞ . \square

Remark 4.12. There exist also anisotropic versions of the s -fractional curvature. Let $K \subseteq \mathbb{R}^d$ be a compact convex set with non-empty interior and symmetric with respect to the origin, and let $|x|_K := \inf\{\lambda : x \in \lambda K\}$, i.e., the norm in \mathbb{R}^d having the set K as unitary ball.

For every $s \in (0, 1)$, the anisotropic s -fractional perimeter of a measurable set $E \subset \mathbb{R}^d$ is defined by

$$(4.33) \quad \mathcal{P}_K^s(E) := \int_E \int_{\mathbb{R}^d \setminus E} \frac{1}{|x - y|_K^{d+s}} dy dx$$

and its first variation, i.e., the anisotropic s -fractional curvature [12] is formally given by

$$(4.34) \quad \mathbb{H}_K^s(x, E) := \lim_{r \rightarrow 0^+} \int_{\mathbb{R}^d \setminus B_r(x)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|x - y|_K^{d+s}} dy, \quad x \in \partial E.$$

It is easy to check that Proposition 4.5 applies also to these curvatures (see [12]), hence they are nonlocal curvatures in the sense of Subsection 2.1.

In [25], using suitable integration formulas, the convergence (pointwise and in the sense of Γ -convergence) of \mathcal{P}_K^s to \mathcal{P}_{ZK} as $s \rightarrow 1$ is proved. The limit \mathcal{P}_{ZK} is given by

$$(4.35) \quad \mathcal{P}_{ZK}(E) = \int_{\partial^* E} |\nu_E(x)|_{Z^*K} d\mathcal{H}^{d-1}(x)$$

where $|y|_{Z^*K} := \frac{d+1}{2} \int_K |y \cdot z| dz$. It is easy to check that $Z^*K = \{x : |x|_{Z^*K} \leq 1\}$ is strictly convex.

We expect that by similar methods it is possible to show that $(1-s)\mathbb{H}_K^s$ converge in the sense of Definition 3.1 as $s \rightarrow 1^-$ to a local anisotropic curvature \mathbb{H}_{ZK}^1 , which is given by the first variation of the anisotropic perimeter (4.35). Since Z^*K is strictly convex, then \mathbb{H}_{ZK}^1 is a curvature in the sense of Subsection 2.1, and (UB) is trivially satisfied, so Theorem 3.2 on the convergence of the corresponding geometric flows should also apply.

5. THE RIESZ CURVATURE FLOW

In this section we introduce and analyze a new nonlocal geometric flow, where the curvature is the first variation of a Riesz interaction energy.

Let $s \in (-d, 0)$. For every $E \in \mathfrak{C}$ and for every $x \in \partial E$, we set

$$(5.1) \quad \mathcal{K}^s(x, E) := \begin{cases} -2 \int_E \frac{1}{|x - y|^{d+s}} dy, & E \subset\subset \mathbb{R}^d, \\ 2 \int_{\mathbb{R}^d \setminus E} \frac{1}{|x - y|^{d+s}} dy, & \mathbb{R}^d \setminus E \subset\subset \mathbb{R}^d. \end{cases}$$

Note that, for E compact, \mathcal{K}^s is the first variation of the perimeter-like Riesz interaction functional

$$(5.2) \quad \mathcal{J}^s(E) := - \int_E \int_E \frac{1}{|x - y|^{d+s}} dy dx.$$

First of all we observe the following.

Proposition 5.1. *For every $s \in (-d, 0)$ the functionals \mathcal{K}^s satisfy the properties (M), (T), (C'), and (S) in Subsection 2.1. Moreover, if $s \in [-1, 0)$, then \mathcal{K}^s satisfies also property (B).*

Proof. The validity of properties (M), (T), (C'), and (S) can be proven by arguing exactly as in Proposition 4.5. As for (B), we observe, denoting with B'_ρ the ball in \mathbb{R}^{d-1} with radius ρ ,

that

$$\begin{aligned}
\mathcal{K}^s(0, B_\rho(\rho e_d)) &= -2 \int_{B_\rho(\rho e_d)} \frac{1}{|y|^{d+s}} dy \\
&= -2 \int_{B'_\rho} \int_{\rho - \sqrt{\rho^2 - |y'|^2}}^\rho \frac{1}{(|y'|^2 + y_d^2)^{\frac{d+s}{2}}} dy_d dy' - 2 \int_{B_\rho(\rho e_d) \cap \{y_d \geq \rho\}} \frac{1}{|y|^{d+s}} dy \\
&\geq -2 \int_{B'_\rho} \frac{\sqrt{\rho^2 - |y'|^2}}{|y'|^{d+s}} dy' - \frac{\omega_d}{\rho^s} \geq \frac{2\omega_{d-1}\rho}{s\rho^s} - \frac{\omega_d}{\rho^s}.
\end{aligned}$$

Therefore (B) is satisfied for $s \in [-1, 0)$. \square

Existence of global solution and uniqueness of s -Riesz curvature flows for $s \in [-1, 0)$ is established by the following result that is a direct consequence of Theorem 2.9 and of Proposition 5.1.

Theorem 5.2. *For every $s \in [-1, 0)$ and for every uniformly continuous function $u_0 \in C(\mathbb{R}^d)$ constant outside a compact set, there exists a unique viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{K}^s .*

Remark 5.3. Note that there holds

$$\mathcal{K}^s(0, B_\rho(\rho e_d)) \leq -2 \int_{B_\rho(\rho e_d) \cap \{y_d \geq \rho\}} \frac{1}{|y|^{d+s}} dy \leq -\frac{\omega_d}{5^{(d+s)/2} \rho^s}.$$

Therefore, for $s \in (-d, -1)$, balls blow up in finite time. Nevertheless, in this case one could prove at least local in time existence of the viscosity solution.

Theorem 5.4. *Let $\{s_n\}_{n \in \mathbb{N}} \subset (-d, 0)$ be such that $s_n \rightarrow 0^-$ as $n \rightarrow +\infty$. Let $\{E_n\}_{n \in \mathbb{N}} \subset \mathfrak{C}$ be such that $E_n \rightarrow E$ in \mathfrak{C} for some $E \in \mathfrak{C}$. For every $x \in \partial E \cap \partial E_n$ it holds*

$$(5.3) \quad \lim_{n \rightarrow +\infty} \mathcal{K}^{s_n}(x, E_n) - \frac{d\omega_d}{s_n} = H^0(x, E),$$

where H^0 is defined in (4.3). In particular,

$$(5.4) \quad \lim_{n \rightarrow +\infty} s_n \mathcal{K}^{s_n}(x, E_n) = d\omega_d.$$

Proof. The proof follows along the lines of that of Theorem 4.7; we briefly sketch it.

Let $x \in \partial E_n \cap \partial E$ for all $n \in \mathbb{N}$. First, notice that all the curvatures we are dealing with satisfy assumption (S) and are invariant by rotations (and translations). In particular, we can assume without loss of generality that E_n and E are compact, that $x = 0$ and $\nu_E(0) = \nu_{E_n}(0) = e_d$, where we recall that for all $F \in \mathfrak{C}$ and $y \in \partial F$, $\nu_F(y)$ denotes the outer normal to ∂F at y .

Setting $\eta := 2 \operatorname{diam}(E)$, we have $E, E_n \subset B_\eta$ for n large enough. Since $E_n \rightarrow E$ in \mathfrak{C} , we deduce that E_n and E satisfy a uniform interior and exterior ball condition. More precisely, there exists $\delta > 0$ such that $B_\delta(-\delta e_d) \subseteq E_n$ and $B_\delta(\delta e_d) \subseteq \mathbb{R}^d \setminus E_n$ and the same for E . Due

to this fact, and to the symmetry property of the kernel, we get

$$\begin{aligned}
(5.5) \quad \mathcal{K}^{s_n}(0, E_n) &= -2 \int_{E_n} \frac{1}{|y|^{d+s_n}} dy \\
&= - \int_{B_\delta(-\delta e_d) \cup B_\delta(\delta e_d)} \frac{1}{|y|^{d+s_n}} dy - 2 \int_{E_n \setminus B_\delta(-\delta e_d)} \frac{1}{|y|^{d+s_n}} dy \\
&= - \int_{B_\eta} \frac{1}{|y|^{d+s_n}} dy + \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{1}{|y|^{d+s_n}} dy - 2 \int_{E_n \setminus B_\delta(-\delta e_d)} \frac{1}{|y|^{d+s_n}} dy \\
&= - \int_{B_\eta} \frac{1}{|y|^{d+s_n}} dy + \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy \\
&= \frac{d\omega_d \eta^{-s_n}}{s_n} + \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy.
\end{aligned}$$

By Lemma 4.3 and using the $C^{\ell, \beta}$ convergence of E_n to E and the Dominate Convergence Theorem, we get

$$\begin{aligned}
(5.6) \quad & \lim_{n \rightarrow +\infty} \int_{B_\eta \setminus (B_\delta(-\delta e_d) \cup B_\delta(\delta e_d))} \frac{\chi_{\mathbb{R}^d \setminus E_n}(y) - \chi_{E_n}(y)}{|y|^{d+s_n}} dy \\
&= \int_{B_\eta \setminus B_\delta(-\delta e_d) \cup B_\delta(\delta e_d)} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^d} dy \\
&= \lim_{r \rightarrow 0^+} \int_{B_\eta \setminus B_r} \frac{\chi_{\mathbb{R}^d \setminus E}(y) - \chi_E(y)}{|y|^d} dy,
\end{aligned}$$

where the last equality follows from Lemma 4.4.

By (5.5) and (5.6), in view of (4.4), we can conclude that (5.3) holds true. \square

Next two results are devoted to the analysis of the s -Riesz curvature flow as $s \rightarrow 0$.

Theorem 5.5. *Let $\{s_n\}_{n \in \mathbb{N}} \subset [-1, 0)$ with $s_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function, constant outside a compact set.*

For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{K}^{s_n} , and set $v^n(x, t) := u^n(x, -s_n t)$ for all $x \in \mathbb{R}^d$, $t \geq 0$. Then, $v^n \rightarrow v^\infty$ locally uniformly where $v^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by $-d\omega_d$.

Proof. For every $n \in \mathbb{N}$ we set $\mathcal{H}^n := -s_n \mathcal{K}^{s_n}$ and $\mathcal{H}^\infty := -d\omega_d$. By Proposition 5.1, \mathcal{H}^n are nonlocal curvatures in the sense of Subsection 2.1 and satisfy (C') and (S). Trivially, also \mathcal{H}^∞ is a nonlocal curvature satisfying (C') and (S). Moreover, from the scaling property

$$(5.7) \quad \mathcal{K}^s(\lambda x, \lambda E) = \lambda^{-s} \mathcal{K}^s(x, E) \quad \text{for all } s \in (-d, 0), E \in \mathfrak{C}, x \in \partial E,$$

we deduce that

$$\mathcal{H}^n(\rho x, \overline{B}_\rho) = \rho^{-s_n} \mathcal{H}^n(x, \overline{B}_1) \quad \text{for all } \rho > 0, n \in \mathbb{N}, x \in \partial B_1.$$

Since \mathcal{H}^n is negative and $-s_n \in (0, 1]$, it follows that

$$\max\{\rho, 1\} \mathcal{H}^n(x, \overline{B}_1) \leq \mathcal{H}^n(\rho x, \overline{B}_\rho) \leq \min\{\rho, 1\} \mathcal{H}^n(x, \overline{B}_1) \quad \text{for all } \rho > 0, n \in \mathbb{N}, x \in \partial B_1,$$

which, in view of Theorem 5.4, implies that the sequence $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies also property (UB). Again by Theorem 5.4 (in particular, by (5.4)), we get that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ in the sense of Definition 3.1.

One can easily check that v^n are viscosity solutions to (2.4) with \mathcal{H} replaced by \mathcal{H}^n , so that, by Theorem 3.2 we can conclude that $v^n \rightarrow v^\infty$ locally uniformly, where v^∞ is the viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^∞ . \square

Theorem 5.6. *Let $\{s_n\}_{n \in \mathbb{N}} \subset [-1, 0)$ with $s_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function, constant outside a compact set.*

For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by $\mathcal{K}^{s_n} - \frac{d\omega_d}{s_n}$. Then, $u^n \rightarrow u^\infty$ locally uniformly where $u^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by \mathcal{H}^0 .

Proof. For every $n \in \mathbb{N}$ we set $\mathcal{H}^n := \mathcal{K}^{s_n} - \frac{d\omega_d}{s_n}$ and $\mathcal{H}^\infty := \mathcal{H}^0$. By Proposition 5.1, \mathcal{H}^n and \mathcal{H}^∞ are nonlocal curvatures in the sense of Subsection 2.1 and satisfy (C') and (S). In view of (5.7), we have

$$(5.8) \quad \mathcal{H}^n(\rho x, \overline{B}_\rho) = \rho^{-s_n} \mathcal{H}^n(x, \overline{B}_1) + d\omega_d \frac{\rho^{-s_n} - 1}{s_n} \quad \text{for all } \rho > 0, x \in \partial B_1.$$

For n large enough we have $s_n \geq -\frac{1}{2}$, and hence, for $\rho > 1$, by Lagrange Theorem, there exists $\xi_n \in (-\frac{1}{2}, 0)$ such that

$$(5.9) \quad \frac{\rho^{-s_n} - 1}{s_n} = -\rho^{-\xi_n} \log \rho \geq -\rho^{\frac{1}{2}} \log \rho;$$

therefore, by (5.8), (5.9), Theorem 5.4, and (4.9), we have that there exists a constant $K > 0$ such that

$$(5.10) \quad \underline{c}^{\inf}(\rho) := \inf_{n \in \mathbb{N}} \underline{c}^n(\rho) \geq -K\rho \quad \text{for all } \rho > 1.$$

Moreover, again by (5.8), (5.9), and Theorem 5.4, it is easy to see that there exist two constants $C_1, C_2 > 0$ such that

$$(5.11) \quad \bar{c}^{\sup}(\rho) := \sup_{n \in \mathbb{N}} \bar{c}^n(\rho) \leq C_1(\rho + |\log \rho|) + C_2 \quad \text{for all } \rho > 0.$$

Therefore, by (5.10) and (5.11) we deduce that $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies also property (UB), and again by Theorem 5.4 (in particular, by (5.3)), we get that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ in the sense of Definition 3.1. One can thus apply Theorem 3.2 in order to get the claim. \square

6. THE FLOW GENERATED BY THE REGULARIZED r -MINKOWSKI CONTENT

As a final example, we consider the asymptotic behavior of the flow generated by the regularized r -Minkowski content introduced in [3] in the framework of two-phase image segmentation. This flow has been considered also in [10, 11] (see also [19]) where, in particular, it has been proved existence and uniqueness of the corresponding level set solution.

Let $r > 0$ be fixed. For every measurable set E , we define the r -Minkowski content of E as

$$(6.1) \quad J_r(E) := \frac{1}{2r} \int_{\mathbb{R}^d} \text{osc}_{B_r(x)}(\chi_E) \, dx,$$

where $\text{osc}_A(u) = \text{ess sup}_A u - \text{ess inf}_A u$. One can check (see for instance [10]) that $J_r(E)$ coincides with the measure of the r -neighborhood of the essential boundary of E divided by $2r$; moreover, one can show that, under mild regularity assumptions on E , $J_r(E)$ converges (pointwise and in sense of Γ -convergence) to the standard perimeter. In [10] it has been

proved that (6.1) is a generalized perimeter, and the corresponding curvature, i.e., its first variation with respect to inner variations, has been introduced.

Let $E \in \mathfrak{C}$. For every $x \in \partial E$, we set

$$(6.2) \quad \kappa_r(x, E) = \kappa_r^{\text{ou}}(x, E) + \kappa_r^{\text{in}}(x, E),$$

where

$$\kappa_r^{\text{ou}}(x, E) = \begin{cases} \frac{1}{2r} \det(I + rD\nu_E(x)) & \text{if } d(x + r\nu_E(x), E) = r, \\ 0 & \text{otherwise,} \end{cases}$$

$$\kappa_r^{\text{in}}(x, E) = \begin{cases} -\frac{1}{2r} \det(I - rD\nu_E(x)) & \text{if } d(x - r\nu_E(x), \mathbb{R}^d \setminus E) = r. \\ 0 & \text{otherwise.} \end{cases}$$

In the above formulas, $\nu_F(x)$ denotes the outer normal unit vector to ∂F at x whereas $d(y, F)$ is the distance between the point y and the set F .

The curvature $\kappa_r(x, E)$ is not continuous and it is the true first variation of J_r only for a strict subset of \mathfrak{C} . In order to deal with a well defined curvature for all $E \in \mathfrak{C}$, in [10], a regularization of J_r , by an averaging procedure, has been defined as follows. Fix a function $f : \mathbb{R} \rightarrow [0, +\infty)$ which is even, smooth and nonincreasing in $[0, +\infty)$, with support in $[-1, 1]$ and define $f_r(s) := \frac{1}{2r} f(s/r)$. Set

$$(6.3) \quad J_r^f(E) := \int_{\mathbb{R}^d} f_r(d_E(x)) dx = \int_0^r (-2sf'_r(s)) J_s(E) ds = \int_0^1 (-sf'(s)) J_{rs}(E) ds,$$

where d_E is the signed distance from ∂E and the second equality is obtained by exploiting coarea formula.

For every $E \in \mathfrak{C}$, $x \in \partial E$, the first variation of $J_r^f(E)$ at x is now well defined [10] and given by

$$(6.4) \quad \kappa_r^f(x, E) := \int_0^1 (-sf'(s)) [\kappa_{rs}^{\text{ou}}(x, E) + \kappa_{rs}^{\text{in}}(x, E)] ds.$$

Existence and uniqueness of flows driven by the curvature κ_r^f have been established in [11, Section 6.4] (see also [10]). We briefly recall such results in next theorem.

Theorem 6.1. *Let $f : \mathbb{R} \rightarrow [0, +\infty)$ be even, smooth and nonincreasing in $[0, +\infty)$, with support in $[-1, 1]$. For every $r > 0$, the functionals κ_r^f satisfy the properties (M), (T), (C'), (B) and (S) in Subsection 2.1.*

For every $r > 0$ and for every uniformly continuous function $u_0 \in C(\mathbb{R}^d)$ constant outside a compact set, there exists a unique viscosity solution to (2.4) with \mathcal{H} replaced by κ_r^f .

We now show the convergence of κ_r^f as $r \rightarrow 0$ in the sense of Definition 3.1.

Theorem 6.2. *Let $f : \mathbb{R} \rightarrow [0, +\infty)$ be even, smooth and nonincreasing in $[0, +\infty)$, with support in $[-1, 1]$. Let $\{r_n\}_{n \in \mathbb{N}} \subset (0, 1)$ be such that $r_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $\{E_n\}_{n \in \mathbb{N}} \subset \mathfrak{C}$ be such that $E_n \rightarrow E$ in \mathfrak{C} for some $E \in \mathfrak{C}$. Then, for every $x \in \partial E \cap \partial E_n$ it holds*

$$(6.5) \quad \lim_{n \rightarrow +\infty} \kappa_{r_n}^f(x, E_n) = c_f H^1(x, E),$$

where $H^1(x, E)$ is the scalar mean curvature of the set ∂E at x in (4.25), and $c_f = \int_0^1 f(s) ds$.

Proof. Let r^{in} be the maximal radius r such that E satisfies the interior ball condition with radius r , and let r^{ou} be defined analogously. Clearly, r^{in} and r^{ou} are continuous with respect to smooth inner variations. For n large enough the sets E_n satisfy the interior and exterior ball condition with radius $\bar{r} := \min\{r^{\text{in}}, r^{\text{ou}}\}/2$.

Fix $s \in (0, 1)$; for n large enough $r_n \leq \bar{r}$, and hence

$$\kappa_{sr_n}(x, E_n) = \frac{\det(I + sr_n D\nu_{E_n}(x)) - \det(I - sr_n D\nu_{E_n}(x))}{2sr_n} = \text{tr}D\nu_{E_n}(x) + o(1).$$

Since $E_n \rightarrow E$ in \mathfrak{C} , we conclude that

$$\lim_{r_n \rightarrow 0} \kappa_{sr_n}(x, E_n) = \text{tr}D\nu_E(x) = H^1(x, E),$$

which, by the Dominated Convergence Theorem, implies

$$\lim_{n \rightarrow +\infty} \kappa_{r_n}^f(x, E_n) = H^1(x, E) \int_0^1 (-sf'(s)) ds = H^1(x, E) \int_0^1 f(s) ds.$$

□

Finally, we conclude with the asymptotic result.

Theorem 6.3. *Let $f : \mathbb{R} \rightarrow [0, +\infty)$ be even, smooth and nonincreasing in $[0, +\infty)$, with support in $[-1, 1]$ and let $r_n \rightarrow 0$ as $n \rightarrow +\infty$. Let $u_0 \in C(\mathbb{R}^d)$ be a uniformly continuous function, constant outside a compact set.*

For every $n \in \mathbb{N}$ let u^n be the viscosity solution to (2.4) with \mathcal{H} replaced by $\kappa_{r_n}^f$. Then, $u^n \rightarrow u^\infty$ locally uniformly where $u^\infty : \mathbb{R}^d \times [0, +\infty) \rightarrow \mathbb{R}$ is the (unique) viscosity solution to (2.4) with \mathcal{H} replaced by $c_f H^1$.

Proof. For every $n \in \mathbb{N}$ we set $\mathcal{H}^n := \kappa_{r_n}^f$ and $\mathcal{H}^\infty := H^1$. By Theorem 6.1, \mathcal{H}^n are nonlocal curvatures in the sense of Subsection 2.1 and satisfy (C') and (S). Trivially, also \mathcal{H}^∞ is a nonlocal curvature satisfying (C') and (S).

To check that $\{\mathcal{H}^n\}_{n \in \mathbb{N}}$ satisfies property (UB), we note that

$$(6.6) \quad \kappa_{r_n}^f(\rho x, B_\rho) = \frac{1}{\rho} \kappa_{\frac{r_n}{\rho}}^f(x, B_1) \quad \text{for all } n \in \mathbb{N}, \rho > 0, x \in \partial B_1.$$

Then we compute for $r > 0$,

$$(6.7) \quad \kappa_r^{\text{ou}}(x, B_1) = \frac{1}{2r} (1+r)^d \quad \forall r > 0 \quad \text{and} \quad \kappa_r^{\text{in}}(x, B_1) = \begin{cases} -\frac{1}{2r} (1-r)^d & r < 1 \\ 0 & r \geq 1 \end{cases}.$$

It is immediate to deduce that $\kappa_r(x, B_1) \geq 0$ for all $r > 0$ and then also $\kappa_r^f(x, B_1) \geq 0$ and, in view of (6.6), $\mathcal{H}^n(\rho x, \bar{B}_\rho) \geq 0$ for all $x \in \partial B_1$, $\rho > 0$. Moreover, by (6.7), it follows that

$$(6.8) \quad \begin{aligned} \kappa_r^f(x, B_1) &\leq \int_0^1 (-sf'(s)) \frac{(1+sr)^d}{2sr} ds = \frac{1}{2} \sum_{k=0}^d \binom{d}{k} r^{k-1} \int_0^1 (-f'(s)) s^k ds \\ &\leq C_1^f + C_2^f r^{d-1}, \end{aligned}$$

for some constants $C_1^f, C_2^f > 0$ depending on f . This fact, together with (6.6), implies that \mathcal{H}^n satisfies also the second property of (UB) is satisfied.

Moreover, by Theorem 6.2 (in particular, by (6.5)), we get that $\mathcal{H}^n \rightarrow \mathcal{H}^\infty$ in the sense of Definition 3.1. One can thus apply Theorem 3.2 in order to get the claim. □

REFERENCES

- [1] N. Abatangelo, E. Valdinoci: A notion of nonlocal curvature. *Numer. Funct. Anal. Optim.* **35** (2014), no. 7-9, 793–815.
- [2] L. Ambrosio, G. De Philippis, L. Martinazzi: Gamma-convergence of nonlocal perimeter functionals. *Manuscripta Math.* **134** (2011), no. 3-4, 377–403.
- [3] M. Barchiesi, S. H. Kang, T. M. Le, M. Morini, and M. Ponsiglione: A variational model for infinite perimeter segmentations based on Lipschitz level set functions: denoising while keeping finely oscillatory boundaries. *Multiscale Model. Simul.* **8** (2010), no. 5, 1715–1741.
- [4] G. Barles, P. Cardaliaguet, O. Ley, and R. Monneau: Global existence results and uniqueness for dislocation equations. *SIAM J. Math. Anal.* **40** (2008), no. 1, 44–69.
- [5] G. Bellettini, M. Novaga: Comparison Results between Minimal Barriers and Viscosity Solutions for Geometric Evolutions. *Ann. Scuola Norm. Sup. Pisa Cl. Sci.* **26** (1998), no. 1, 97–131.
- [6] L. Caffarelli, E. Valdinoci: Regularity properties of nonlocal minimal surfaces via limiting arguments. *Adv. Math.* **248** (2013), 843–871.
- [7] A. Cesaroni, S. Dipierro, M. Novaga, E. Valdinoci: Fattening and nonfattening phenomena for planar nonlocal curvature flows. *Math. Ann.* **375** (2019), no. 1-2, 687–736.
- [8] A. Cesaroni, M. Novaga: Second order asymptotics of the fractional perimeter as $s \rightarrow 1$. arXiv preprint 2020, <https://arxiv.org/abs/2001.11415>.
- [9] A. Cesaroni, V. Pagliari: Convergence of nonlocal geometric flows to anisotropic mean curvature motion. arXiv preprint 2018, <https://arxiv.org/abs/1811.01732>.
- [10] A. Chambolle, M. Morini, M. Ponsiglione: A non-local mean curvature flow and its semi-implicit time-discrete approximation. *SIAM J. Math. Anal.* **44** (2012), no. 6, 4048–4077.
- [11] A. Chambolle, M. Morini, M. Ponsiglione: Nonlocal curvature flows. *Arch. Ration. Mech. Anal.* **218** (2015), no. 3, 1263–1329.
- [12] A. Chambolle, M. Novaga, B. Ruffini: Some results on anisotropic fractional mean curvature flows. *Interfaces Free Bound.* **19** (2017), no. 3, 393–415.
- [13] Y.G. Chen, Y. Giga, S. Goto: Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *J. Differ. Geom.* **33** (1991), no. 3, 749–786.
- [14] H. Chen, T. Weth: The Dirichlet problem for the logarithmic Laplacian. *Comm. Partial Differential Equations* **44** (2019), no. 11, 1100–1139.
- [15] F. Da Lio, N. Forcadel, R. Monneau: Convergence of a non-local eikonal equation to anisotropic mean curvature motion. Application to dislocation dynamics. *J. Eur. Math. Soc. (JEMS)* **10** (2008), no. 4, 1061–1104.
- [16] E. De Giorgi: *Barriers, boundaries, motion of manifolds*. Conference held at the Department of Mathematics of Pavia, March 1994.
- [17] L. De Luca, M. Novaga, M. Ponsiglione: The 0-fractional perimeter between fractional perimeters and Riesz potentials. arxiv preprint 2019, <https://arxiv.org/abs/1906.06303>.
- [18] S. Dipierro, A. Figalli, G. Palatucci, E. Valdinoci: Asymptotics of the s -perimeter as $s \searrow 0$. *Discrete Cont. Dynam. Syst.* **33** (2013), no. 7, 2777–2790.
- [19] S. Dipierro, M. Novaga, E. Valdinoci: On a Minkowski geometric flow in the plane: evolution of curves with lack of scale invariance. *J. Lond. Math. Soc.* **99** (2019), no. 1, 31–51.
- [20] S. Dipierro, E. Valdinoci: *Nonlocal minimal surfaces: interior regularity, quantitative estimates and boundary stickiness*. Recent developments in nonlocal theory, 165–209, De Gruyter, Berlin, 2018.
- [21] L.C. Evans, J. Spruck: Motion of level sets by mean curvature I. *J. Differ. Geom.* **33** (1991), no. 3, 635–681.
- [22] C. Imbert: Level set approach for fractional mean curvature flows. *Interfaces Free Bound.* **11** (2009), no. 1, 153–176.
- [23] H. Ishii, P. Souganidis: Generalized motion of noncompact hypersurfaces with velocity having arbitrary growth on the curvature tensor. *Tohoku Math. J. (2)* **47** (1995), no. 2, 227–250.
- [24] V. Julin, D. La Manna: Short time existence of the classical solution to the fractional Mean curvature flow. arXiv preprint 2019, <https://arxiv.org/abs/1906.10990>.
- [25] M. Ludwig: Anisotropic fractional perimeters. *J. Differential Geom.* **96** (2014), no. 1, 77–93.
- [26] C. Mantegazza: *Lecture notes on mean curvature flow*. Progress in Mathematics, 290. Birkhäuser/Springer Basel AG, Basel, 2011.

- [27] V. Maz'ya, T. Shaposhnikova: On the Bourgain, Brezis and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces. *J. Funct. Anal.* **195** (2002), no. 2, 230–238. Erratum: *J. Funct. Anal.* **201** (2003), no. 1, 298–300.
- [28] S. Osher, J.A. Sethian: Fronts propagating with curvature-dependent speed: algorithms based on Hamilton-Jacobi formulations. *J. Comput. Phys.* **79** (1988), no. 1, 12–49.
- [29] E. Valdinoci: A fractional framework for perimeters and phase transitions. *Milan J. Math.* **81** (2013), no. 1, 1–23.
- [30] Y. Zhao, L. Rada, K. Chen, S.P. Harding, Y. Zheng: Automated Vessel Segmentation Using Infinite Perimeter Active Contour Model with Hybrid Region Information with Application to Retinal Images. *IEEE Transactions on Medical Imaging* **34** (2015), no. 9, 1797–1807.

(Annalisa Cesaroni) DIPARTIMENTO DI SCIENZE STATISTICHE, UNIVERSITÀ DI PADOVA, VIA CESARE BATTISTI 241/243, I-35121 PADOVA, ITALY
E-mail address: `annalisa.cesaroni@unipd.it`

(Lucia De Luca) IAC-CNR, VIA DEI TAURINI 19, I-00184 ROMA, ITALY
E-mail address, L. De Luca: `lucia.deluca@cnr.it`

(Matteo Novaga) DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, I-56127 PISA, ITALY
E-mail address, M. Novaga: `matteo.novaga@unipi.it`

(Marcello Ponsiglione) DIPARTIMENTO DI MATEMATICA “GUIDO CASTELNUOVO”, SAPIENZA UNIVERSITÀ DI ROMA, PIAZZALE ALDO MORO 2, I-00185 ROMA, ITALY
E-mail address, M. Ponsiglione: `ponsigli@mat.uniroma1.it`